



*Associate Directorate for Environmental Management*  
 P.O. Box 1663, MS M992  
 Los Alamos, New Mexico 87545  
 (505) 606-2337



*Environmental Management*  
 P. O. Box 1663, MS M984  
 Los Alamos, New Mexico 87545  
 (505) 665-5658/FAX (505) 606-2132

Date: **SEP 29 2017**  
 Refer To: ADEM-17-0246  
 LAUR: 17-28553

John Kieling, Bureau Chief  
 Hazardous Waste Bureau  
 New Mexico Environment Department  
 2905 Rodeo Park Drive East, Building 1  
 Santa Fe, NM 87505-6303

**Subject: Screening-Level Ecological Risk Assessment Methods, Revision 5**

Dear Mr. Kieling:

Enclosed please find two hard copies with electronic files of the Screening-Level Ecological Risk Assessment Methods, Revision 5. Changes to this version include the following:

- An update to the boundary map for Los Alamos National Laboratory (the Laboratory).
- Updates to threatened and endangered species based on the Laboratory's latest Habitat Management Plan.
- Incorporation of an updated vegetation land cover map.
- Updates to wetlands information based on information from the U.S. Army Corps of Engineers.
- Replacement of the desert cottontail with the mountain cottontail and the red fox with the gray fox as receptors.
- Updates to exposure parameter values and references based on recent literature reviews.

If you have any questions, please contact Kent Rich at (505) 665-4272 (krich@lanl.gov) or Arturo Duran at (505) 665-7772 (arturo.duran@em.doe.gov).

Sincerely,

Bruce Robinson, Program Director  
 Environmental Remediation Program  
 Los Alamos National Laboratory

Sincerely,

David S. Rhodes, Director  
 Office of Quality and Regulatory Compliance  
 Environmental Management  
 Los Alamos Field Office

BR/DR/KR:sm

Enclosures: Two hard copies with electronic files – Screening-Level Ecological Risk Assessment Methods, Revision 5 (EP2017-0132)

Cy: (w/enc.)  
Arturo Duran, DOE-EM-LA  
Kent Rich, ADEM ER Program

Cy: (w/electronic enc.)  
Laurie King, EPA Region 6, Dallas, TX  
Steve Yanicak, NMED-DOE-OB, MS M894  
emla.docs@em.doe.gov  
Public Reading Room (EPRR)  
ADESH Records  
PRS Database

Cy: (w/o enc./date-stamped letter emailed)  
lasomailbox@nnsa.doe.gov  
Peter Maggiore, DOE-NA-LA  
Steve Goodrum, DOE-NA-LA  
Arturo Duran, DOE-EM-LA  
David Rhodes, DOE-EM-LA  
Kent Rich, ADEM ER Program  
Bruce Robinson, ADEM ER Program  
Randy Erickson, ADEM  
Jocelyn Buckley, ADESH-EPC-CP  
Mike Saladen, ADESH-EPC-CP  
John Bretzke, ADESH-EPC-DO  
Michael Brandt, ADESH  
William Mairson, PADOPS  
Craig Leasure, PADOPS



LA-UR-17-28553  
September 2017  
EP2017-0132

# **Screening-Level Ecological Risk Assessment Methods, Revision 5**

Prepared by the Environmental Programs Directorate

## EXECUTIVE SUMMARY

This document provides guidance to conduct screening-level ecological assessments at the Los Alamos National Laboratory (LANL or the Laboratory). This guidance promotes consistency, rigor, and defensibility in ecological screening assessments and in reporting the results. The purpose of the screening assessment is to provide information to the risk managers to make informed risk-management decisions. The information presented in this document has been updated to reflect the current approaches implemented since the last revision of the screening-level ecological assessment methods.

The Laboratory-wide information needed for the screening-level ecological risk problem formulation, including the environmental setting, contaminant fate and transport, exposure pathways, and food webs, is presented. Screening assessments are performed on solid waste management units (SWMUs) or areas of concern (AOCs); the area may also be a collection of SWMUs and/or AOCs in a consolidated unit or some other aggregate. In this document, the term *site* is used broadly to include these different possibilities.

The purpose of the screening evaluation is to identify chemicals of potential concern (COPCs) that should be retained as chemicals of potential ecological concern (COPECs). The screening evaluation focuses investigations on important ecological concerns of potentially contaminated sites and identifies those sites that do not have COPECs. Sites with no COPECs do not need further ecological evaluation. The outcome of the screening is protective of receptors from potential adverse ecological effects but is not intended to be predictive of ecological risk. Thus, protective assumptions are made throughout the screening evaluation to ensure contaminants, exposure pathways, and sensitive species are not missed.

The key components of the screening evaluation are the ecological screening levels (ESLs) that are developed for each chemical and receptor and are media-specific. This document presents the basis for, and the elements used in, calculating ESLs for the screening assessment. The ESLs are determined such that if a site has concentrations of a chemical above the ESL in any medium, the site warrants further consideration because the chemical concentration(s) may pose a potential risk to ecological receptors. To evaluate the potential risk for each COPC, the ESL and the site exposure point concentration are used to calculate a hazard quotient (HQ). If the HQ for a COPC at a site with only a single COPC is greater than 1 or the HQ for a COPC is greater than 0.3 for a site with multiple COPCs, then that COPC is identified as a COPEC. The HQs are calculated for each receptor/COPEC combination and are the ratio of a receptor's exposure at the site to an acceptable effects level (i.e., the ESL). Because ESLs are specific to each medium evaluated (soil, sediment, or water), they do not account for exposure to multiple media. The potential hazard posed by multiple chemicals is summed as a hazard index (HI) for each wildlife receptor. If the HI is greater than 1, then the site may pose an ecological risk.

This document also describes the uncertainty analysis that follows the COPEC identification and the key sources of uncertainty in the screening assessment. This analysis includes a more refined screening assessment using ESLs based on the low observed adverse effect level (LOAEL) rather than on the no observed adverse effect level (NOAEL). The LOAEL analysis is less conservative and is designed to provide a more realistic, but still protective, estimate of potential risk. The LOAEL analysis is based on the lowest-effect ecological screening levels (L-ESLs).

The ESL, L-ESL, HQ, and HI calculations require toxicity information, including toxicity reference values (TRVs), and knowledge of bioconcentration and bioaccumulation factors for all chemicals for all receptors and media. The Laboratory's ECORISK Database provides the necessary information and supporting detailed documentation for TRVs, ESLs, L-ESLs, and related information. The database includes values for the TRVs used to develop ESLs and L-ESLs, information on other studies considered for TRVs, transfer and bioaccumulation factors, and exposure parameters for the representative receptor species.



## CONTENTS

<b>1.0</b>	<b>INTRODUCTION .....</b>	<b>1</b>
<b>2.0</b>	<b>GENERIC PROBLEM FORMULATION FOR ECOLOGICAL RISK-SCREENING ASSESSMENTS .....</b>	<b>2</b>
2.1	Environmental Setting .....	2
2.1.1	Geographic Setting.....	2
2.1.2	Climate .....	4
2.1.3	Geology and Soil .....	4
2.1.4	Hydrology .....	6
2.1.5	Biology.....	6
2.1.6	Wetlands .....	9
2.2	Contaminant Fate and Transport.....	10
2.2.1	Hydrologic Transport.....	12
2.2.2	Mass Wasting and Mass Deposition .....	12
2.2.3	Atmospheric Transport.....	13
2.3	Exposure Pathways .....	13
2.4	Functional Food Web.....	14
2.5	Assessment Endpoints .....	17
2.6	Screening Receptors .....	17
<b>3.0</b>	<b>DERIVATION OF ESLs .....</b>	<b>22</b>
3.1	Screening-Level Ecological Risk Calculation Overview .....	25
3.2	Types and Sources of Ecological Effects Information .....	25
3.3	Wildlife Exposure and Effects Evaluations .....	27
3.4	ESLs for Chemicals .....	33
3.4.1	Soil ESLs .....	34
3.4.2	Burrow Air ESLs (Volatile-Phase Contaminants Only) .....	37
3.4.3	Sediment ESLs .....	37
3.4.4	Water ESLs .....	40
3.5	Radionuclide ESLs .....	43
3.5.1	Radionuclide Dose Limits and Effects Calculations.....	44
3.5.2	Soil ESLs.....	44
3.5.3	Sediment ESLs.....	45
3.5.4	Water ESLs .....	46
<b>4.0</b>	<b>SCREENING-LEVEL ECOLOGICAL RISK ASSESSMENT .....</b>	<b>48</b>
4.1	Scoping Evaluation .....	48
4.1.1	Ecological Scoping Checklist .....	49
4.2	Screening Evaluation.....	53
4.3	Uncertainty Analysis .....	53
4.3.1	Development of AUFs .....	55
4.3.2	Exposure-Related Parameters.....	56
4.3.3	Toxicity-Related Parameters .....	58
4.3.4	L-ESL Analysis .....	59
4.3.5	Comparison with Previous Investigations .....	60
4.4	Risk Interpretation.....	60
<b>5.0</b>	<b>REFERENCES .....</b>	<b>60</b>



## Figures

Figure 2.1-1	Laboratory TAs in relation to surrounding landholdings .....	3
Figure 2.1-2	Topography of the Los Alamos area.....	5
Figure 2.2-1	Key transport pathways .....	11
Figure 2.4-1	Terrestrial food web based on feeding relationships of biota on the Pajarito Plateau.....	15
Figure 2.4-2	Aquatic food web based on feeding relationships of biota on the Pajarito Plateau .....	16
Figure 2.6-1	Screening receptors for terrestrial food web .....	23
Figure 2.6-2	Screening receptors for aquatic food web .....	24
Figure 3.4-1	Ecological conceptual site model (CSM) for soil pathways .....	35
Figure 3.4-2	Aquatic CSM for sediment pathways.....	38
Figure 3.4-3	Terrestrial CSM for sediment pathways (to account for bioaccumulation concerns) .....	39
Figure 3.4-4	Aquatic CSM for water pathways.....	41
Figure 3.4-5	Terrestrial CSM for water pathways.....	42
Figure 3.5-1	Aquatic CSM for sediment pathways.....	46
Figure 3.5-2	Aquatic CSM for water pathways.....	47
Figure 4.1-1	Terrestrial receptor conceptual exposure and transport model .....	51
Figure 4.1-2	Aquatic receptor conceptual exposure and transport model .....	52

## Tables

Table 2.1-1	Sensitive Species Occurring or Potentially Occurring at the Laboratory .....	7
Table 2.6-1	List of Receptors Selected for Screening at the Laboratory .....	19
Table 3.3-1	Measures Required for the Wildlife Exposure Model.....	29
Table 3.4-1	Ecological Screening Receptors for Chemicals.....	33

## Appendixes

Appendix A	Derivation of Ecological Screening Levels
Appendix B	Ecological Scoping Checklist

## Plate

Plate 1	Vegetation land cover map of the LANL area
---------	--

## Acronyms

ACR	acute-chronic ratio
ADEM	Associate Directorate for Environmental Management
AOC	area of concern
ARCS	Assessment and Remediation of Contaminated Sediments
asl	above sea level
AUF	area use factor
AWQC	ambient water-quality criteria
BCF	bioconcentration factor
bgs	below ground surface
BMP	best management practice
Bq	Becquerel
BW	body weight
CCC	criterion continuous concentration
CCME	Canadian Council of Ministers of the Environment
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CMC	criterion maximum concentration
COPC	chemical of potential concern
COPEC	chemical of potential ecological concern
CSL	cleanup screening level
CSM	conceptual site model
DCF	dose conversion factor
DOE	Department of Energy (U.S.)
EP	Environmental Programs (Directorate)
EPA	Environmental Protection Agency (U.S.)
EPC	exposure point concentration
EqP	equilibrium partitioning
ERM	effect range median
ESA	Endangered Species Act
ESH	Environment, Health, and Safety (Directorate)
ESL	ecological screening level
FSOC	Federal Species of Concern
FSS	Forest Service Sensitive Species
HI	hazard index
HMP	Habitat Management Plan
HQ	hazard quotient
HR	home range
IR	inhalation rate

IAEA	International Atomic Energy Agency
LANL	Los Alamos National Laboratory
L-ESL	lowest effect ESL
LOAEL	lowest observed adverse effect level
LOEC	lowest observed effect concentration
MC	moisture content
MeV	million electron volts
MPC	maximum permissible concentration
NC	negligible concentration
NCRP	National Council on Radiation Protection
NMAC	New Mexico Administrative Code
NME	New Mexico Endangered
NMED	New Mexico Environment Department
NMS	New Mexico Sensitive Taxa
NMSOC	New Mexico Species of Concern
NMT	New Mexico Threatened
NOAA	National Oceanic and Atmospheric Administration
NOAEL	no observed adverse effect level
NOEC	no observed effect concentration
PAUF	population area use factor
PEC	probable effect concentration
PTSE	primary toxicity study evaluation
PTV	primary toxicity value
RPF	Records Processing Facility
S1	Heritage New Mexico: Critically Imperiled in New Mexico
SEL	severe effect level
SI	International System of Units
SLERA	screening-level ecological risk assessment
SMDP	scientific management decision point
SQS	sediment quality standard
SQuiRT	Screening Quick Reference Tables
Sv	sievert
SWMU	solid waste management unit
T&E	threatened and endangered
TA	technical area
TEC	threshold effect concentration
TEL	threshold effect level
TF	transfer factor

TL	threshold level
TRV	toxicity reference value
UCL	upper confidence limit
UET	upper effects threshold
UF	uncertainty factor
USACOE	U.S. Army Corps of Engineers
VOC	volatile organic compound
WQC	water-quality criteria





## **1.0 INTRODUCTION**

This revised methodology document describes the approach used by the Los Alamos National Laboratory (LANL or the Laboratory) Environmental Programs Directorate for screening-level assessments of potential impacts to ecological resources resulting from exposure to contaminants. The approach is consistent with the U.S. Environmental Protection Agency's (EPA's) "Ecological Risk Assessment Guidance for Superfund" (EPA 1997, 059370); the "Guidelines for Ecological Risk Assessment" (EPA 1998, 062809); "Issuance of Final Guidance: Ecological Risk Assessment and Risk Management Principles for Superfund Sites" (EPA 1999, 070086); and the "Guidance for Developing Ecological Soil Screening Levels" (EPA 2003, 085643). This guidance incorporates the assessment endpoints developed in "Generic Assessment Endpoints for Ecological Risk Assessment at the Los Alamos National Laboratory" (LANL 1999, 064137). The guidance in this document is consistent with the New Mexico Environment Department's (NMED's) "Risk Assessment Guidance for Site Investigations and Remediation" (NMED 2017, 602274; NMED 2017, 602273). The approach to ecological risk screening for radionuclides provided in this document is also consistent with the U.S. Department of Energy's (DOE's) "Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota" (DOE 2002, 085637) and DOE's "RESRAD-BIOTA: A Tool for Implementing a Graded Approach to Biota Dose Evaluation, User's Guide, Version 1" (DOE 2004, 085639). This version of the document incorporates guidance on and direction for conducting ecological risk-screening assessments and is consistent with risk-assessment conducted by the Laboratory.

EPA guidance requires that initial screening-level assessments use conservative assumptions to evaluate the potential for adverse ecological impacts. The rationale behind this requirement is to provide high confidence that all potential adverse impacts to ecological receptors resulting from exposure to contaminants are identified in the initial investigation. Thus, the screening-level assessment may be used to identify sites that clearly pose no threat to the environment as well as sites that need corrective action. However, for the many sites that do not fall into one of these two categories, screening-level evaluations must be followed by a series of progressively more in-depth and site-specific evaluations to characterize risks accurately and to provide adequate information for risk-management decisions. The screening-level assessment helps to focus these more detailed (and often more complex) site-specific investigations by identifying important contaminants, receptors, ecological endpoints, and spatial scales. The screening-level evaluation also employs a common metric for comparing risks among different sites, thus providing a tool for prioritizing site investigations and corrective actions.

This document presents the ecological screening process for individual solid waste management units (SWMUs) or areas of concern (AOCs) as well as clusters of SWMUs and/or AOCs. Application of this methodology to larger spatial aggregates is not explicitly considered. The approach assesses present-day risk at the site where contamination has been investigated and characterized. However, these methods, coupled with the appropriate transport models, may be used to assess the potential for future ecological risk at areas affected by off-site transport of contaminants. The discussion and evaluation of transport models, other than to emphasize their importance, is beyond the scope of this document.

## **2.0 GENERIC PROBLEM FORMULATION FOR ECOLOGICAL RISK-SCREENING ASSESSMENTS**

As noted in the Comprehensive Environmental Response, Compensation, and Liability Act– (CERCLA- or Superfund-) specific ecological risk guidance (EPA 1997, 059370), problem formulation is the most critical step of an ecological risk assessment. The EPA guidance identifies (among others) the following issues for the screening-level problem formulation:

- Environmental setting (physical and biological)
- Contaminant fate and transport
- Food webs
- Screening receptors
- Exposure pathways
- Assessment endpoints

Therefore, problem formulation requires understanding the physical and biological setting of the Laboratory. The physical setting greatly influences the potential contaminant transport pathways, which also influence the potential exposure pathways for ecological receptors. The biological setting is important for receptor selection because receptors must represent the broad spectrum of plant and animal species present at the Laboratory. One key exposure pathway is expressed through the food web (section 2.4), which structures information on the feeding relationships among animals and plants to develop representative groups of ecological receptors. Receptor groupings based on feeding relationships are an efficient and effective way to represent all relevant biota. In the following sections, the general physical setting of the Laboratory and the surrounding area is summarized, followed by descriptions of the salient biotic features.


### **2.1 Environmental Setting**

The Laboratory is situated on the Pajarito Plateau, which consists of a series of fingerlike mesas separated by deep east-to-west-oriented canyons cut by intermittent streams. Mesa tops range in elevation from approximately 2377 m (7800 ft) on the flanks of the Jemez Mountains to about 1890 m (6200 ft) at their eastern termination above the Rio Grande. The climate, geographic setting, geology, hydrology, and biology of the Laboratory are described briefly below.

#### **2.1.1 Geographic Setting**

The Laboratory and residential and commercial areas of Los Alamos and White Rock are located in Los Alamos County, in north-central New Mexico, approximately 60 mi northeast of Albuquerque and 20 mi northwest of Santa Fe. The surrounding land is largely undeveloped, with large tracts of land north, west, and south of the Laboratory held by the Santa Fe National Forest, Bureau of Land Management, Bandelier National Monument, General Services Administration, and Los Alamos County. The Pueblo of San Ildefonso borders the Laboratory to the east.

The Laboratory is divided into technical areas (TAs) that are used for building sites, experimental areas, waste disposal locations, roads, and utility rights-of-way (Figure 2.1-1). However, these uses account for only a small part of the total land area. Most land provides buffer areas for security and safety and is held in reserve for future use. Thus, much of the Laboratory is undeveloped land that supports diverse and abundant organisms.



**Figure 2.1-1 Laboratory TAs in relation to surrounding landholdings**

### **2.1.2 Climate**

The average diurnal temperature at Los Alamos is 13°C (55°F). Winter temperatures range from –1°C to 10°C (30°F to 50°F) during the day, to –9°C to –4°C (15°F to 25°F) during the night. Summer temperatures range from 21°C to 31°C (70°F to 88°F) during the day to 10°C to 15°C (50°F to 59°F) during the night. The average annual precipitation (including both rain and water equivalent of frozen precipitation) is 48 cm (19 in.). Details are available at <http://weather.lanl.gov/> and are discussed in the “Installation Work Plan for Environmental Restoration Project, Revision 8” (LANL 2000, 066802, p. 2-41).

The semiarid, temperate, mountain climate in Los Alamos County influences weather and soil development as well as biotic assimilation in the region. Both weather and soil conditions influence transport of contaminants at the Laboratory and potential exposure of ecological receptors to contamination. The speed, frequency, direction, and persistence of wind influence the airborne transport of contaminants. High winds, common in the spring, can result in atmospheric transport of contaminants. The role of climate in the atmospheric contaminant pathway is considered part of the site-specific scoping evaluation.

Intense thunderstorms in the summer can cause erosion of unstable sediment or soil. The form, frequency, intensity, and evaporation potential of precipitation strongly influence surface water runoff and infiltration of contaminants. As discussed below, fires also change hydrological regimes, and small precipitation events may lead to large amounts of runoff.

### **2.1.3 Geology and Soil**

Geologic and hydrologic information provides the basis for the discussion of hydrologic transport of contaminants. The likelihood of hydrologic transport is considered in the site-specific scoping evaluation (section 4.1). The geologic and hydrologic characteristics in and around the Laboratory as they relate to the potential for contaminant transport are complex. Additional literature on the hydrology and geology of the Los Alamos region may be found in an annotated bibliography of geologic, hydrogeologic, and environmental studies related to SWMUs and AOCs at the Laboratory (LANL 1990, 047588).

#### **Geology**

The Laboratory extends over the east-sloping dissected tableland of the Pajarito Plateau and is bounded on the west by the eastern Jemez Mountains and on the east by White Rock Canyon of the Rio Grande. The geology of the Pajarito Plateau primarily reflects ancient volcanism in the Jemez Mountains and surrounding areas. The Rio Grande rift lies to the east of the plateau, forming a series of north-south-trending fault troughs from southern Colorado to southern New Mexico. Most of the fingerlike mesas in the Los Alamos area (Figure 2.1-2) are formed in Bandelier Tuff, which includes ash fall, ash-fall pumice, and rhyolite tuff. The tuff is more than 305 m (1000 ft) thick in the western part of the plateau and thins to about 79 m (260 ft) eastward above the Rio Grande. It was deposited as a result of major eruptions in the Jemez Mountains’ volcanic center about 1.2 to 1.6 million years ago. Deep canyons are incised into the Bandelier Tuff and exposed to depths of up to several hundred feet below the upper elevation of the plateau. Some of the deeper canyons expose older lava deposits and sedimentary rocks. Permeable units in the floors that outcrop below saturated alluvium create the potential for recharge to deeper groundwater zones and form a source for springs and seeps in the area. Faults, cooling joints, and fractures potentially occur throughout the Pajarito Plateau (LANL 2000, 066802, pp. 2-23–2-24).





**Figure 2.1-2 Topography of the Los Alamos area**

On the western part of the Pajarito Plateau, the Bandelier Tuff overlaps onto the Tschicoma Formation, which consists of older volcanic rock that comprises most of the Jemez Mountains. The conglomerate of the Puye Formation in the central plateau and near the Rio Grande underlies the tuff. Chino Mesa basalts intertwine with the conglomerate along the river. These formations overlay the sediment of the Santa Fe Group, which extend across the Rio Grande Valley and are more than 1006 m (3300 ft) thick. Most Laboratory facilities are located on tuff, covered by thin discontinuous soil on mesa tops and alluvial deposits of variable thickness on canyon floors.

## **Soil**

Soil erodability is important to understanding the potential for contaminant transport. Characterizing soil erodability is also important for accurately completing the “contaminant transport information” in site-specific scoping evaluations (section 4.1). Soil on the Pajarito Plateau was initially mapped and described by Nyhan et al. (1978, 005702). A large variety of soil and sediment has developed on the Pajarito Plateau as the result of interactions of the underlying bedrock, slope, biota, and climate. Mesa tops may consist of soil derived from Bandelier Tuff, lavas, basalts, sedimentary rocks, and alluvium. Canyon floors generally contain poorly developed, deep, well-drained soil (Nyhan et al. 1978, 005702). General patterns of soil erosion rates are summarized by the following text from section 2.2.1.6 of the installation work plan (LANL 2000, 066802, p. 2-25):

Erosion rates vary considerably on the mesa tops; the highest rates occur in and near drainage channels and in areas of locally steeper slope gradient. The lowest rates occur



on relatively gently sloping portions of the mesa tops removed from channels. Areas where runoff is concentrated by roads and other development are especially prone to accelerated erosion. The rates and processes of erosion may differ significantly between the north and south slopes of the canyons. Given current vegetation and climate, the more extensive exposure of bedrock on south-facing sides and greater soil cover on north-facing sides suggest that erosion rates of fine-grained material that can be transported by runoff are higher on the drier, less-vegetated, south-facing sides of canyons, although this material is largely retained on the north-facing slopes.

The mesa tops generally consist of finer-textured soil, and the canyon bottoms consist of relatively coarse sediment. The finer-textured soil of mesa tops is prone to overland runoff, whereas soil fines may accumulate in canyon bottoms. The latter are subject to mobilization during flood events.

#### **2.1.4 Hydrology**

Surface water on the Pajarito Plateau occurs as streams that are ephemeral (flowing in response to precipitation), intermittent (flowing in response to availability of snowmelt or groundwater discharge), perennial (flowing continuously), or interrupted (alternating perennial, ephemeral, and intermittent reaches). Some surface water arises from natural flows that originate in canyon heads in the upper Jemez Mountains north and west of the Laboratory. Other surface water originates from mesa-top storm water drainage and permitted Laboratory discharges. Perennial springs on the flanks of the Jemez Mountains supply base flow into the upper reaches of some canyons, but the volume is insufficient to maintain surface flows across the Laboratory site before they are depleted by the processes of evaporation, transpiration, and infiltration described in the “Core Document for Canyons Investigations” (LANL 1997, 055622).

The Rio Grande is the highest-order river in north central New Mexico. Much of the surface water flow and groundwater discharge from the Pajarito Plateau canyon systems ultimately arrive at the Rio Grande through drainages that extend from the Laboratory in a southwest direction but not as continuous flow. Only five of the canyons within Laboratory boundaries contain reaches with perennial water flow. These canyons are Los Alamos Canyon, Pajarito Canyon, Water Canyon, Ancho Canyon, and Chaquehui Canyon. In addition to these limited natural perennial reaches, several effluent-supported reaches also exist within the watersheds (LANL 2000, 066802).

Groundwater in the Los Alamos area occurs in three forms: (1) water in shallow alluvium in canyons, (2) perched water (a body of groundwater above a less permeable layer separated from the underlying regional aquifer by an unsaturated zone), and (3) the regional aquifer of the Los Alamos area. Groundwater hydrology for this region, including the potential for contamination, is complex.

#### **2.1.5 Biology**

The biota within the Laboratory includes approximately 500 plant species, 29 mammal species, 200 bird species, 19 reptile species, 8 amphibian species, and 1000s of insect species (LANL 2000, 066802).

The Laboratory’s Habitat Management Plan (HMP) is a document prepared to provide for the protection of federally listed threatened and endangered species and their habitats within the Laboratory boundaries. The HMP is designed to be a comprehensive landscape-scale management plan that balances the current operations and future development needs of the Laboratory with the habitat requirements of threatened and endangered species. It also facilitates DOE compliance with the Endangered Species Act (ESA) and related federal regulations (LANL 2015, 602156).

Currently, the HMP covers the following threatened and endangered species occurring or potentially occurring within the Laboratory boundaries: southwestern willow flycatcher (*Empidonax trailii extimus*), Jemez Mountains salamander (*Plethodon neomexicanus*), Mexican spotted owl (*Strix occidentalis lucida*), the western distinct population segment of the yellow-billed cuckoo (*Coccyzus americanus*), and the New Mexico meadow jumping mouse (*Zapus hudsonius luteus*).

Results of preliminary risk assessments for the bald eagle (*Haliaeetus leucocephalus*), Mexican spotted owl, and southwestern willow flycatcher are available in Gallegos et al. (1997, 057915); Gonzales et al. (1997, 062879); Gonzales (1998, 062349); Gonzales (1998, 062350); and Gonzales et al. (2004, 085207). Information on the biology and ecology of these species relevant to risk from contaminants can also be found in these references.

There are also state-listed or other sensitive species within Laboratory boundaries that are not protected under the ESA. These species are identified by federal or state agencies or non-governmental organizations. Table 2.1-1 presents the current list of sensitive species potentially occurring within Laboratory boundaries (Hathcock et al. 2015, 602499).

**Table 2.1-1**  
**Sensitive Species Occurring or Potentially Occurring at the Laboratory**

Scientific Name	Common Name	Protected Status <sup>a</sup>	Potential to Occur <sup>b</sup>
<i>Gila pandora</i>	Rio Grande chub	NMS	Moderate
<i>Falco peregrinus anatum</i>	American peregrine Falcon	NMT, FSOC	High
<i>Falco peregrinus tundrius</i>	Arctic peregrine Falcon	NMT, FSOC	Moderate
<i>Haliaeetus leucocephalus</i>	Bald eagle	NMT, S1	High
<i>Cynanthus latirostris magicus</i>	Broad-billed hummingbird	NMT	Low
<i>Accipiter gentilis</i>	Northern goshawk	NMS, FSOC	High
<i>Lanius ludovicianus</i>	Loggerhead shrike	NMS	High
<i>Vireo vicinior</i>	Gray vireo	NMT	Moderate
<i>Amazilia violiceps</i>	Violet-crowned hummingbird	NMT	Low
<i>Myotis ciliolabrum melanorhinus</i>	Western small-footed Myotis bat	NMS	High
<i>Myotis volans interior</i>	Long-legged Myotis bat	NMS	High
<i>Euderma maculatum</i>	Spotted bat	NMT	High
<i>Corynorhinus townsendii pallescens</i>	Pale Townsend's big-eared bat	NMS, FSOC	High
<i>Nyctinomops macrotis</i>	Big free-tailed bat	NMS	High
<i>Bassariscus astutus</i>	Ringtail	NMS	High
<i>Vulpes vulpes</i>	Red fox	NMS	Moderate
<i>Ochotona princeps nigrescens</i>	Goat Peak pika	NMS, FSOC	Low
<i>Lilium philadelphicum var. andinum</i>	Wood lily	FSOC, NME	High
<i>Cypripedium parviflorum var. pubescens</i>	Greater Yellow Lady's Slipper	FSOC, NME	Moderate
<i>Speyeria nokomis nitocris</i>	New Mexico silverspot butterfly	FSOC	Moderate
<i>Mentzelia springeri</i>	Springer's blazing star	NMSOC, FSOC, FSS	Moderate

<sup>a</sup> NMS = New Mexico Sensitive Taxa (informal); S1 = Heritage New Mexico: Critically Imperiled in New Mexico; NMT = New Mexico Threatened; NME = New Mexico Endangered; FSOC = Federal Species of Concern (no longer maintained); FSS = Forest Service Sensitive Species; NMSOC = New Mexico Species of Concern.

<sup>b</sup> Low = No known habitat exists on Laboratory property; Moderate = Habitat exists, although the species has not been recorded recently; High = Habitat exists and the species is recorded to occur on Laboratory property.

Knowledge of the vegetative communities at the Laboratory and the animal fauna found in association with these complexes is used in the ecological risk-screening process for predicting the presence of species at the site or in the surrounding areas. For example, areas containing mature, mixed conifer stands are important to Mexican spotted owls. Knowledge and expectations from biological assessments associated with the site are then used to identify potential pathways and exposures to ecological receptors, including threatened and endangered (T&E) species.

The Laboratory has developed a vegetation land cover map (Plate 1) to support region-wide environmental studies. The land cover map identifies areas by the dominant overstory vegetation. The resulting cover types include major vegetation zones and physiognomic types important to the distribution and abundance of several T&E species (McKown et al. 2003, 087150). The individual conducting the site scoping verifies the vegetation cover type during the site visit that supports the site-specific problem scoping.

The land cover types can be subdivided to correspond with the National Vegetation Classification System (McKown et al. 2003, 087150). The elevation and climatic gradients in the region of the Laboratory most strongly influence distribution of three vegetative cover types defined by their dominant tree species and by their structural characteristics; these include piñon-juniper woodlands, ponderosa pine forests, and mixed conifer-spruce-fir forests. In contrast, aspen-riparian-wetland areas, grass species areas, shrub species areas, open water, and urban-sparse-bare rock lands are influenced less by elevation and climatic gradients. Instead, their distribution is most strongly influenced by topographic features, soil and geologic conditions, and moisture levels.

**Mixed conifer-spruce-fir forests.** Mixed conifer forests may be found above 2070 m (6900 ft) above sea level (asl), blended with ponderosa pine communities, but they also extend to lower elevations on north-facing slopes of canyons. These communities continue to the highest elevations of the Sierra de los Valles, 3150 m (10,500 ft). Douglas fir (*Pseudotsuga menziesii*) and white fir (*Abies concolor*) are the typical overstory dominants in mixed conifer forests. At elevations above 2700 m (9000 ft), Engelmann spruce (*Picea engelmannii*) becomes more important. Ponderosa pine (*Pinus ponderosa*) and aspen (*Populus tremuloides*) are also typically present. Limber pine (*Pinus flexilis*) can also be found in mixed conifer forests, especially on rocky ridgelines.

**Aspen-riparian-wetland.** Aspen (*Populus tremuloides*) communities are common at mid-elevations in the mountains, from approximately 2700 m to 3030 m asl (8900 ft to 9950 ft asl). Below 2820 m (9250 ft), aspen stands occupy north and northeast facing slopes, whereas above this elevation they are found mostly on southeast- to southwest-facing slopes. At higher elevations and on south-facing slopes, aspen typically exceeds 45% coverage and may be the only species present in the overstory. At lower elevations and on north-facing slopes, white fir, Engelmann spruce, and Douglas fir may collectively contribute up to 30% of the overstory coverage. Depending on the fire history of the specific stand, other tree species, such as ponderosa pine and limber pine, may be blended with aspen. Riparian areas and wetlands are also included in this vegetation land cover type.

**Grass species.** Grass species areas are dominated by grasses, narrow-leaf plants (e.g., yucca), and colonizing species that invade disturbed areas. Forbs and other nonshrubby species may be dominant components of these communities. Shrubs and trees are absent or rare. The grass species cover type

may include areas undergoing post-fire succession, abandoned homestead areas, montane meadows, and subalpine grasslands.

**Shrub species.** These areas include evergreen, microphyllus shrubs, and temperate, cold-deciduous shrub species. Post-fire shrub-sized sprouts of aspen, Gambel oak (*Quercus gambelii*), and New Mexico locust (*Robinia neomexicana*) are also included in this vegetation type.

**Ponderosa pine.** This vegetation consists of open-canopied woodlands with needle-leaved evergreen trees, primarily ponderosa pine. An understory of Gambel oak or grasses and bare ground may occur between the trees.

**Piñon-juniper.** This vegetation cover also consists of open-canopied woodlands with needle-leaved evergreen trees, primarily piñon pines (*Pinus edulis*) and one-seed junipers (*Juniperus monosperma*); bare soil may be under the trees or an understory of Basin big sage (*Artemisia tridentata*), and blue grama grass (*Bouteloua gracilis*) may grow.

**Open water.** This cover type includes all land that is at least periodically flooded or is open water. In the wettest of these sites, the vegetative cover is limited to plant species that require or prefer permanent or seasonally mesic conditions. The Rio Grande borders the Laboratory on its eastern boundary.

**Urban-sparse-bare rock.** This land type includes all undeveloped land covered by less than 7% vegetation. These land surfaces are dominated by cobbles, boulders, bedrock, or bare ground, including tuffaceous cliffs, basalt cliffs, felsenmeers, and basalt talus. Areas of sparse vegetation resulting from development, such as the Los Alamos townsite, the town of White Rock, and some TAs, are also part of the vegetation land cover class.

#### 2.1.6 Wetlands

Wetlands are generally defined as areas of the environment containing water or moisture that support a host of aquatic plants and animals. More specifically, wetlands are defined on the basis of properties related to hydrophytes and hydrophilic plants, hydric soil, and the hydrology as described in 10 Code of Federal Regulations 1022, "Compliance with Floodplain and Wetland Environmental Review Requirements." In and around the Laboratory, these systems occur primarily in the canyon bottoms of the Pajarito Plateau and along the banks of the Rio Grande. Wetlands may also be associated with effluent and storm water outfalls from Laboratory and county facilities. The Los Alamos Site Office, in March 2005, requested the services of and entered into contract with the U.S. Army Corps of Engineers (USACOE) Albuquerque District for the purpose of identifying and delineating wetlands at the Laboratory. Thirty wetlands occupying portions of 14 different TAs met the criteria of the 1987 USACOE Wetlands Delineation Manual, Routine Method and were identified and delineated totaling 33.955 acres (USACE 2005, 092220). The approximate locations of many of the larger wetlands are shown on Plate 1. Some of the larger wetlands on the Laboratory are located in upper Sandia Canyon, Pajarito Canyon, Mortandad Canyon, and Cañon de Valle.

The protection of wetland ecosystems at the Laboratory from the impacts of contaminants is especially important because of the diversity of associated fauna and because wetlands provide significant potential contaminant uptake pathways. These pathways include food web, direct media contact, and gamma radiation exposure pathways. Additionally, aquatic organisms occupying wetlands may experience higher exposures to contaminants because of continuous contact with water and specialized respiration mechanisms. Wetlands are of critical importance to both terrestrial and aquatic biota. Functional aspects of wetlands include food web contribution, breeding habitat, sediment retention, erosion prevention, flood and runoff storage, groundwater recharge, and nutrient retention.


## **2.2 Contaminant Fate and Transport**

The geomorphology of the Pajarito Plateau, with its alternating mesas and canyons, determines the primary contaminant transport pathways for sources of environmental contamination. Figure 2.2-1 shows the key transport pathways:

- hydrologic transport (e.g., surface water and groundwater)
- physical transport (e.g., mass wasting of cliffs) and
- atmospheric transport (e.g., dust resuspension)

These pathways are discussed briefly below. Pathways relevant to a particular site should be discussed in the applicable, site-specific reports.





**Figure 2.2-1 Key transport pathways**

Source: F4-1, CORE DOC, 032497  
modified for F3.2-1, SLERA, Rev.2, 122004, ptm; modified 101512, ptm

## **2.2.1 Hydrologic Transport**

### **2.2.1.1 Surface Water and Sediment Transport**

Surface water flows provide the primary mechanism for redistributing and transporting the contaminants that remain from early Laboratory operations. The primary mechanisms affecting mobilization of contaminants within the canyons include sediment transport, contaminant dissolution and desorption, runoff, infiltration, and percolation. The water flowing across Laboratory property, especially in canyon systems, is used by wildlife, constituting a major potential contaminant exposure pathway to these receptors. Much of the surface water flow, including groundwater discharge from springs, from the Pajarito Plateau ultimately arrives at the Rio Grande.

Sediment transport by surface water may be the predominant mechanism for redistributing contaminants at the Laboratory. Carried by storm-event runoff, contamination from mesa-top release sites could enter surface-water drainages. Contaminants have also been released directly into stream channels by effluent discharges. Most environmental contaminants are adsorbed onto sediment particles, preferentially binding to particles with high surface areas and/or charged particles, such as silt and clay. The more soluble contaminants may remain in solution, which makes them available for vertical transport to perched aquifers and for later emergence in springs.

Erosion of soil and transport as sediment via surface water by runoff has significantly increased in areas of the Laboratory within or downstream of the Cerro Grande (2000) and Las Conchas (2011) burn scars. In addition to an increase in the mass of sediment transported in the 2 to 3 yr following the fires, the concentrations of both nonradionuclides and radionuclides in sediment also increased significantly (e.g., Kraig et al. 2002, 085536; Gallaher and Koch 2004, 088747). The sediment is transported downstream and deposited at some locations where these elevated concentrations are potentially available to both terrestrial and aquatic receptors. Increased flow also leads to erosion of sediment deposits in other settings and contaminants in the mobilized sediment would mix with post-fire material and other upstream sediment sources.

### **2.2.1.2 Groundwater Transport**

The primary mechanism for contaminant transfer between the surface and underlying groundwater is infiltration of surface water carrying colloidal and dissolved contaminants (LANL 1997, 055622). The potential for significant infiltration from mesa-top settings is typically limited by the general lack of ponded water that might create hydraulic head. In canyon settings, however, the potential for significant infiltration exists, given the presence of perennial or intermittent surface water and coarse-grained sediment in most parts of the canyon systems and the high vertical hydraulic gradients beneath canyon streams.

Saturated groundwater zones beneath the Pajarito Plateau may be recharged in part by the vertical migration of water from canyon-floor alluvium, which may be partly directed and accelerated by faults and fractures. Unsaturated zones are considered only an occasional transport pathway.

## **2.2.2 Mass Wasting and Mass Deposition**

Physical transport of surface or subsurface materials is most dramatically possible through a mechanism termed *mass wasting*. Mass wasting is the process in which blocks of soil and rock break off the cliffs and are deposited violently into the canyons. Mass wasting is an episodic phenomenon and could be an important mechanism of contaminant transport for mesa-top sites located near canyon walls. Exposure to ecological receptors would result if subsurface contamination became surficial contamination through

mass wasting into the canyons. The transport pathways would then be similar to media subject to surface-water transport.

### **2.2.3 Atmospheric Transport**

Atmospheric transport may occur through transport of windblown particles or vaporization of volatile chemicals. Transport of soil or fine sediment particles by wind is a means of dispersing contaminants. Wind resuspension and transport of surficial contaminant-laden soil or sediment is not a significant transport pathway because the volume of contaminated media mobilized by this pathway is small compared to the total amount of soil to which the receptor is exposed. Exposure of surface-dwelling animals to vapors does not represent a significant pathway because vapors disperse in the open atmosphere. Within burrows, however, vapors from subsurface contamination may accumulate and result in potentially significant exposures to animals.

## **2.3 Exposure Pathways**

Contaminants associated with surface soil may be available to biological receptors through the following exposure pathways:

- Rain splash or saltation-creep of contaminated soil onto plants
- Root uptake of water-soluble contaminants
- Incidental ingestion of soil
- Dermal contact with soil
- Inhalation of volatile chemicals by animals in burrows
- Deposition of particulates on foliage
- Deposition of particulates on animals, and subsequent ingestion during grooming
- Food web transport (consumption of contaminated plants and animals)
- Direct exposure to soil containing gamma-emitting radioactive contaminants

Contaminants associated with sediment or surface water may be taken up by biota primarily through the following exposure pathways:

- Ingestion of surface water
- Root uptake of surface water
- Root uptake of water-soluble contaminants from sediment
- Incidental ingestion of sediment
- Rain splash or saltation-creep of contaminated sediment onto plants
- Dermal contact with surface water or sediment
- Exposure to aquatic animals through respiration
- Food web transport (consumption of contaminated plants and animals)

- Direct exposure to sediment containing gamma-emitting radioactive contaminants
- Direct exposure to surface water containing gamma-emitting radioactive contaminants (immersion)

When groundwater becomes surface water in springs or seeps, the previous exposure pathways also apply. In addition, shallow groundwater, particularly alluvial water, may be taken up by deep-rooted plants (e.g., chamisa) and enter the food web primarily through the ingestion of contaminated plants.

Contaminants present in air as vapors are available for uptake by biota through the following exposure pathways:

- Inhalation of contaminants present as vapors by animals in burrows
- Uptake by plants of contaminants present as vapors

## 2.4 Functional Food Web


A food web diagram is important for evaluating dietary exposure pathways and for specifying ecologically relevant groups of organisms for an exposure assessment. The food web structure captures functionally relevant biotic assimilation and associated relationships and is important for selecting receptors that may be vulnerable to contaminants by virtue of dietary exposure. A food web diagram also shows pathways of food consumption in a biotic system by means of boxes and connecting arrows. Boxes in a food web diagram represent biota (e.g., functional assemblages or taxonomic groups), and arrows define the major direction of energy flow between biota (e.g., from prey to predators).

For the purposes of this ecological screening-level risk-assessment methodology, it is more useful to design a food web where biological receptors are classified into functional groups with similar feeding roles instead of a taxonomic classification. Taxonomically based food webs use phylogenetic classification to organize species into evolutionarily related natural assemblages (genera, families, orders) and are not sensitive to potentially similar feeding habits among taxa. Figures 2.4-1 and 2.4-2 represent the terrestrial and aquatic functional food webs for the Laboratory, respectively. The food webs are organized into functional guilds based on feeding (trophic) relationships. Thus, a *feeding guild* is a collection of species sharing common food consumption roles. For example, animals that eat seeds (granivores) are considered one feeding guild, browsers/grazers another, and top carnivores yet another. Feeding guilds may be organized in many ways, from general to specific.


A food web organized by feeding guilds forms a basis for selecting individual species from each guild that represent the guild as a whole. This approach forms the basis of receptor selection for the ecological screening assessments at the Laboratory. The food webs for the Laboratory include three fundamental trophic positions: producers (vascular and nonvascular plants); consumers (herbivores, omnivores, carnivores, and parasites); and decomposers. Within these basic trophic levels, several feeding guilds have been identified. For example, one group of consumers is herbivores, consisting of six feeding guilds: seed eaters (granivores), fruit eaters (frugivores), foliage or leaf eaters (folivores), nectar and pollen feeders (nectarivores/pollen eaters), fungi eaters (fungivores), and browser/grazers. Since the Laboratory food web included multiple levels of organization, it was necessary to choose receptors that were broadly representative of these levels.

Consumers

Carnivores




Parasites




Omnivores

Herbivores





Parasites

Producers

Vascular



Non-vascular

Decomposers

NUTRIENTS

Figure 2.4-1 Terrestrial food web based on feeding relationships of biota on the Pajarito Plateau




Figure 2.4-2 Aquatic food web based on feeding relationships of biota on the Pajarito Plateau

Aquatic environments on the Laboratory are of limited spatial extent and typically occur in canyon settings. Therefore, the primary connection between the terrestrial and aquatic food webs is not riparian species but rather aerial insectivores for which receptors are designated as part of the terrestrial food web discussed in section 2.6. Separate screening receptors are developed for the terrestrial and aquatic food webs described in section 2.6 because of the limited connectivity between the aquatic and terrestrial systems at the Laboratory. Vertebrate herbivores, omnivores, and carnivores are listed on the aquatic food web to represent the trophic positions of fish species. The dashed lines in Figure 2.4-1, enclosing a number of guilds in a single rectangle, represent broad categories for which a single member may suffice as a screening receptor.

## 2.5 Assessment Endpoints

To represent the feeding guilds in the food webs as described in section 2.4, some attribute of that receptor must be selected as an assessment endpoint, an explicit expression of the environmental value to be protected. These endpoints should be ecologically relevant and should help sustain the natural structure, function, and biodiversity of an ecosystem or its components (EPA 1998, 062809). In a screening-level assessment, assessment endpoints are any adverse effects on ecological receptors, where receptors are populations and communities (EPA 1997, 059370).

Superfund guidance also indicates an ecological risk assessment should be designed to protect local populations and communities of biota rather than individual organisms, except for listed or candidate T&E or treaty-protected species (EPA 1999, 070086). The protection of individuals within these designated protected species could also be protected at the population level; the populations of these species tend to be small, and the loss of an individual adversely affects the species.

In accordance with this guidance, the Laboratory developed generic assessment endpoints (LANL 1999, 064137) to ensure values at all levels of ecological organization are considered in the ecological screening process. These general assessment endpoints can be measured using impacts on reproduction, growth, and survival to represent categories of effects that may adversely impact populations. In addition, specific receptor species, described in section 2.6, are chosen to represent each functional group. The receptor species were chosen based on their presence at the site, their sensitivity to the chemicals of potential concern (COPCs), and their potential for exposure to those COPCs. These categories of effects and the chosen receptor species were used to select the types of effects seen in toxicity studies considered in the development of the toxicity reference values (TRVs). Toxicity studies used in the development of TRVs included only studies in which the adverse effect evaluated affected reproduction, survival, and/or growth.

The selection of receptors and assessment endpoints is designed to be protective of both the representative species used as screening receptors and the other species within their feeding guilds and the overall food web for the terrestrial and aquatic ecosystems. Focusing assessment endpoints on these general characteristics of species that affect populations (versus biochemical and behavior changes that may affect only the studied species) also ensures applicability of the estimates of the effect to the ecosystems of concern.

## 2.6 Screening Receptors

As described in section 2.1, Laboratory property supports numerous habitats with a variety of plant and animal species. The selection of a set of receptors that includes representatives of every class of biota for every trophic level would result in an unwieldy number of receptors for ecological screening. Therefore, the rationale behind receptor selection is to choose an appropriate set of receptors that address the

primary feeding relationships outlined in section 2.4. Receptor selection facilitates the determination of potential adverse ecological impacts across the Laboratory and satisfies the following criteria (based on Fordham and Reagan 1991, 063081).

- The receptor is representative of an exposure pathway, including dietary pathways specified in the functional food web, and nondietary exposure pathways.
- The receptor is representative of a major feeding guild as defined in the functional food web.
- Protection of the receptor is protective of the integrity of ecosystem structure and function.
- The receptor is representative of potentially exposed populations or communities.
- Protection of the receptor is protective of T&E and other species of special interest or concern.
- Toxicity information is available that indicates the receptor is sensitive to contaminants occurring at the Laboratory.
- Exposure information for the species is available, and these data show the species has greater exposure per unit body mass than other candidate species (small species typically have greater intake rates per unit body mass based on allometric relationships [e.g., EPA 1993, 059384]).
- The home range (HR) of the receptor is of an appropriate spatial scale for ecological evaluations at the SWMU or AOC or site aggregate scale, leading to selecting species of small body weight (BW) and therefore small HR to maximize exposure at most SWMUs or AOCs (<0.1 ha to several ha in area).

Given these criteria, the selection of receptors for the Laboratory is outlined below. The selection of terrestrial receptors, including those with links to the aquatic food chain, follows directly from the above logic. The selection of aquatic receptors for radiological contamination is also in direct accord with the logic provided. For nonradionuclide contaminants in aquatic environs, however, the Laboratory has selected methods that are more broadly protective of aquatic ecosystems. These methods include the use of water and sediment benchmarks in ecological screening assessments for aquatic environments. For example, the application of benchmarks for water is targeted at protecting roughly 95% of all aquatic organisms, and thus is inclusive of all trophic guilds illustrated in Figure 2.4-2. The use of benchmarks for screening aquatic environments is recommended in EPA guidance (EPA 1996, 062792). Table 2.6-1 summarizes the factors that led to the selection of the terrestrial, aquatic, and aerial insectivores used for screening.

### **Terrestrial Receptors**

The use of a “generic” plant is indicative of the broad-base taxonomic concern for plants in general rather than any particular species. Additionally, plants are primary producers and form much of the physical habitat structure used by animal species. By using a generic plant, a broadly protective view of the methods for development of ecological screening levels was chosen.



**Table 2.6-1**  
**List of Receptors Selected for Screening at the Laboratory**

Receptor Category	Receptor Species	Selection Factors
Terrestrial autotroph (producer)	Plant	<p>Food source for many animals</p> <p>Provides habitat structure and functional base for terrestrial animals</p> <p>Represents culturally important plants</p> <p>Representative of T&amp;E plant species</p> <p>Direct exposure to contaminated soil</p> <p>Representative of all terrestrial plant species</p>
Soil-dwelling invertebrate	Earthworm	<p>Represents decomposer group important for nutrient cycling</p> <p>Although earthworms are not present in all environmental settings at the Laboratory (absent from more arid locations), they are present in sufficient locations to justify their selection.</p> <p>Large body of toxicity data available for earthworms and other soil-dwelling invertebrates</p> <p>Direct exposure to contaminated soil and detritus</p> <p>Represents a food source for terrestrial vertebrates as discussed below</p> <p>Representative of all soil-dwelling invertebrates</p>
Mammalian herbivore	Mountain cottontail	<p>Food source for carnivores</p> <p>Ubiquitous and abundant</p> <p>Exposure data and toxicity data available</p> <p>Surrogate for socially important browsers (deer and elk)</p>
Mammalian omnivore	Deer mouse	<p>Food source for carnivores</p> <p>Ubiquitous and abundant</p> <p>Exposure data and toxicity data available</p> <p>Surrogate for T&amp;E (New Mexico meadow jumping mouse)</p>
Mammalian insectivore	Montane shrew	<p>Food source for carnivores</p> <p>High fraction of soil in diet relative to rabbit and deer mouse</p> <p>Diet is 100% invertebrates (earthworms) and thereby maximizes this potentially bioaccumulative exposure pathway</p> <p>Surrogate for all terrestrial insectivores, including T&amp;E (Jemez Mountain salamander)</p>
Three diets modeled: Avian omnivore Avian herbivore Avian insectivore	American robin	<p>Food source for some carnivores</p> <p>Exposure data available</p> <p>Large fraction of soil in diet based on eating prey like earthworms</p>
Two diets modeled: Intermediate carnivore Top carnivore	American kestrel	<p>Surrogate for T&amp;E (Mexican spotted owl) by assuming 100% flesh diet</p> <p>Ubiquitous</p> <p>Exposure data available</p> <p>Addresses potential biomagnification from soil</p> <p>Conservative choice for this category, given the food intake to BW ratio</p>

Table 2.6-1 (continued)

Receptor Category	Receptor Species	Selection Factors
Top carnivore	Gray fox	Exposure data available Addresses potential biomagnification from soil Conservative choice for this category, given the food intake to BW ratio
Burrowing mammal	Pocket gopher (air pathway only)	Representative for potential inhalation exposure inside a burrow for fossorial or semifossorial mammals (mouse, gopher, rabbit, fox) Exposure through air pathway only and evaluated only for vapor-phase COPCs
Aquatic community, sediment	Invertebrates	Food source for aquatic animals Ubiquitous and abundant Exposure and toxicity data available
Mammalian aerial insectivore	Occult little brown myotis bat	100% diet may be assumed to come from emergent aquatic insects Allows the consideration of bioaccumulation from aquatic sources to a high-level mammalian receptor
Avian aerial insectivore	Violet-green swallow	100% diet may be assumed to come from emergent aquatic insects Allows the consideration of bioaccumulation from aquatic sources to a high-level avian receptor
Aquatic community, water	Multiple	Generally representative organisms so ecological screening levels (ESLs) are broadly protective of most aquatic species Food source for aquatic animals Ubiquitous and abundant Exposure and toxicity data available
Aquatic autotroph (producer)	Algae (radionuclides only)	Food source for aquatic animals Provides structure (substrate) for animals Ubiquitous and abundant Exposure and toxicity data available
Aquatic omnivore/herbivore	Daphnids (radionuclides only)	Food source for higher trophic levels High exposure to contaminated water and sediment Ubiquitous and abundant Exposure and toxicity data available <i>Daphnia</i> and <i>Ceriodaphnia</i> typically the most sensitive aquatic organisms for a variety of contaminants
Aquatic herbivore (grazer)	Aquatic snails (radionuclides only)	Food source for higher trophic levels High exposure to contaminated sediment Ubiquitous and abundant Exposure and toxicity data available
Aquatic intermediate carnivore	Fish (radionuclides only)	Representative of potential waterborne contaminant effects in the Rio Grande High potential exposure to contaminants; potentially sensitive to persistent bioaccumulators and biomagnifiers

The earthworm (terrestrial worms of the subclass *Oligochaeta*) was selected because it represents the functional category of mechanical decomposers, which are important for nutrient cycling. In addition,

earthworms have a higher exposure to contaminants than other invertebrates because of their high soil intake and intimate soil contact. Earthworms are also present in some Laboratory ecological settings and are food for some middle trophic level vertebrates as discussed below. The earthworm is considered generally protective of all terrestrial invertebrate species, including insects, arachnids, crustaceans, and other taxa.

The mountain cottontail (*Sylvilagus nuttallii*) was selected because it is a strict herbivore (browser/grazer) and can be used as a functional surrogate to evaluate potential effects on large mammalian browsers/grazers (e.g., deer and elk). The deer mouse (*Peromyscus maniculatus*) was selected because of its omnivorous food habits and largely to represent the importance of rodents as a food source for higher consumers (carnivores and omnivores), making it important in the functional food web. The montane shrew (*Sorex monticolus*) was selected largely because of its high exposure to contaminants from grubbing for invertebrates in soil and because of its high-level intake of soil-dwelling invertebrates (including earthworms). The montane shrew also acts as a good receptor when considering a food chain model that includes bioaccumulation of contaminants from soil. The gray fox (*Urocyon cinereoargenteus*) was selected because it represents a mammal with relatively high contaminant biomagnification potential, given its largely carnivorous feeding habits.

The American robin (*Turdus migratorius*) was selected because it is representative of birds that forage for ground-dwelling invertebrates (including earthworms) and fruits, with relatively high potential exposure to contaminants from its diet because of its high food consumption rate per unit body mass. The American robin is considered in several functional roles for avian receptors: an insectivore, herbivore, and omnivore (invertebrate/plant). The American kestrel (*Falco sparverius*) was selected as a top avian carnivore because it serves as a representative of T&E bird species at the Laboratory, namely the Mexican spotted owl. Additionally, abundant information has been gathered for the kestrel's biology, and the kestrel represents an organism with high susceptibility to contaminant biomagnification via terrestrial pathways.

The little brown myotis bat (*Myotis lucifugus occultus*) and the violet-green swallow (*Tachycineta thalassina*) were chosen as receptors for modeling the effects of contaminants bioaccumulated from sediment to insects to aerial insectivores. The former is a species of special concern and considered rare in the Jemez Mountains, although it has been trapped on Laboratory grounds. A large fraction of the brown myotis bat's diet consists of emergent aquatic insects because the habitats surrounding water are favorite hunting areas. The violet-green swallow is common on Laboratory grounds, and some portion of its diet consists of emergent aquatic insects, although its feeding habits are less specialized than that of the brown myotis bat. Nonetheless, both aerial insectivores may be modeled for maximum uptake of aquatic sediment-borne contamination, and information is available on their general biology.

The pocket gopher (*Thomomys bottae*) was selected as the receptor for air inhalation within a burrow because it represents several fossorial and semifossorial species (small mammals like rabbits and foxes) that may occupy burrows at sites with subsurface vapor-phase COPCs present. Gophers spend most of their time underground. Although small mammals like the deer mouse and shrew have smaller BWs and higher weight-normalized air inhalation rates, these species spend much less time underground relative to the gopher. Thus, pocket gophers are a protective representative for all the burrowing mammal species.

Figure 2.6-1 shows the terrestrial food web with a box representing each screening receptor species superimposed over the feeding guilds represented by that receptor. All terrestrial receptors were selected partially on the basis of information available regarding life history habits of the same or similar species (e.g., EPA 1993, 059384).

## Aquatic Receptors

No specific aquatic receptors were selected for the screening assessment of nonradiological contaminants. Methods adopted for screening are considered by the EPA (e.g., 60 Federal Register 15366, “Final Water Quality Guidance for the Great Lakes System, Final Rule”; EPA 1996, 062792) and others (e.g., Jones et al. 1997, 059813) to be protective of aquatic organisms at large (plants, invertebrates, and vertebrates). Although few vertebrates reside in the aquatic realms of the Laboratory, it was considered prudent to adopt methods that are otherwise considered broadly protective and that include organisms that may be found in the Rio Grande (e.g., fish). The aquatic food web, as shown in Figure 2.4-2, is useful for organizing the scoping portion of screening, but for contaminant-based ecological screening comparisons for nonradionuclides, the methods employed broadly cover all species represented in all trophic guilds.

Four aquatic receptors were selected for screening exposure to radionuclides. Algae were selected to represent the producer functional group. Daphnids (*Crustacea*) and snails (*Gastropoda*) were selected to represent the aquatic omnivore and herbivore functional subgroups. The daphnid’s diet in freshwater systems consists primarily of phytoplankton and zooplankton, while snails typically obtain food from scraping lithic and vegetative surfaces for incidental free and attached algae. Some daphnids (e.g., *Daphnia* and *Ceriodaphnia*) represent the most sensitive aquatic organisms to most environmental contaminants. Lastly, although fish are not found on Laboratory property, a “generic” bony fish was selected to represent intermediate carnivores exposed to contaminants.

Figure 2.6-2 shows the aquatic food web with a box representing each screening receptor species superimposed over the feeding guilds represented by that receptor. This figure is specific to receptors for radionuclides but also generally applicable to the aquatic receptors of water and sediment communities. No direct representative is available for the Jemez Mountain salamander, an endangered species with both aquatic and terrestrial life stages. Juvenile salamanders are associated with water, while adults inhabit terrestrial environments. Adult Jemez Mountain salamanders are invertebrate consumers and may be considered functionally similar to shrews; therefore, they are covered by terrestrial screening procedures. It is assumed that juvenile salamanders or other amphibians are represented by the aquatic herbivore and omnivore receptors described above.

## 3.0 DERIVATION OF ESLs

ESLs provide a simple way to characterize the potential for ecological risks in a screening-level ecological risk assessment (SLERA). The Laboratory has developed methods for deriving ESLs for radionuclides and nonradionuclides. Because these methods differ substantially, sections 3.4 and 3.5 describe the methods used to derive ESLs for nonradiological and radiological COPCs, respectively, for soil, sediment, and water. This document describes how screening-level ecological effect levels are calculated (section 3.1), lists the types and sources of ecological effects information (section 3.2), and presents an overview of wildlife exposure and effects evaluations (section 3.3). Although methods for ESL derivation are presented, the ESLs and the supporting information are not included. The Laboratory’s ECORISK Database (LANL 2016, 601838, or latest version) provides the necessary information and documentation as well as the ESLs. Additional details on the derivation of ecological screening levels are presented in Appendix A.




Figure 2.6-1 Screening receptors for terrestrial food web




Figure 2.6-2 Screening receptors for aquatic food web

### 3.1 Screening-Level Ecological Risk Calculation Overview

The Laboratory uses a hazard quotient (HQ) model as a screening-level risk calculation. For multiple chemicals or radionuclides, the hazards are summed into an index. This approach is consistent with both EPA and NMED guidance (EPA 1997, 059370; NMED 2017, 602274; NMED 2017, 602273). The HQ is a ratio between exposure and the effect level of interest (Equation 3.1-1). The hazard index (HI) is a sum of HQs for chemicals of potential ecological concern (COPECs) with similar toxicological modes of action (Equation 3.1-2).

$$HQ_{ij} = \frac{exposure_{ij}}{effect_{ij}} \quad \text{Equation 3.1-1}$$

$$HI_i = \sum_{j=1}^n HQ_{ij} \quad \text{Equation 3.1-2}$$

where  $HQ_{ij}$  is the hazard quotient for receptor  $i$  to COPEC/COPEC  $j$  (unitless)

$exposure_{ij}$  is the exposure concentration for COPEC  $j$  for receptor  $i$

$effect_{ij}$  is the effect level used for screening COPEC  $j$  for receptor  $i$

$HI_i$  is the hazard index for receptor  $i$  to  $n$  COPECs (unitless)

Risks are evaluated relative to an HQ or HI equal to 1. An HQ of less than 1 indicates the COPEC by itself is unlikely to be associated with adverse ecological effects, while an HI of less than 1 indicates the combination of COPECs is unlikely to be associated with adverse ecological effects (EPA 1997, 059370).

The units associated with literature effect levels differ by receptor and type of contaminant (radionuclide or nonradionuclide). The units for exposure (numerator) must match those for the effect level (denominator).

### 3.2 Types and Sources of Ecological Effects Information

The following are the kinds of effect level information for receptors and contaminants available in the literature:

- **Medium concentration** (e.g., mg of contaminant per kg soil). Such effect levels are available for plants, invertebrates, or aquatic organisms where the ecotoxicity studies are based on direct exposures to a contaminated medium (soil, sediment, or water).
- **Wildlife dose** (e.g., mg of contaminant ingested per day [mg/d]). Such effect levels are available for wildlife based on ecotoxicity studies with contaminated food or water.
- **Radiological dose** (e.g., rad per day [rad/d]). These effect levels only apply to radionuclides and can apply to wildlife or other biota for a variety of contaminated medium exposures.

Screening-level ecological risk calculations are simple for those receptors and contaminants that have effect levels with same units as the abiotic media being screened. For wildlife or radionuclide effect levels, some calculations are required to derive ESLs with the proper units.

The Laboratory uses the no observed adverse effect level (NOAEL) or no observed effect concentration (NOEC) as the basis for the ESLs. These values are also used as the toxicity reference values or TRVs. The dose limit for radionuclides is 0.1 rad/d (IAEA 1992, 062802; DOE 2002, 085637). EPA defines the NOAEL or NOEC as the “highest level of a stressor evaluated in a toxicity test or biological field survey

that causes no statistically significant difference in effect compared with controls or a reference site” (EPA 1997, 059370).

ESLs are used to evaluate potential hazards associated with chemicals and radionuclides. The Laboratory has developed chemical-, media-, and receptor-specific ESLs using a tiered TRV development approach, as described in the ECORISK Database (LANL 2016, 601838, or latest version). The Laboratory develops and maintains ESLs as part of the ECORISK Database, which archives the ESLs, TRVs, associated exposure parameters, and all supporting documentation.

The development of an ESL is a two-step process. The first step involves identifying or developing a TRV. In the second step, the TRV and exposure parameters are used to calculate ESLs for chemicals and ecological receptors representative of the ecosystems at the Laboratory. The receptors were selected to be representative of mammals, birds, plants, and invertebrates inhabiting terrestrial and aquatic ecosystems at the Laboratory (Table 2.6-1).

A TRV represents an exposure rate associated with an acceptable risk from chronic exposure of an ecological receptor to a specific contaminant via a specific exposure pathway. In other words, exposures exceeding the TRV may pose adverse effects to wildlife species, while exposures below the TRV are not expected to result in adverse effects (EPA 2005, 089448).

TRVs are important parameters in ESL calculations because “they represent the component of the model that determines whether a contaminant in a media may present potential harm to ecological receptors in the area” (Podolsky et al. 2001, 072586). For any given chemical, TRVs vary among government agencies and private sectors because the methods used to develop them vary according to the site-specific concerns of the organization that developed them (i.e., receptor species, chemical, type of exposure pathway, type, and magnitude of uncertainty factors applied).

The ideal TRV for a SLERA is one that is based on literature representing the most ecologically relevant effects (reproduction/development, survival and/or adult weight/size change); exposure routes (oral ingestion via food or drinking water for birds and mammals, inhalation for mammals, uptake via seed coat and/or roots for plants, and direct contact exposure for invertebrates and aquatic community organisms); exposure media (food and drinking water for birds and mammals, air for mammals, soil for plants and invertebrates, and water and sediment for aquatic community organisms); exposure period (chronic); and effect levels (NOAEL for vertebrates or NOEC for plants and invertebrates). A TRV based on these characteristics is considered protective of the wildlife; aquatic community, plant, and invertebrate populations; and sensitive individuals because it represents an exposure that is not associated with adverse impacts of low-level, long-term chemical effects (i.e., adverse effects on ability of individuals to develop into viable organisms, search for mates, breed successfully, and produce live and equally viable offspring). Therefore, NOAELs and NOECs used for the ESL are meant to be protective of all receptors and levels of biological organization (e.g., individuals and populations). To provide information for a bounding analysis of the potential for ecological risks, lowest-effect ESLs or L-ESLs are also presented in the ECORISK Database.

Laboratory guidance (LANL 2010, 110623) includes guidelines for the literature search, data extraction, default value assignment, and exception ruling for various fields of data entry in customized primary toxicity study evaluation (PTSE) databases, primary toxicity value (PTV) calculation, and TRV derivation. Before performing a PTSE, the primary toxicity literature for the organism and for the exposure pathway and chemical scenario of concern must be identified and collected.

ESLs are chemical- and medium-specific screening levels pertaining to a given receptor (e.g., avian omnivore, earthworm) and medium (sediment, soil, water, and/or air). The TRV is used in the receptor-specific ESL calculation. This equation converts the toxicity value from a dose (mg-contaminant/kg BW/d)



to an environmental concentration (e.g., mg-contaminant/kg-soil) using factors to estimate the transfer of chemical from soil, sediment, or water to dietary media (e.g., soil-to-plant transfer factor [TF]) and receptor-specific exposure parameters (e.g., ingestion/inhalation rates and BW). In the case of plants, earthworms, and aquatic organisms, the ESL is equal to the TRV. Similarly, the L-ESLs are based on lowest observed adverse effect levels [LOAELs] for vertebrates or lowest observed effect concentrations [LOECs] for plants and invertebrates.

### 3.3 Wildlife Exposure and Effects Evaluations

To determine if wildlife receptors receive COPC doses equal to the NOAEL, a wildlife exposure model is developed and used. This wildlife exposure model considers various dietary and nondietary exposure pathways for wildlife. Modeling is not needed to evaluate exposure to nonwildlife species (e.g., plants, soil invertebrates, and aquatic organisms) because it is assumed most of the exposure to these organisms is not related to dietary pathways. Instead, it is assumed plants, soil invertebrates, and aquatic organisms are exposed by direct contact to, and uptake from, a contaminated medium. For example, root uptake for plants is the primary exposure pathway. If site-specific scoping indicates that foliar uptake may be a primary exposure route for a contaminant, the lack of foliar uptake in the plant toxicity testing is addressed in the uncertainty analysis.

Wildlife exposure is derived by intake of COPCs from various sources, including diet, incidental ingestion of contaminated media, dermal contact, and respiration. This general model is presented as Equation 3.3-1 and is based on EPA's general wildlife exposure models (EPA 1993, 059384).

$$E_{total} = E_{oral} + E_{dermal} + E_{respiration} \quad \text{Equation 3.3-1}$$

where  $E_{total}$  is total exposure to a COPC (units are mg/kg/d)

$E_{oral}$  is oral exposure (diet and direct ingestion of contaminated media, with units of mg/kg/d)

$E_{dermal}$  is dermal exposure (with units of mg/kg/d)

$E_{respiration}$  is exposure through respiration or inhalation (with units of mg/kg/d)

For terrestrial wildlife inhabiting the soil surface, it is assumed most contaminant exposure to nonradiological chemicals is through the oral exposure pathway (Sample et al. 1998, 062807). The dermal contact pathway is not typically assessed quantitatively in ecological risk assessments, based on guidance indicating the ingestion route is most important to terrestrial animals (EPA 1997, 059370). Dermal exposure to wildlife is mitigated by the fur or feathers covering the bodies of most vertebrates. In addition, the incidental consumption of soil during grooming is included in the direct soil ingestion estimates. Soil exposure pathway analysis has shown that dermal pathways contribute a small fraction of the dose obtained orally (EPA 2003, 085643). Therefore, the exposure pathways considered in the development of the ESLs for a site capture the primary exposures for wildlife receptors. Inhalation exposures may contribute a significant component of exposure to volatile organic compounds (VOCs) for species occupying burrows for a significant fraction of the time. Therefore, ESLs have been developed for inhalation exposure for VOCs only for burrowing mammals. For other receptor species and for burrowing mammals, for COPCs other than VOCs, the terrestrial wildlife exposure model for nonradionuclides simplifies to Equation 3.3-2.

$$E_{total} = E_{oral} \quad \text{Equation 3.3-2}$$

Although the oral pathway is dominant in most cases, the site-specific scoping should assess the potential importance of the dermal and respiration/inhalation pathways. In cases where dermal and respiration may represent significant exposure pathways, the models presented by Hope (1995, 062783) should be used

to evaluate these pathways. The oral exposure model used for terrestrial wildlife is from EPA's Wildlife Exposure Factors Handbook (EPA 1993, 059384) and is provided in Equation 3.3-3:

**Equation 3.3-3**

$$E_{oral} = C_{soil} \times I_{soil} \times AUF_{soil} + C_{water} \times I_{water} \times AUF_{water} \times (1 / d_{water}) + C_{food} \times I_{food} \times AUF_{food}$$

where  $E_{oral}$  is the estimated oral daily dose for a COPC (mg COPC/kg-BW/day; simplified to mg/kg-BW/d)

$C_{soil}$  is the concentration of chemical constituent x in soil (mg/kg-dry wt)

$I_{soil}$  is the normalized daily soil ingestion rate (kg soil/kg-BW/d, simplified to kg/kg/d)

$AUF_{soil}$  is the area use factor that represents the fraction of soil ingested from a contaminated area (this fraction is set to 1 for the initial screening)

$C_{water}$  is the concentration of chemical constituent x in water (mg/L)

$I_{water}$  is the normalized daily water ingestion rate (kg water/kg-BW/d, simplified to kg/kg/d)

$AUF_{water}$  is the fraction of water ingested from a contaminated area (this fraction is set to 1 for the initial screening)

$d_{water}$  is the density of water (1 kg/L)

$C_{food}$  is the concentration of COPC in food (mg/kg-dry wt)

$I_{food}$  is the normalized daily dietary ingestion rate (kg-food dry wt/kg-BW/d)

$AUF_{food}$  is the fraction of the diet derived from a contaminated area (this fraction is set to 1 for the initial screening)

This model provides an estimate of the oral exposure associated with a concentration of an inorganic or organic chemical toxicant in soil, food, and water, given an organism's normalized daily ingestion rate. Soil ingestion is calculated from a fraction of the dietary intake of soil (EPA 1993, 059384). As a protective assumption appropriate for ecological risk screening, the area use factor (AUF) is set to 1 to indicate the animal receives all its exposure from the contaminated site. An additional conservative assessment is made if the maximum detected concentration is used to represent concentrations in contaminated media and food. The implications of these assumptions should be addressed in the uncertainty analysis.

An implicit assumption of this model is that the bioavailability of the COPC from the environmental media is comparable with the bioavailability of the contaminant in a toxicological experiment. Because little information currently exists on bioavailability conversions, a bioavailability term was not included in the general wildlife exposure model. If bioavailability of a COPC is known and site-specific adjustments to bioavailability are possible, this information should be included in the site-specific uncertainty analysis.

The above model requires all measures of ingestion (except water) to be on a dry-weight basis. Because the EPA presents most normalized food ingestion rates on a wet-weight basis, these dietary constituents must undergo wet-to-dry weight conversions (EPA 1993, 059384). Food intakes rates are provided in units of dry weight, and any conversion factors used in this calculation are also provided. Parameters required for calculations of the general wildlife exposure model, conversions, and other elements of the model are provided for terrestrial vertebrate receptors in Table 3.3-1. The information provided in Table 3.3-1 is for the screening receptors adopted by the Laboratory. It is also important to note that exposure parameters provided in Table 3.3-1 represent conservative upper estimates of potential exposure. More realistic exposure information may be considered in the uncertainty analysis. Information about BW and inhalation rates, which are not required by Equation 3.3-1, is provided to assist with

alternate forms of the wildlife exposure model. For example, the exposure models discussed by Hope (1995, 062783) require these additional parameters.

**Table 3.3-1**  
**Measures Required for the Wildlife Exposure Model**

Receptor	Parameter	Value	Units	Reference	Notes
American kestrel	BW	0.103	kg	EPA 1993, 059384, p. 2-112	Lowest male average weight was 103 g used to provide more conservative ESL value
	Food intake <sup>a</sup>	0.148	kg-food dry wt/kg-BW/d	Nagy 2001, 253420	Estimated using Nagy (2001, 253420) allometric scaling formula for all birds
	Food moisture content	0.68	Proportional	EPA 1993, 059384, p. 4-13	Diet includes insects, birds, mammals, other (EPA 1993, 059384, p. 2-113) (value assumes mammals, birds)
	Fraction soil in diet	0.02	Unitless	none	Assumed
	Soil invertebrate diet	0.5 or 0 <sup>b</sup>	Unitless	EPA 1993, 059384, p. 2-113	Rounded EPA value to 50% to equally expose receptor to potentially contaminated invertebrates and flesh; strict flesh-eater is used to mimic the diet of the Mexican spotted owl
	Flesh diet	0.5 or 1 <sup>b</sup>	Unitless	EPA 1993, 059384, p. 2-113	Rounded EPA value to 50% to equally expose receptor to potentially contaminated invertebrates and flesh; strict flesh-eater is used to mimic the diet of the Mexican spotted owl
	Daily water ingestion rate	0.12	L/kg/d	EPA 1993, 059384, p. 2-112	Used higher of two estimated values
	Home range	106	ha	EPA 1993, 059384	Average of all HR data for woods, forests, and agricultural areas
	Population area	4240	ha	Calculated	40 times HR (see text for explanation)
American robin	BW	0.0773	kg	EPA 1993, 059384, p. 2-197	Used lowest weight of 77 g to provide a conservative ESL
	Food intake <sup>a</sup>	0.35	kg-food dry wt/kg-BW/d	EPA 1993, 059384, p. 2-197	Higher of two empirical values fresh weight food intake rate for robins feeding primarily on fruits, 1.52 kg-food fresh wt/kg-BW/d, multiplied by (100–77)% to account for food moisture content
	Food moisture content	0.77	Proportional	EPA 1993, 059384, pp. 4-13,14	Diet includes invertebrates, plants (fruits), assumed fruit
	Fraction soil in diet: herbivore, omnivore, and insectivore	0.139, 0.152, 0.164	Unitless	Beyer et al. 1994, 062785, Table 1	Used 90th percentile dove value for herbivore diet, 90th percentile woodcock value for insectivore diet, and average of these two species for omnivore diet

Table 3.3-1 (continued)

Receptor	Parameter	Value	Units	Reference	Notes
American robin (continued)	Plant diet	0, 0.5, or 1 <sup>c</sup>	Unitless	None	Modeled with three diets: herbivore, omnivore, and insectivore
	Soil invertebrate diet	1, 0.5, or 0 <sup>c</sup>	Unitless	None	Modeled with three diets: herbivore, omnivore, and insectivore
	Daily water ingestion rate	0.14	L/kg/d	EPA 1993, 059384, p. 2-197	Estimated by EPA from allometric equations
	Home range	0.42	ha	EPA 1993, 059384, p. 2-199	HR data represent average territory size in an open, semiurban environment
	Population area	16.8	ha	Calculated	40 times HR (see text for explanation)
Deer mouse	BW	0.02	kg	EPA 1993, 059384, p. 2-295	For females that have lower BWs and therefore are used to provide a conservative ESL
	Food intake <sup>a</sup>	0.2	kg-food dry wt/kg-BW/d	EPA 1993, 059384, p. 2-296	Based on empirical fresh weight food intake of 0.22 kg-food fresh wt/kg-BW/d (diet of lab chow, 8% to 10% moisture), multiplied by (100–10)% to account for food moisture
	Food moisture content	0.1	Proportional	EPA 1993, 059384, p. 2-296	Moisture content of lab chow used to determine food intake
	Fraction soil in diet	0.02	Unitless	Beyer et al. 1994, 062785, Table 1	For white-footed mouse, most closely related of species available
	Plant diet	0.5	Unitless	EPA 1993, 059384, p. 2-297	Rounded EPA value to 50% to equally expose receptor to potentially contaminated plants and invertebrates
	Soil invertebrate diet	0.5	Unitless	EPA 1993, 059384, p. 2-297	Rounded EPA value to 50% to equally expose receptor to potentially contaminated plants and invertebrates
	Daily water ingestion rate	0.19	L/kg/d	EPA 1993, 059384, p. 2-296	Adult male or female
	Home range	0.077	ha	EPA 1993, 059384, p. 2-298	Average of data from representative environments
	Population area	3.0	ha	Calculated	40 times HR (see text for explanation)
Mountain cottontail	BW	0.560	kg	Sowls 1957, 602507	Minimum of range of reported values, used desert cottontail as surrogate
	Food intake <sup>a</sup>	0.0816	kg-food dry wt/kg-BW/d	Nagy 2001, 253420	Estimated using Nagy 2001, 253420 allometric scaling formula for herbivores
	Fraction soil in diet	0.063	Unitless	Arthur III and Gates 1988, 602506	For black-tailed jackrabbit at Idaho National Laboratory
	Plant diet	1	Unitless	EPA 1993, 059384, p. 2-356	Assume strict herbivore diet

Table 3.3-1 (continued)

Receptor	Parameter	Value	Units	Reference	Notes
Mountain cottontail (continued)	Daily water ingestion rate	0.097	L/kg/d	EPA 1993, 059384, p. 2-355	Estimated by EPA from allometric equations
	Home range	3.1	ha	EPA 1993, 059384, p. 2-357	Average of all HR data for a woodlot and for mixed habitats (used eastern cottontail as surrogate)
	Population area	124	ha	Calculated	40 times HR (see text for explanation)
Montane shrew	BW	0.0054	kg	Bennett et al 1999, 082652	Average of 17 males and females from Sandia Canyon
	Food intake <sup>a</sup>	0.198	kg-food dry wt/kg-BW/d	EPA 1993, 059384, p. 2-213	Higher of two empirical fresh weight food intakes, 0.62 kg-food fresh wt/kg-BW/d, multiplied by (100–68)% to account for food moisture in diet of beef liver
	Food moisture content	0.68	Proportional	EPA 1993, 059384, p. 4-13	Laboratory feeding study used beef liver
	Fraction soil in diet	0.03	Unitless	EPA 2007, 602500, Attachment 4-1, Table 3	Used 90 <sup>th</sup> of the calculated soil intake for the shrew
	Soil invertebrate diet	1	Unitless	EPA 1993, 059384, p. 2-214	Assume strict insectivore diet
	Daily water ingestion rate	0.223	L/kg/d	EPA 1993, 059384, p. 2-213	Only value reported
	Home range	0.39	ha	EPA 1993, 059384, p. 2-214	Reported average HR for one environment.
	Population area	15.6	ha	Calculated	40 times HR (see text for explanation)
Pocket gopher	BW	0.104	kg	Gonzales et al. 2000, 085653	Laboratory-specific minimum measured field value used to provide a conservative ESL
	Inhalation rate	0.089	m <sup>3</sup> /d	EPA 1993, 059384, p. 3-12	Calculated from BW by Equation 3-20 in cited EPA guidance
	Home range	0.06	ha	EPA 1993, 059384, p. 2-214	Reported HR of up to 700 yd <sup>2</sup> (Controlling Pocket Gophers in New Mexico, New Mexico State University Guide L-109; <a href="http://aces.nmsu.edu/pubs/_I/L-109.pdf">http://aces.nmsu.edu/pubs/_I/L-109.pdf</a> )
	Population area	2.4	ha	Calculated	40 times HR (see text for explanation)
Gray fox	BW	3.94	kg	EPA 1993, 059384, p. 2-224	Lowest of four mean values used to provide a conservative ESL (used red fox as a surrogate)
	Food intake <sup>a</sup>	0.0448	kg-food dry wt/kg-BW/d	EPA 1993, 059384, p. 2-224	Red fox female after whelping, empirical fresh weight food intake is 0.14 kg-food fresh wt/kg-BW/d for an unknown diet, multiplied by assumed food moisture content (100–68)%
	Food moisture content	0.68	Proportional	EPA 1993, 059384, p. 4-13	Mean value for mammals and passerine birds
	Fraction soil in	0.028	Unitless	Beyer et al. 1994,	For red fox as a surrogate for gray fox

Table 3.3-1 (continued)

Receptor	Parameter	Value	Units	Reference	Notes
	diet			062785, Table 1	
Gray fox (continued)	Flesh diet	1	Unitless	EPA 1993, 059384, p. 2-224	Rounded diet to 100% flesh
	Daily water ingestion rate	0.086	L/kg/d	EPA 1993, 059384, p. 2-224	Higher of two estimated values
	Home range	1038	ha	EPA 1993, 059384, p. 2-226	Average of all HR data for the red fox over a variety of unspecified environments
	Population area	41,520	ha	Calculated	40 times HR (see text for explanation)
Violet-green swallow	BW	0.0139	kg	Dunning 1993, 073795	Average BW of females for <i>Tachycineta thalassina</i>
	Food intake <sup>a</sup>	0.274	kg-food dry wt/kg-BW/d	Nagy 2001, 253420	Estimated using Nagy (2001, 253420) allometric scaling formula for passerines
	Fraction soil or sediment in diet	0	Unitless	None	Assume no soil or sediment exposure for aerial insectivores
	Invertebrate diet	1	Unitless	None	Assume 100% invertebrate diet
	Daily water ingestion rate	0.242	L/kg/d	EPA 1993, 059384, p. 3-8	Estimated from allometric scaling formula for birds
	Home range	0.68	ha	Bowman 2003, 087148	Using general allometric equation of $10^{[1.8+\log(\text{BW}) \times 1.06]}$
	Population area	27.2	ha	Calculated	40 times HR (see text for explanation)
Occult little brown myotis bat	BW	0.00875	kg	Whitaker 1980, 062889	Used midpoint of reported BW range for <i>Myotis lucifugus</i> (3.1 g to 14.4 g)
	Food intake <sup>a</sup>	0.179	kg-food dry wt/kg-BW/d	Nagy 2001, 253420	Estimated using Nagy (2001, 253420) allometric scaling formula for all mammals
	Food moisture content	0.69	Proportional	EPA 1993, 059384, p. 4-13	Used value for grasshoppers and crickets as surrogate for emergent aquatic insects
	Fraction soil or sediment in diet	0	Unitless	None	Assume no soil or sediment exposure for aerial insectivores
	Invertebrate diet	1	Unitless	None	Assume 100% invertebrate diet
	Daily water ingestion rate	0.159	L/kg/d	EPA 1993, 059384, p. 3-10	Estimated from allometric scaling formula for mammals
	Home range	100	ha	Menzel et al. 2003, 087151	Minimum of 100- to 500-ha HR given for southeastern myotis bat
	Population area	4000	ha	Calculated	40 times HR (see text for explanation)

<sup>a</sup> Normalized ingestion rates are presented in units of kg of food (dry weight) / [kg of BW × d].

<sup>b</sup> Two variants on the American kestrel are used: one more realistically models its actual diet (half invertebrate and half flesh), and the strict flesh-eater is used to mimic the diet of the Mexican spotted owl.

<sup>c</sup> Three variants on the American robin are used: one modeled as a strict herbivore, one an omnivore eating 50% plants and 50% invertebrates, and one modeled as a strict insectivore.

Table 3.3-1 presents information on the spatial scales for exposure to the representative receptors. The HR reflects the area from which individuals may be exposed to contamination. However, EPA guidance is to manage the ecological risk to populations rather than to individuals, with the exception of T&E species (EPA 1999, 070086). One approach to addressing the potential effects on populations is to estimate the spatial extent of the area inhabited by the local population that overlaps with the contaminated area. The population area for each receptor is based on the individual receptor HR and its dispersal distance (Bowman et al. 2002, 073475). Bowman et al. (2002, 073475) estimate that the median dispersal distance for mammals is 7 times the linear dimension of the HR (i.e., the square root of the HR area). If only the dispersal distances for the mammals with HRs within the range of the screening receptors are used, the median dispersal distance becomes 3.6 times the square root of the HR ( $R^2 = 0.91$ ) (Bowman et al. 2002, 073475). If it is assumed the receptors can disperse over the same distance in any direction, the population area is circular, and the dispersal distance is the radius of the circle. Therefore, the population area for each receptor can be derived by  $\pi(3.6\sqrt{\text{HR}})^2$  or approximately 40HR. Table 3.3-1 presents receptor population areas based on 40HR.

### 3.4 ESLs for Chemicals

This section provides an overview of the approach used to develop ESLs for nonradionuclides for soil, burrow air, sediment, and water. Table 3.4-1 summarizes the receptors and diet compositions used in equations for ESL development for each exposure medium.

**Table 3.4-1**  
**Ecological Screening Receptors for Chemicals**

Medium	Receptor Group	Receptor Name	Diet Composition
Soil	Bird	American kestrel	50% invertebrate/50% flesh
		American kestrel	100% flesh
		American robin	100% invertebrate
		American robin	50% invertebrate/50% plant
		American robin	100% plant
	Mammal	Mountain cottontail	100% plant
		Deer mouse	50% invertebrate/50% plant
		Gray fox	100% flesh
		Montane shrew	100% invertebrate
	Plant	Plant	Not applicable <sup>a</sup>
	Invertebrate	Earthworm	Not applicable <sup>a</sup>
Water <sup>b</sup>	Bird	American kestrel	No food, water only <sup>c</sup>
		American robin	No food, water only <sup>c</sup>
		Swallow	No food, water only <sup>c</sup>
	Mammal	Mountain cottontail	No food, water only <sup>c</sup>
		Deer mouse	No food, water only <sup>c</sup>
		Gray fox	No food, water only <sup>c</sup>
		Montane shrew	No food, water only <sup>c</sup>
		Bat	No food, water only <sup>c</sup>
	Aquatic	Multiple aquatic receptors that represent most aquatic organisms	Not applicable <sup>a</sup>

**Table 3.4-1 (continued)**

Medium	Receptor Group	Receptor Name	Diet Composition
Sediment <sup>b</sup>	Bird	Swallow	100% invertebrate
	Mammal	Bat	100% invertebrate
	Aquatic	Multiple aquatic receptors that represent most aquatic organisms	Not applicable
Burrow Air <sup>b</sup>	Mammal	Pocket gopher	Not applicable <sup>d</sup>

<sup>a</sup> ESLs were not calculated for these receptors based on exposure and effects calculations. The ESLs and L-ESLs are values obtained from the literature (NOECs and LOECs).

<sup>b</sup> Water, sediment, and burrow air ESLs are used only to evaluate whether those media may have significant exposure pathways and COPCs because ESLs for one media do not account for exposure to the same COPC in another media. In all cases where a site has sediment or water contaminated, a multimedia assessment is expected.

<sup>c</sup> The water ESL for these terrestrial receptors reflects only exposure from contaminated water from the site. Therefore, a multimedia exposure assessment may be required to address the potential cumulative effects from soil (or sediment) and water for these receptors.


<sup>d</sup> The burrow air ESL applies only to burrowing mammals and only for COPCs that are considered VOCs. The air ESL reflects only exposure from vapors in the air within the burrow. The mammalian herbivore feeding guild has been modeled with the mountain cottontail, so a multimedia exposure assessment to address the potential cumulative effects from soil, water, and air is not possible for this representative species.

### 3.4.1 Soil ESLs

As described in the Laboratory background document for soil, sediment, and Bandelier Tuff, “soil” is defined as material overlaying intact bedrock that has been modified by the addition of organic material or by movement of clay-size particles and by development of ferric hydroxides (LANL 1998, 059730). For the purposes of ecological risk screening, imported fill or disturbed soil is evaluated as well-developed soil because the exposure and transport pathways are similar. Even though tuff and bedrock are not generally considered accessible media to ecological receptors (LANL 2002, 073791), these media are evaluated for ecological risk for purposes of conservatism. For purposes of wildlife exposure, soil is generally assumed to represent the 0.0–5.0-ft interval, but site-specific scoping should present a rationale and justification for the depth interval assumed to represent surface soil.

Although soil ESLs are based on exposure to terrestrial receptors—plants, invertebrates (earthworms), and wildlife—they are determined differently for each receptor. The different approaches are required because of the different ways that toxicological experiments are performed for these organisms. For plants, earthworms, and other soil-dwelling invertebrates, the effects are based on the concentration of a COPC in soil. Therefore, ESLs are directly based on effects concentrations and modeling is not required. Exposure to wildlife, however, is dependent on exposure of the organism to a chemical constituent from a given medium (such as soil or foodstuff) through direct and indirect means and serves as the model for terrestrial exposure calculation (EPA 1993, 059384). The transport and exposure pathways likely to be complete for sites with soil contamination are shown in Figure 3.4-1. Pathways included in all the ESL calculations are designated as “evaluated” in this figure. The pathway for respiration of air vapors is evaluated only for burrow air of terrestrial mammals (section 3.4.2).





Notes: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways. Complete pathways for soil exposure are gray and lines bolded; evaluated pathways are included in the soil ESL calculations.

\* For burrowing animals only.

**Figure 3.4-1 Ecological conceptual site model (CSM) for soil pathways**

For wildlife receptors, ESLs are based on the dietary regimen of the receptor, including consumption of plants, invertebrates, and vertebrate flesh, with some incidental soil ingestion. The general wildlife exposure model is presented in section 3.3. The conversion of soil concentration to dose ingested requires an inversion of the wildlife exposure model (with the intake of contaminated water assumed to be zero) discussed below. This inversion is possible because the food intake value may be related to the concentration in soil. The general basis for this relationship is shown in Equation 3.4-1.

$$C_{food} = C_{soil} \times TF_{food} \quad \text{Equation 3.4-1}$$

where  $C_{food}$  is the concentration of the COPC in food (units are mg/kg)

$C_{soil}$  is the concentration in soil (mg/kg)

$TF_{food}$  is a transfer factor from soil to food (mg/kg dry wt food per mg/kg dry wt soil)

Thus, the general wildlife exposure model can be rewritten in the following form, after setting the AUF to 1 and using the relationship between  $C_{soil}$  and  $C_{food}$  shown in Equation 3.4-2.

$$E_{oral} = C_{soil} \times I_{soil} + C_{soil} \times TF_{food} \times I_{food} \quad \text{Equation 3.4-2}$$

where  $E_{oral}$  is the estimated oral daily dose for a COPC (mg-COPC/kg-BW/d)

$C_{soil}$  is the concentration of chemical constituent x in soil (mg/kg dry wt)

$I_{soil}$  is the normalized daily soil ingestion rate (kg-soil dry wt/kg-BW/d)

$TF_{food}$  is a transfer factor from soil to food (mg/kg dry wt food per mg/kg dry wt soil)

$I_{food}$  is the normalized daily dietary ingestion rate (kg-food dry wt/kg-BW/d)

Because the intake of soil can be related to the intake of food, Equation 3.4-2 can be further simplified to Equation 3.4-3. This manner of modeling soil intake rate is conservative because it assumes incidental soil intake in addition to food intake. An alternate model would be based on total oral intake, and in this alternate model soil and food intake would add up to 100% of the total intake.

$$E_{oral} = C_{soil} \times I_{food} \times [fs + TF_{food}] \quad \text{Equation 3.4-3}$$

where  $E_{oral}$  is the estimated oral daily dose for a COPC (mg-COPC/kg-BW/d)

$C_{soil}$  is the concentration of chemical constituent x in soil (mg/kg dry wt)

$I_{food}$  is the normalized daily dietary ingestion rate (kg-food dry wt/kg-BW/d)

$fs$  is the fraction of soil ingested, expressed as a fraction of the dietary intake

$TF_{food}$  is a transfer factor from soil to food (mg/kg dry wt food per mg/kg dry wt soil)

Because the HQ is a ratio between exposure and effect level of interest (Equation 3.1-1), the right-hand side of Equation 3.4-3 can be the numerator (i.e., exposure), and the TRV can be the denominator (i.e., effect) as shown in Equation 3.4-4.

$$HQ = 1 = \frac{E_{oral}}{TRV} = \frac{C_{soil} \times I_{food} \times [fs + TF_{food}]}{TRV} \quad \text{Equation 3.4-4}$$

Equation 3.4-4 can be rearranged to a basic equation for calculating the soil ESL as shown in Appendix A (Equation A-1.1-1).

The mathematical basis for calculating wildlife ESLs for herbivore, omnivore, insectivore, and carnivore functional groups is presented in Appendix A. These equations show the ESLs are proportional to the effect level. Thus, larger TRVs lead to larger ESL values, which indicate the receptor may be more tolerant of the COPC. The opposite relationship holds for the variables in the denominator of the wildlife ESL model (i.e., a receptor with higher feeding rates or one that eats more contaminated prey has a lower ESL). A receptor with higher exposure will have lower ESLs for the same TRV value as a receptor with lower exposure. The wildlife L-ESLs are calculated with the same equations using the LOAEL for the TRV term.

The minimum soil ESL for each COPC is the lowest receptor-specific soil ESL value available among plants, invertebrates, robin, kestrel, shrew, mouse, cottontail, and fox. For plants and invertebrates, the soil ESL is the NOEC and the L-ESL is the LOEC. Information supporting the selected effect level is provided in the Ecorisk Database (LANL 2016, 601838, or latest version). For wildlife, the soil ESL is calculated as the soil concentration of the COPC that results in an exposure dose equal to the NOAEL.

The wildlife L-ESL is the soil concentration of the COPC that results in an exposure dose equal to the LOAEL.

### 3.4.2 Burrow Air ESLs (Volatile-Phase Contaminants Only)

Quantitative evaluations of ecological risk do not typically include the inhalation pathway because ingestion-related exposure is relatively more important for most chemicals. However, burrow air exposure is potentially a significant exposure pathway for burrowing mammals at some Laboratory SWMUs and AOCs. These SWMUs and AOCs are typically colonized by pocket gophers (*Thomomys bottae*) and other ecological receptors exposed to vapor-phase contaminants (i.e., VOCs) in burrows. Simple fate and transport models indicate vapor-phase contaminants are at much lower concentrations in surface air (Markwiese et al. 2003, 087149), and, therefore, quantitative evaluation of surface air inhalation as a pathway to ecological receptors is not warranted. Vapor-phase contaminants are not prone to bioaccumulation, so the pathways considered for burrow air ESLs are limited to inhalation or respiration of vapors. The pocket gopher is designated as the representative receptor for burrowing mammals. The best estimate of burrow air concentrations is obtained by using soil pore-gas data collected from depths corresponding to those occupied by pocket gophers. Exposure parameters for the pocket gopher are provided in Table 3.3-1. The gopher's inhalation rate (IR) is based on BW according to the allometric equation from Stahl (1967, 063119) shown in Equation 3.4-5:

$$IR = 0.5458 \times BW^{0.80} \quad \text{Equation 3.4-5}$$

It is assumed the gopher stays in its burrow 100% of the time; the exposure through air is described by Equation 3.4-6:

$$E_{air} = \frac{C_{air} \times I_{air}}{BW} \quad \text{Equation 3.4-6}$$

where  $E_{air}$  is the estimated inhalation daily dose for a COPC (mg-COPC/kg-BW/d)

$C_{air}$  is the concentration of chemical constituent x in air inside the burrow (mg/m<sup>3</sup>)

$I_{air}$  is the daily inhalation rate for the pocket gopher (m<sup>3</sup>/d)

$BW$  is the body weight for the pocket gopher (kg)

Appendix A provides additional information and the equation used to calculate burrow air ESLs.


### 3.4.3 Sediment ESLs

Sediment generally exists as young alluvium occurring within or near stream channels, which would be classified as A or C generic horizons in soil nomenclature (LANL 1998, 059730). This definition includes sediment in active channels, inactive channels, and floodplain fluvial geomorphic settings. Sediment can also be found in lentic systems (ponds or lakes), but no lakes and few ponds exist on Laboratory property. Inactive channel and floodplain sediment typically have associated terrestrial ecological communities and, therefore, are more akin to soil from an ecological risk evaluation perspective. Thus, soil ESLs apply to inactive channel and floodplain sediment. Aquatic ecological communities are often associated with perennial and seasonally intermittent aquatic environments and, therefore, sediment-based ESLs are applicable to active channel and pond geomorphic settings with developed aquatic communities.

Because of the typical association of sediment with water, application of sediment ESLs leads to an incomplete evaluation of the potential ecological effects associated with contaminated sediment/water settings. Thus, surface water and multimedia exposure assessments are required in all cases where contaminated sediment is identified. The intent of developing sediment ESLs is to assist in determining the sensitive receptors and major and minor exposure pathways from contaminated sediment, which, in turn, assists in developing an appropriate multimedia exposure model.


Sediment ESLs for the protection of aquatic life are derived from information on direct effects of contaminated sediment on aquatic organisms. Only limited modeling is needed to develop sediment ESLs. Modeling is used to evaluate the potential effects of contaminated sediment on terrestrial receptors through accumulation of COPCs in emergent insects.

General discussion of the transport and exposure pathways considered in the development of sediment ESLs is needed to evaluate the applicability of sediment screening values to the results of site-specific scoping. Pathways of sediment transport to aquatic environs include water as a primary contaminated media through discharge of effluents, directly or indirectly, into perennial and intermittent water bodies; surface water runoff from contaminated soil; infiltration of surface water into shallow and/or deep groundwater; mass wasting; and wind-driven transport of soil-borne COPCs into water courses/bodies. Of primary concern are the first three transport mechanisms included in Figures 3.4-2 and 3.4-3. Rare instances where mass wasting or wind-blown soil may significantly influence the sediment load of a water body are identified during site-specific problem scoping. With the limited water resources in the region, the primary focus should be on pathways of sediment transport from areas adjacent to or contiguous with permanent or seasonally intermittent surface water resources.



Notes: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways. Complete pathways for sediment exposure to aquatic receptors are gray and lines bolded; evaluated pathways are included in the sediment ESL calculations.

**Figure 3.4-2 Aquatic CSM for sediment pathways**



**Figure 3.4-3 Terrestrial CSM for sediment pathways (to account for bioaccumulation concerns)**

Protecting sediment quality is increasingly viewed as a logical extension of water-quality protection, which helps to emphasize the interrelationship between sediment and water as exposure media. Chapman (1989, 062902) cites several reasons for the requirement of sediment ESLs, including

- various toxic contaminants found only in trace amounts in the water column that accumulate in sediment to elevated levels;
- sediment that serves as both a reservoir and a source of contaminants to the water column;
- sediment that integrates contaminant concentrations over time, whereas water column contaminant concentrations are much more variable and dynamic;
- sediment contaminants, in addition to water column contaminants, that affect benthic and other sediment-associated organisms; and
- sediment that is an integral part of the aquatic environment, providing habitat, feeding, and rearing areas for many aquatic organisms.

The general methodologies adopted for screening aquatic receptors to contaminated sediment conform to those proposed by the EPA for developing ecotoxicity thresholds (EPA 1996, 062792). Methods for

screening sediment are based on the assumption that aquatic organisms are generally exposed to the greatest fraction of contamination by means of direct media contact (i.e., continuous bodily contact with sediment). Thus, the exposure pathways for aquatic receptors (using EPA methods) include bioconcentration and, for radionuclides only, external gamma exposure (Figure 3.4-2). Aquatic ecological screening pertains to receptors generally associated with benthic surfaces. Generally, to be protective of aquatic plant and animal species, the EPA methods used in this document have been derived with the intent of protecting a large fraction of species found in aquatic environs at large.

Although sediment ESLs are primarily developed to protect against potential effects on aquatic receptors, pathways from sediment to terrestrial receptors are also evaluated to ensure bioaccumulation concerns have been addressed. A simple wildlife exposure model is developed to evaluate bioaccumulation potential of COPCs in sediment to aerial insectivores (bat and swallow) via emergent insects. The terrestrial receptor exposure model for sediment pathways is provided in Figure 3.4-3. This conceptual model indicates several exposure pathways are complete, but only the food web transport pathway is evaluated because other pathways make only minor contributions. Additionally, the uptake of COPCs from sediment is much more significant for aquatic plants and animals in direct contact with the sediment, which is covered by the sediment pathways model (Figure 3.4-2) and screening methods.

Sediment ESLs come from a variety of sources and may be derived from different measurement endpoints. Further information on sediment benchmark selection for the aquatic community as well as the sediment ESL selection process for the aquatic sediment community (Figure A-1.3-1) is provided in Appendix A. To address transport of COPCs from sediment through the food chain, a wildlife ESL model based on the insectivore soil ESL model has been developed and is presented in Appendix A (Equation A-1.3-1).

In summary, sediment ESLs that are protective of the aquatic community may be derived from a variety of literature sources. Additionally, ESLs are developed for aerial insectivores based on wildlife exposure and effects calculations (Appendix A). The lowest of these values is used as the minimum sediment ESL to ensure that the potential for adverse ecological effects from both direct exposure and ingestion are considered.


#### **3.4.4 Water ESLs**

Water of potential concern to ecological receptors at the Laboratory includes surface water and shallow groundwater. For the purposes of ecological screening, only exposure pathways related to surface water and groundwater that emerges at the surface are evaluated. For those sites where exposure to shallow groundwater is an issue, a discussion of this exposure medium should be included in the uncertainty analysis.

Water samples may be filtered (suspended solids removed) or unfiltered. Unfiltered samples have greater or equal concentrations of COPCs than filtered samples. As a conservative measure of potential exposure, unfiltered water can be used in screening evaluations. If unfiltered samples show no potential risk, no further evaluation of the filtered samples may be needed. If unfiltered samples show potential problems, water samples for inorganic chemical content should be evaluated on the basis of filtered samples because the filtered sample is considered the bioavailable fraction of these constituents in water (EPA 1996, 062792).

Methods for screening water are based on exposure pathways to the aquatic community and to wildlife. For aquatic organisms, the screening approach assumes they are generally exposed to the greatest fraction of contamination by means of direct media contact (i.e., continuous bodily contact with water). Ecological screening for waterborne COPCs pertains to receptors associated with benthic surfaces and

the free water column of both lentic and lotic systems. To be broadly protective of aquatic plant and animal species, New Mexico and the EPA have developed methods to calculate water-quality standards and criteria intended to protect a large fraction (roughly 95%, unless otherwise stated) of species found in aquatic environs. By using these methods, it is assumed that any particular species selected to be representative of feeding guilds in the aquatic realms of the Laboratory will be protected. The exposure model for water pathways to aquatic receptors is provided in Figure 3.4-4. To evaluate potential effects of contaminated water on terrestrial receptors, a wildlife exposure model is developed (Figure 3.4-5). The terrestrial conceptual model is based on exposure to contaminated drinking water.




Notes: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways. Complete pathways for water exposure to aquatic receptors are gray and lines bolded; evaluated pathways are included in the water ESL calculations.

**Figure 3.4-4 Aquatic CSM for water pathways**

Considering the impacts from waterborne contamination to aquatic receptors requires evaluating a number of water-quality criteria or benchmarks that come from a variety of sources, all based upon toxicological information from primary studies. These criteria differ in the methods for their development and/or in the rigor of their development. Consequently, water-quality criteria or benchmarks must be evaluated in a hierarchical fashion, based upon an evaluation of their conservatism or certainty for the protection of approximately 95% of aquatic species.

Water ESLs are selected using water-quality criteria or benchmarks in the order presented below:

1. Section 20.6.4.900 of the New Mexico Administrative Code 20.6.4.900;
2. Ambient Water-Quality Criteria set forth by EPA ([www.epa.gov/ost/criteria/wgctable](http://www.epa.gov/ost/criteria/wgctable));
3. National Oceanic and Atmospheric Administration Screening Quick Reference Tables (Buchman 2008, 206414); and
4. Other sources (see LANL 2016, 601838, or latest version).



**Figure 3.4-5 Terrestrial CSM for water pathways**

Values reported as chronic are used for the ESLs, and those reported as acute are used for L-ESLs. More information on water ESLs for the aquatic community as well as the water ESL selection process for the aquatic community is provided in Appendix A (Figure A-1.4-1).

To address the drinking water exposure pathway to wildlife, an ESL model was developed as described and presented in Appendix A (Equation A-1.4-1), and the parameter values are provided in Table 3.3-1. This model is based on Equation 3.3-3, which is the general wildlife exposure model. To screen the drinking water pathway, it is assumed all oral exposure to water is derived from drinking water. Thus, dose is calculated as follows:

$$E_{water} = C_{water} \times I_{water} \quad \text{Equation 3.4-7}$$

where  $E_{water}$  is the estimated oral daily dose for a COPC (mg-COPC/kg-BW/d)  
 $C_{water}$  is the concentration of chemical constituent x in water (mg/L)  
 $I_{water}$  is the normalized daily water ingestion rate (L of water/kg-BW/d)



### 3.5 Radionuclide ESLs

The methods presented in this section were developed before DOE guidance on the ecological evaluation of radionuclides was established in “A Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota” (DOE 2002, 085637) and “RESRAD-BIOTA: A Tool for Implementing a Graded Approach to Biota Dose Evaluation, User’s Guide, Version 1” (DOE 2004, 085639). However, the methods are consistent with DOE guidance and with the conceptual basis presented by NMED for evaluating ecological effects of radionuclides (NMED 2000, 087104).

The graded approach DOE developed considers the potential for adverse effects on terrestrial, aquatic, and riparian receptors based on three tiers of assessment (DOE 2002, 085637). The first tier provides only a single screening value for each medium (soil, sediment, or water) and thus is similar to the minimum ESLs. However, the first tier of the DOE methods does not provide any way to evaluate the set of receptors and trophic levels considered in this document. Thus, the Laboratory has retained the methods described in this section so screening assessments of radionuclides and nonradionuclides are based on the same set of receptors. When the current Laboratory method is used for radionuclides, the minimum ESLs for soil are lower than those developed under Tier I by DOE for most radionuclides. The notable exceptions are cesium-134, cesium-137, and strontium-90; the minimum ESLs for these radionuclides exceed the DOE screening levels by at least an order of magnitude. These DOE screening levels and their potential impact on the results of the screening assessment should be discussed in the uncertainty analysis.

Radionuclide ESLs are calculated by the dose rate received by individual plants and animals. Radionuclide dose is related to the energy of the specific radioactive decay emission and the amount or mass of the radionuclide. Thus, the basic radionuclide dose model is

$$\text{Dose} = \text{Effective Energy} \times \text{Amount} \quad \text{Equation 3.5-1}$$

For calculating radionuclide ESLs, “dose” is expressed in units of rad/d, while the “amount” of the radionuclide is expressed in units of pCi/g, which is an activity (decay per unit time) per unit mass of media or organism. Thus, effective energy has units of rad/d per pCi/g, which indicates the effective energy term can also be viewed as a dose conversion factor (DCF).

Radionuclide ESLs require calculations to account for the dose received from internal (within the organism) and external (from contaminated media) sources. The difference between the radionuclide and nonradionuclide wildlife models is that the radionuclide models require calculating the internal concentration or body burden and the nonradionuclide models require calculating the exposure to the contaminant. Conversion factors are also required to account for the effective energy for different types of radionuclides in different media. The same receptor species are used to model terrestrial exposure to radionuclides and nonradionuclides, with the exception that aquatic receptors for radionuclides consist of four specific groups (algae, daphnids, snails, and fish); aquatic ESLs for nonradionuclides are based on standards and benchmarks considered to be broadly protective of all aquatic species.

### 3.5.1 Radionuclide Dose Limits and Effects Calculations

Radionuclide dose limits are the equivalent of the NOAELs used to develop nonradionuclide ESLs. The International Atomic Energy Agency (IAEA) has concluded doses protective of human health are protective of ecological resources, except under the following conditions when doses protective of human health may not provide adequate protection of ecological resources (IAEA 1992, 062802):

- human access is restricted but access by biota is not restricted,
- unique exposure pathways exist,
- T&E species are present, or
- other stresses are significant.

For these four situations, IAEA recommends a dose limit of 0.1 rad/d. Because this dose limit is considered appropriately conservative and is consistent with the results of the National Council on Radiation Protection (NCRP) reviews (NCRP 1991, 062803) and Eisler (1988, 600338), the Laboratory has adopted 0.1 rad/d as the dose limit for all ecological receptors for screening purposes. Thus, the basic model for calculating acceptable dose for radionuclides is

$$\text{Total Acceptable Dose} = 0.1 \text{ (rad/d)} = \text{Internal Dose} + \text{External Dose} \quad \text{Equation 3.5-2}$$

DOE has also recommended 0.1 rad/d as the dose limit for wildlife, but DOE has specified 1 rad/d as the basis for plant and aquatic animal screening values, and DOE has not developed screening levels that are specifically protective of soil invertebrates (DOE 2002, 085637). Thus, the Laboratory has selected a more protective dose limit for plant and aquatic receptors as the no effect level for the ESL. For the L-ESLs, the Laboratory increases the DOE's dose limit by a factor of 10 to 1 rad/d.

As discussed below, all radionuclide ESLs are calculated using the target dose level. The target dose level is the denominator in the HQ calculation (Equation 3.1-1), and the numerator is the receptor and medium-specific internal and external dose. The radionuclide ESLs are calculated as the concentrations predicted to be equal to the target dose (i.e., would yield an HQ = 1).

### 3.5.2 Soil ESLs

Section 3.4.1 presents the operational definition of soil. Radionuclide soil ESLs are based on exposure of terrestrial receptors to contaminated soil. The minimum radionuclide soil ESL is the lowest receptor-specific ESL among the terrestrial receptors. ESLs are developed to account for dose from a single radionuclide.

The radiological dose to terrestrial biota is the sum of the dose from internally deposited radionuclides and the external dose from the same radionuclides in soil. The transport pathways included in the calculations for radionuclides in soil are identical to those for nonradionuclides (Figure 3.4-1).

Conservative assumptions about the size of the organism, its diet, the geometry of the contaminated source, and the location of the receptor relative to the contaminated source are used in the methods presented in this document to estimate internal and external doses. Thus, the calculations overestimate dose and are used for screening purposes only. The calculations for estimating internal and external doses from radionuclides in soil are derived from Higley and Kuperman (1996, 062804). The basic model for calculating acceptable dose from soil for radionuclides is

$$\text{Dose}_j = C_{\text{organism},j} \times DCF_{\text{int},j} + C_{\text{soil},j} \times DCF_{\text{ext},j} \quad \text{Equation 3.5-3}$$

where  $Dose_j$  is the total acceptable dose from radionuclide  $j$  (rad/d)

$C_{organism,j}$  is the internal concentration of radionuclide  $j$  (pCi/g of organism)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g BW)

$C_{soil,j}$  is the concentration of radionuclide  $j$  in soil (pCi/g)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g soil)

Internal dose results from exposure to radionuclides through plant uptake, incidental soil ingestion, and food web uptake (Figure 3.4-1). External dose is based on exposure to gamma-emitting radionuclides from contaminated soil (Figure 3.4-1). The basis for calculating internal and external dose and radionuclide ESLs is provided in Appendix A (section A-2.1).

Nonradionuclide and radionuclide ESL calculations share many common variables. Thus, much of the discussion concerning uncertainty in the nonradionuclide ESLs is directly relevant to the radionuclide ESLs. Three variables—the retention time, the TF from food to blood, and the DCFs—are unique to radionuclides. The retention time and blood TFs vary between species and are based on laboratory experimental data. Thus, some uncertainty in these values exists. However, the retention time typically does not impact the ESL, except for radionuclides with short biological clearance times (like tritium). The DCFs are based on the physical properties of each radionuclide and typically have less uncertainty, especially in the screening context where worst-case assumptions are made. The soil radiological L-ESLs are calculated with the same equations in Appendix A using 1 rad/d as the dose limit.

### 3.5.3 Sediment ESLs

The operational definition of sediment is discussed in section 3.4.3. Radionuclide sediment ESLs are based on exposure of contaminated sediment to aquatic receptors and to the bat and swallow through ingestion of contaminated prey. The minimum radionuclide sediment ESL is the lowest receptor-specific ESL among the aquatic receptors as well as the bat and swallow. ESLs are developed to account for dose from a single radionuclide.

An ESL calculation for aquatic organisms exposed to sediment is based on the models presented by Baker and Soldat (1992, 062801). The radiological dose to aquatic organisms is the external dose from the radionuclide in sediment; the internal dose from sediment radionuclides is accounted for in the water ESL calculations for aquatic organisms for radionuclides (Baker and Soldat 1992, 062801; DOE 2002, 085637). Sediment-based thresholds used for screening values do not exist for radionuclides, so algae, daphnids, snails, and fish have been selected as surrogates for organisms living in aquatic environments at the Laboratory. Transport pathways from sediment to aquatic organisms are presented in Figure 3.5-1.

To address bioaccumulation and some biomagnification, the bat and swallow have been selected as higher-trophic-level terrestrial receptors that feed primarily upon insects emerging from sediment in aquatic environments. ESLs calculated for these receptors assume they are feeding 100% upon aquatic invertebrates. The pathways for bat and swallow exposure to sediment are the same as those presented in Figure 3.4-3.

The basic model for calculating acceptable dose from sediment to aquatic organisms for radionuclides is


$$Dose_j = C_{sediment,j} \times DCF_{ext,j} \quad \text{Equation 3.5-4}$$

where  $Dose_j$  is the total acceptable dose from radionuclide  $j$  (rad/d)

$C_{sediment,j}$  is the concentration of radionuclide  $j$  in sediment (pCi/g dry sediment)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry sediment)

More information on the basis for deriving radionuclide ESLs in sediment is provided in Appendix A (section A-2.2). The sediment radiological L-ESLs are calculated with the equations in Appendix A using 1 rad/d as the dose limit.




**Figure 3.5-1 Aquatic CSM for sediment pathways**

### 3.5.4 Water ESLs

The operational definition of water is discussed in section 3.4.4. Radionuclide water ESLs are based on exposure of contaminated surface water to aquatic and terrestrial receptors. The minimum radionuclide water ESL is the lowest receptor-specific ESL among the four aquatic and eight wildlife receptors. ESLs are developed to account for dose from a single radionuclide. Calculation of ESLs for aquatic organisms is based on the models presented by Baker and Soldat (1992, 062801). The radiological dose to aquatic receptors is the sum of the dose from internally deposited radionuclides and the external dose from the same radionuclides in water. In this model, the internal dose calculated for water ESLs for aquatic receptors includes the internal component associated with sediment as well because the bioaccumulation factor considers the partitioning of the radionuclide between sediment and water (Baker and Soldat 1992, 062801; DOE 2002, 085637). Thus, paired data for water and sediment are needed to assess the radionuclide dose. Media-based screening values for radionuclides do not exist, so algae, daphnids,

snails, and fish have been selected as assessment endpoint surrogates for receptors living in aquatic environments at the Laboratory. Transport pathways to aquatic organisms are presented in Figure 3.5-2.



Notes: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways. Complete pathways for water exposure to aquatic receptors are gray and lines bolded; evaluated pathways are included in the water ESL calculations.

**Figure 3.5-2 Aquatic CSM for water pathways**

The only water exposure pathway considered for terrestrial receptors is ingestion of drinking water (Figure 3.4-5). The basic model for calculating acceptable dose from water for radionuclides is

$$Dose_j = C_{organism,j} \times DCF_{int,j} + C_{water,j} \times DCF_{ext,j} \quad \text{Equation 3.5-5}$$

where  $Dose_j$  is the total acceptable dose from radionuclide  $j$  (rad/d)

$C_{organism,j}$  is the internal concentration of radionuclide  $j$  (pCi/g of organism)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g)

$C_{water,j}$  is the concentration of radionuclide  $j$  in water (pCi/mL)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/mL)

More information on the basis for deriving radionuclide ESLs in water is provided in Appendix A (section A-2.3). The water radiological L-ESLs are calculated with the equations in Appendix A using 1 rad/d as the dose limit.

## 4.0 SCREENING-LEVEL ECOLOGICAL RISK ASSESSMENT

The SLERA is conducted only for sites known or suspected to have COPCs present in soil, sediment, or water. The SLERA consists of three steps:

1. The scoping evaluation (or problem formulation) described in section 4.1;
2. The screening evaluation (or the screening-level risk and uncertainty analysis) described in sections 4.2 and 4.3; and
3. Risk interpretation (or screening-level risk characterization) described in section 4.4.

During the initial step, the ecological risk assessor should determine if COPCs are known or expected to occur at the site. If not, the site should be recommended as requiring no further ecological evaluation in the risk assessment. Although these recommendations are made for an individual SWMU or AOC, in the rest of this document, the term *site* is used broadly to represent a SWMU or AOC or an aggregate of SWMUs and/or AOCs. The information presented in this section is an overview of the SLERA. Risk assessors are also referred to recent investigation reports for the steps and details involved in performing a SLERA.

### 4.1 Scoping Evaluation

Sites being investigated to determine the nature and extent of contamination as well as the potential need for corrective actions must undergo ecological scoping, including conducting a site visit and completing the ecological scoping checklist (Appendix B). The ecological exposure CSM is developed during scoping using the ecological scoping checklist. Fate and transport issues relative to ecological concerns are assessed during scoping. The scoping evaluation should address whether a SWMU or AOC should be combined (aggregated) on an appropriate scale to support risk-based corrective action decision-making with neighboring SWMUs or AOCs for the purposes of the SLERA. Sites may be combined based on size, geography, common contaminants, common transport pathways, common land use, common receptors and/or habitat, or on programmatic considerations. For ecological risk, sites may be aggregated on a larger scale than might be used to consider human health risk. Any aggregation of the SWMUs and/or AOCs under consideration should be determined before the SLERA begins.

After the scoping evaluation, if the risk assessor determines the site poses no threat to the environment because no ecological receptors and/or no pathways to receptors exist, a recommendation is made that no further assessment of ecological risk is necessary. The justification for this recommendation is documented in the risk assessment.

During scoping, a decision is made about the adequacy of the data and the CSM for the screening evaluation. At a minimum, the SLERA must be performed for all relevant media (e.g., soil, sediment, or water) that have a complete ecological exposure pathway. Before the screening evaluation can be performed, site-specific data must be deemed adequate for characterizing the nature and extent of contamination. Data adequacy in scoping involves determining whether the geographic and biotic limits of sampling as well as depths and media sampled match the potential extent of contamination at the site. If adequate data do not exist for the site, a recommendation must be made to collect additional data. It should be noted that when data are adequate and appropriately distributed, the upper confidence limit (UCL) of the mean concentration may be used instead of the maximum detected concentration in calculations and comparisons. The UCLs of the mean concentration is calculated using the EPA ProUCL program (<http://www.epa.gov/osp/hstl/tsc/software.htm>).

The goals of the ecological scoping evaluation are to identify sites that need a screening evaluation, assess the need for an aggregate assessment, identify COPCs, determine data adequacy for screening,

evaluate the potential for environmental contaminant transport, and establish likely exposure pathways. The scoping evaluation is equivalent to the site-specific problem-formulation step.

#### 4.1.1 Ecological Scoping Checklist

The purpose of the ecological scoping checklist is to

- describe the site setting and the known form of contaminant releases;
- confirm complete exposure pathways to ecological receptors exist;
- determine if the site should be combined with other sites for screening, and establish the functional/operational boundaries of the assessment;
- determine if adequate data exist for the screening evaluation, primarily as related to the nature and extent of contamination;
- prepare for the screening evaluation by determining whether screening should encompass terrestrial and/or aquatic receptors; and
- gather information to develop the CSM (e.g., what are the dominant/important transport pathways, exposure routes, and receptors).

Completion of the ecological scoping checklist consists of three steps:

1. Assembling and initially interpreting information on the nature of releases, site history and operations, potential for off-site transport, and biological receptors potentially impacted by releases.
2. Visiting the site to validate information from Step 1 and collecting field notes to complete the CSM. The site visit may be used to document the presence or lack of receptors and off-site migration pathways. Notes are also made regarding the applicability of existing data for determining the nature and extent of contamination.
3. Completing the CSM diagrams identifies the complete and incomplete exposure pathways as well as the major and minor pathways.

##### 4.1.1.1 Checklist Step 1: Assemble Existing Information

To prepare for the site visit, the following information should be obtained: (1) the most current biological assessment information for the site (typically the Biological and Floodplain Assessment document for the applicable operable unit and/or TA); (2) information on site erosion potential; (3) investigation work plans or reports that provide information on contamination source, sampling locations, analytical suites, and sampling results; (4) geographic information system maps that show (if applicable) neighboring SWMUs and AOCs, sampling locations, vegetation types, watershed name, and wetlands; and (5) historical and current aerial photographs to help document changes in site operations and conditions. The information obtained is documented in Part A of the ecological scoping checklist (Appendix B). The information required for Part A of the checklist includes (1) site identification; (2) nature of releases (solid, liquid, gaseous, or other); (3) a list of the primary impacted media (surface soil, water, sediment, subsurface soil-tuff, or other); (4) specification of the applicable vegetation classes (open water, aspen-riparian-wetland, mixed conifer-spruce-fir, grassland, shrub land, urban-sparse-bare rock, ponderosa pine, and piñon-juniper); (5) identification of T&E habitat, if present (list species if applicable); (6) a list and description of neighboring/contiguous/upgradient SWMUs and AOCs, and discussion of whether the site should be aggregated with additional SWMUs and/or AOCs for screening, if appropriate; (7) surface water erosion potential; and (8) documentation of other scoping meeting notes (as appropriate).

#### **4.1.1.2 Checklist Step 2: Site Visit**

The main objective of the site visit is to confirm whether ecological receptors are present and can be exposed to site contaminant releases. A secondary objective is a qualitative evaluation of whether site data provide adequate information to determine the nature and extent of contamination. The site visit should be arranged at an appropriate time of year (ideally, spring or summer) to best evaluate biota at the site. If the site visit is planned for another time of year, uncertainties introduced in the initial biological assessment by such timing must be noted.

Maps showing sampling locations and results (if available) and a camera are recommended for the site visit. The need for other equipment or supplies to locate and measure site features should be determined during the scoping meeting. Such additional resources may include a measuring device to approximately locate relevant biological features (measuring tape and/or rangefinder and pin flags or other markers to specify locations for surveying).

Part B of the checklist is completed during the site visit and includes administrative information such as the site identification, date of site visit, and personnel conducting visit. Part B also includes receptor information, primarily aimed at determining whether ecological receptors are present at the site. Contaminant transport information, emphasizing surface water and other modes of transport, is documented in Part B. Part B also provides ecological effect information, including notes on physical disturbance and obvious ecological effects (such as dead vegetation or lack of fossorial activity).

If no complete pathways to receptors and no transport pathways to off-site receptors are present, the remainder of the checklist (Part C) is not completed, and any additional explanation/justification is provided to conclude that the site poses no threat to the environment. An example of “no pathways/no receptors” is a mesa-top site with buried, inaccessible contamination with no potential for off-site transport. However, a site that lacks receptors because of high levels of contamination would not qualify for the “no pathways/no receptors” stopping point.

If receptors and pathways are present, then subsequent questions in Part B involving data adequacy are addressed. Specifically, do existing data provide adequate information on the nature and extent of contamination? Also, do existing data for the site address potential pathways of site contamination and receptor exposure? Based on the evaluation of existing data, additional data may be required to resolve adequacy and/or quality issues. For example, if the COPCs at a site are based on elevated detection limits, the risk assessor should encourage resampling or reanalysis to obtain detection limits that are appropriate and usable in the ecological screening evaluation. Similarly, if vertical and/or lateral extent of the contamination is not defined to permit an ecological risk assessment, a recommendation for additional sampling should be provided. Once data issues are resolved, the process of scoping and screening the site for potential ecological impacts should proceed.


Completion of Part B also includes additional field notes on the site setting and potential ecological receptors to document other site observations relevant to the ecological screening evaluation of the site. Such information may include observations on the variability in the type and density of ecological receptors present at the site. Of particular interest are any field notes that could be used to document factors considered in the uncertainty analysis.

#### **4.1.1.3 Checklist Step 3: Ecological Conceptual Site Model**

Part C of the checklist relates to the CSM for ecological receptors and consists of questions related to contaminant transport and the potential for exposure of biota (Appendix B). Answers to questions in Part C are used to complete the CSM (Figures 4.1-1 and 4.1-2 as well as Ecological Pathways Exposure Model figures in back of checklist). This model is used to select appropriate ecological screening




receptors (terrestrial, aquatic, or both) and helps to interpret the results of the ecological screening assessment in a site-specific manner.



Note: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways.

**Figure 4.1-1 Terrestrial receptor conceptual exposure and transport model**



Note: Boxes with Xs indicate incomplete pathways. Open boxes indicate complete pathways.

**Figure 4.1-2 Aquatic receptor conceptual exposure and transport model**

The generic terrestrial receptor CSM is shown in Figure 4.1-1. The questions provided in the scoping checklist help evaluate the transport and exposure routes to terrestrial receptors. The model evaluates surface soil, groundwater, surface water/sediment, and the subsurface as potentially contaminated media. Figure 4.1-1 also illustrates the transport pathways that may lead to contaminated air, surface water/sediment, or groundwater as secondary contaminated media. Two exposure routes are available to terrestrial receptors from air: respiration of vapors or inhalation/deposition of particulates. Respiration includes exposure to plants and invertebrates, and inhalation refers to exposure to wildlife. Five possible exposure routes are available to terrestrial receptors from contaminated soil: plant uptake, food web transport, incidental ingestion, dermal contact, and external gamma. Five possible exposure routes are available to terrestrial receptors from contaminated water/sediment: plant uptake, food web transport, drinking water ingestion, dermal contact, and external gamma. Groundwater may be an exposure medium for deep-rooted plants but typically does not have complete exposure pathways to animals.

The generic aquatic receptor CSM is shown in Figure 4.1-2. The questions provided in the scoping checklist help evaluate the transport and exposure routes to aquatic receptors. This model shows surface soil, groundwater, surface water/sediment, and the subsurface as possible primary contaminated media. Figure 4.1-2 also shows transport pathways that may lead to surface water/sediment or groundwater as secondary contaminated media. The aquatic model does not consider transport to air because volatile contaminants are rapidly lost from surface water and sediment, and the potential for dust generation in damp sediment is unlikely. Thus, the aquatic model is most relevant to sites with perennial water. Sites with intermittent sources of water may need to be evaluated in both terrestrial and aquatic site conceptual models to ensure all contaminant exposure pathways are evaluated. Three possible exposure routes are available to aquatic receptors from contaminated surface water/sediment: bioconcentration, bioaccumulation, and external gamma. Bioconcentration covers all nontrophic exposure routes, which include respiration and dermal absorption. Bioaccumulation covers only trophic exposure routes (i.e., food web transport).

## 4.2 Screening Evaluation

Once the scoping process is complete, the screening evaluation is conducted. The goal of the screening evaluation is to identify the COPECs by exposure medium, and the outcome of the evaluation is to determine whether contaminants pose a potential unacceptable risk to ecological receptors. The evaluation is intended to be protective of the environment, not predictive of ecological risk. Thus, conservative assumptions are made throughout the screening evaluation to ensure contaminants, exposure pathways, and sensitive species are not missed.

Identification of COPECs first requires assembling exposure point concentrations (EPCs) and ESLs for all media, receptors, and COPCs. All the ESLs for the receptors in a chemical-medium combination are obtained from the Laboratory's ECORISK Database (LANL 2016, 601838, or latest version); the lowest ESL for that chemical in that medium becomes the minimum ESL used for the ecological screening. If the HQ for a COPC at a site with only a single COPC is greater than 1 or the HQ for a COPC is greater than 0.3 for a site with multiple COPCs, then that COPC is identified as a COPEC. The HQs are calculated for each receptor/COPEC combination and are the ratio of a receptor's exposure at the site to an acceptable effects level (i.e., the ESL).

The minimum ESLs are specific to the medium and include values for soil, sediment, and water, as appropriate. Each medium and COPC has a minimum ESL. The minimum ESL is the lowest applicable ESL for a COPC in soil, sediment, and water and is intended to be protective for all ecological receptors in a given functional group for exposure to that single medium. The site EPC and the minimum ESL are used to calculate the COPC and medium-specific HQ and the sum of COPEC HQs or HI for a receptor (section 3.1).

The ESLs and the toxicity and other parameter information required for their calculation are maintained in the Laboratory's ECORISK Database (LANL 2016, 601838, or latest version). The ECORISK Database is available to anyone performing or reviewing ecological screening assessments for the Laboratory, and updates to this database are issued as new information becomes available. The current version of the database is available for downloading at <http://www.lanl.gov/community-environment/environmental-stewardship/protection/eco-risk-assessment.php>.

The ESL comparisons and HQ/HI calculations are followed by an uncertainty analysis that focuses on key sources of uncertainty in the screening assessment and may result in adding or removing COPECs. The main components of the uncertainty analysis are described in section 4.3.

Following the uncertainty analysis, the results of the screening assessment are provided to the risk managers. At this point, an ecological scientific management decision point (SMDP) is required. As part of this SMDP, a risk-management strategy may be recommended by the risk assessors. Possible recommendations and risk-management strategies are discussed in section 4.4.

## 4.3 Uncertainty Analysis

Much of the uncertainty in the screening assessment is addressed by applying exposure and toxicity values designed to be protective of all the receptors. However, the net result is likely to overestimate exposure to ecological receptors from contaminated media. Thus, more accurate estimates of exposure can be evaluated by considering factors such as area use and bioavailability of COPECs (Pastorok et al. 1996, 062784).

Many factors are incorporated in the development of the ESLs, and uncertainty is associated with values for the factors and the model itself. At a minimum, the uncertainty analysis should focus on the

key sources of uncertainty. Examination of the uncertainty can result in adding or deleting COPECs. The uncertainty analysis may qualitatively discuss factors that may overestimate or underestimate the potential risk to ecological receptors at the site.

Uncertainties associated with ESLs fall into two main categories. The first group is associated with COPCs, including toxicity and bioavailability (or TFs between soil/sediment/water and food). The second group relates to receptors, including feeding rates, the amount of incidental soil/sediment/water ingestion and diets. These uncertainties are addressed by selecting inputs to the ESL calculations that represent worst-case conditions. For example, carnivores could have mammalian and avian prey, which would tend to reduce exposure because of the lower fat content of birds versus mammals. Uncertainties are also addressed by using the lowest receptor-specific ESL as the minimum ESL for each COPC to ensure the screening evaluation is protective and inclusive of all COPCs.

Bioavailability is often a key parameter in the evaluation of exposure to wildlife, and mechanistic bioconcentration or bioaccumulation models can be evaluated for their applicability (Jager 1998, 062786). One important factor not considered in developing wildlife ESLs is the potential for biomagnification of COPCs in higher trophic levels. The carnivore is modeled as eating herbivore or insectivore prey, which has consumed potentially contaminated plants or insects. However, this model does not account for top carnivores that may be eating prey with more complex diets (e.g., a raptor that eats a snake that preys on lizards that eat predaceous insects that eat herbivorous insects). Developing models to account for multiple trophic level transfers is complex and beyond the realm of screening. The potential for biomagnification for top carnivores depends on factors relating to the spatial distribution of the COPC relative to the distribution of prey and the biological retention time within the prey. This uncertainty should be discussed on a site-specific basis where potentially biomagnifying COPCs are identified.

Body weight is the main covariate for many of the parameters in the wildlife soil ESL models. Body weight has an allometric relationship to gross food intake rates (Nagy 1987, 062782) and is also used as a normalizing factor for food intake and the NOAEL values. Some studies also show relationships between body size and toxicity (e.g., Newman et al. 1994, 062788). The energy value of the food consumed by the animal also shows a relationship to food intake (Nagy 1987, 062782). For example, an animal consuming a low-energy food source must consume a greater quantity to support its basal metabolism. Thus, interrelationships exist between diet composition, BW, and food intake. Relationships also exist between BW and HR because small animals tend to have smaller HRs (Cotgreave 1993, 062905). Thus, screening receptors were selected to be relatively small species within a feeding guild, which will tend to have smaller HRs and greater food intake per unit body mass.

As noted above, one of the goals of the approach to calculating soil ESLs is to ensure COPECs or pathways are not eliminated prematurely. Thus, more realistic modeling, including the application of nonlinear TF relationships, is viewed as unnecessary for the purposes of screening.

Individual AUFs and population area use factors (PAUFs) may be appropriate to modify the estimate of risk to some receptors at some sites depending on the size of the site. The introduction of area use reduces potential overestimation of risks to receptors whose HRs are larger than the area of contamination being evaluated. These AUFs/PAUFs may be applied to either individual organisms or populations. Area use may be particularly important for species that represent both a feeding guild and serve as a surrogate for a T&E species with a different HR than the surrogate. Because T&E species must be assessed on an individual basis (EPA 1999, 070086), the AUF is used for the Mexican spotted owl. The flesh-eating kestrel represents both the feeding guild of carnivorous birds (using its normal HR) and serves as a surrogate for the Mexican spotted owl (which has a much larger HR).

#### 4.3.1 Development of AUFs

EPA guidance recommends evaluating ecological effects at the population rather than at the individual level (EPA 1999, 070086), except when evaluating T&E species. The initial screening using ESLs generates HQs and HIs designed to estimate the potential for risk to individual ecological receptors, assuming continuous exposure to the representative concentration of the COPC in question. The AUF is calculated based on the ratio of the site area to the HR of an individual receptor to reflect the fact a receptor actually moves around its HR and does not remain stationary in the contaminated site. Therefore, the individual AUF assesses the level of individual exposure based on the area of the HR. The modification of an HQ and/or HI with a PAUF uses the estimated area occupied by the population of a receptor species to assess the likelihood of any individual within the assessment population encountering the contaminated area, while using the same ESL based on effects to individuals to determine the impact of this contact within the contaminated area. The PAUF assumes impacts to some individuals and estimates the average effect on the assessment population of that impact. Application of AUFs and/or PAUFs to the results of the ecological screening is generally beyond the screening level and begins to examine the uncertainty associated with the estimates of potential risk generated by the screening analysis. The PAUF puts exposure from a contaminated site in perspective of possible population impacts and provides a reasonable basis for characterizing potential ecological risks to wildlife.

The AUF is used to account for the amount of time that a receptor is likely to spend within the contaminated area based on the size of the receptor's HR. Because T&E species must be assessed on an individual basis (EPA 1999, 070086), the AUF is used for the Mexican spotted owl based on an HR of 366 ha. The kestrel (top carnivore) is used as the surrogate receptor for the Mexican spotted owl. If Mexican spotted owls are potentially exposed receptors for a site, then the uncertainty analysis should include a discussion of the impact on HQs and HIs of the surrogate species when the HR of the Mexican spotted owl is used instead of the HR of the surrogate.

As discussed in section 4.3, PAUFs are developed based on investigations correlating the HR of a receptor with its dispersal distance (the distance an animal moves from its natal HR). The dispersal distance has been shown to affect population structure, demographics, and spacing patterns and can be used to determine the assessment population boundaries (Bowman et al. 2002, 073475). When HR is expressed as its linear dimension (the square root of HR), it has a good linear correlation with dispersal distance for the same species (Bowman et al. 2002, 073745). For mammals with similar HR sizes to the species used as screening receptors at the Laboratory, dispersal distance is equal to 3.5 times the square root of the HR. The relationship holds well for small mammals such as mice and rabbits but may overpredict dispersal distance for fossorial species and slightly underpredict dispersal distance for some large herbivores such as the white-tailed deer (Ryti et al. 2004, 600901). The mathematical relationship between HR and dispersal distance has been estimated only for mammals, but for the calculations at these sites the same methodology was applied to avian receptors. Bird species have higher median and maximum dispersal distances than similar-sized mammals (Sutherland et al. 2000, 073460), so application of the mammalian relationship is protective of bird species because this relationship underestimates the dispersal distance and, therefore, the avian assessment population area.

The dispersal distance from the center of the HR can be considered the radius of the animal's population area, with the area likely to be occupied by members of that population (the assessment population area) consisting of the circle described by the area covered by the dispersal distance. The assessment population area would therefore be equal to  $\pi r^2$ , which would be equal to  $\pi$  times (3.5 times the HR)<sup>2</sup>. This mathematical relationship can be simplified to 40 times the HR as a representation of the assessment population area in hectares (Ryti et al. 2004, 600901). Once the population area is calculated for each receptor species of interest, the area of the site can be divided by the population area to develop

a site-specific PAUF for that population. HRs and population areas (40HRs) for the receptors are presented in Table 3.3-1.

AUFs and PAUFs cannot be calculated for the plant and earthworm because these receptors do not have an HR that can be related to an individual or population assessment area. The plant and earthworm are evaluated directly against their EPCs. Assessment populations of plants and earthworms are evaluated in a more qualitative manner.

#### **4.3.2 Exposure-Related Parameters**

The CSMs for terrestrial and aquatic ecosystems describe the potential pathways that may apply to soil, sediment, or water at sites being evaluated. These models should be reviewed as part of the uncertainty analysis to determine if complete pathways exist at the site under consideration that were not included in the development of the ESLs. The exposure pathways addressed by the ESL and HQ/HI analysis include all complete exposure pathways, with the exception of foliar uptake by plants, inhalation, and dermal exposure. Although the last two pathways contribute to the dose received by animals, the contribution is relatively small and does not interfere with determining the COPECs. Soil ingestion rates, however, can represent one of the more significant sources of environmental exposure, up to 18% for grazing species in areas of sparse vegetation, and over 10% for some birds and aquatic insects (Beyer et al. 1994, 062785). Therefore, the exposure pathways considered in developing the ESLs used in the screening assessment for a specific site capture the primary exposures for wildlife receptors at this site. ESLs incorporate all the exposure pathways described above; the ESLs overestimate the dose ingested if some of the pathways are not complete at the site, for example, if the contaminated media was buried at a depth inaccessible to wildlife receptors.

For complete pathways used to develop ESLs at a site, the equations used to calculate ESLs from the TRVs include terms for BW, water intake, food intake, and inhalation rate (gopher only). To provide a conservative estimate of the ESL, maximum estimates of intake factors (food, water, air) were combined with lower estimates of BW. This approach maximizes the weight-specific dose to the receptor and is protective of all species within a feeding guild represented by a screening receptor. It may overestimate potential risk to larger-size species or to small-size species with lower intake rates than those used in the model.

As discussed above, risk to far-ranging species may also be overestimated because the area used to develop ESLs is 100%. Depending on the size of the site, this value may be appropriate for small-size species but is likely to overestimate risk for larger-size species with a HR greater than the size of the site.

Uncertainty is associated with the values used for the EPC and the potential risk it represents. The uncertainty analysis should consider whether use of the maximum detected concentration of a COPC as the EPC is likely to overestimate the potential ecological risk to receptors or whether the EPC may underestimate the exposure at a site. Use of the UCL as the EPC is likely to overestimate risk if the receptor has an HR greater than the area over which the UCL was determined. The analysis of uncertainty associated with the EPC should also consider findings of the data review (e.g., precision and bias of sample results for environmental media samples) and the impact of the review on the confidence and representativeness of the concentration estimate.

The uncertainty analysis discusses aspects of the conservative risk-screening process that over- or underestimate potential risk to receptors and thereby affect site decisions. In the case of the SLERA, one uncertainty is related to the exposure of receptors to COPEC concentrations not likely to result in adverse impacts. This overestimation of risk to receptors exposed either to naturally occurring levels or to exposure that cannot be distinguished from naturally occurring levels is described and put in the context

of whether an increased risk to receptors exists. Therefore, the discussion and analysis are appropriate when determining whether COPECs contribute to increased potential risk at a site.

The EPCs (either the UCL, the maximum detected concentration, or the maximum detection limit) are evaluated relative to the concentrations measured in samples of soil, sediment, and tuff from uncontaminated areas of the Pajarito Plateau (LANL 1998, 059730). This uncertainty discussion and analysis are not related to whether an inorganic chemical was detected above background and is a COPC but rather to whether COPCs identified and retained as COPECs result in a potential increased risk to receptors at the concentration representing exposure at the site. Furthermore, the presence of a concentration or concentrations above the background values that resulted in the identification of a COPC does not mean the level of exposure to the COPC poses an increased risk. For example, if the UCL for copper is 8.3 mg/kg and the measured background concentrations range from 0.25 mg/kg to 16 mg/kg for soil and 0.25 mg/kg to 6.2 mg/kg for tuff, the mean exposure to copper across the site is the same as if the receptor were exposed to naturally occurring levels of copper. In addition, because the UCL for copper background concentrations is 6.4 mg/kg and the UCL for site concentrations is 8.3 mg/kg, the difference in the potential risk associated with these concentrations is negligible. Thus, risk from copper to ecological receptors cannot be distinguished from, or does not incrementally increase above, that associated with naturally occurring levels, making any further assessment of copper and risk unnecessary. If, on the other hand, the EPC for copper is 117 mg/kg, exposure across the site is above naturally occurring levels of copper and may pose a potential risk to ecological receptors. In this case, further assessment of copper is conducted to determine if a potential risk exists at this mean exposure level.

The comparison of EPCs with background concentrations is relevant in the context of uncertainty associated with potential risks and exposures to COPECs. If, as defined, the UCL is intended to represent the average concentration of a contaminant that a receptor is exposed to at the site, then an average concentration that is indistinguishable from what occurs naturally does not result in an increased potential exposure or risk. Because the risk is not increased by the EPC over what may be expected from naturally occurring concentrations, the risk is overestimated, and the uncertainty associated with this overestimation should be eliminated from the risk estimate. In addition, if the EPC is the maximum detected concentration or maximum detection limit for that inorganic chemical from 0.0–5.0 ft below ground surface (bgs), then a comparison with background is appropriate.

Although site-to-background comparisons were conducted to identify COPCs, a reevaluation of COPC concentrations relative to background at the risk-assessment stage is warranted because the concentrations used at this point are depth-dependent. In the initial comparison to background, all sampling results, regardless of depth, are used for each medium. However, in the case of ecological risk, only data from 0.0–5.0 ft bgs are included. This approach may result in a subset of data being used to assess risk and warrants a reevaluation of concentrations to background, especially if the maximum detected concentration or the maximum detection limit is used as the EPC. EPCs based on 95% UCLs would not necessarily be greater than 95% UCLs calculated for the background data set, and it is incorrect to assume that exposure represented by 95% UCLs for inorganic COPECs would be different on average than exposure to naturally occurring levels.

For example, manganese is a COPEC with an EPC (95% UCL) of 670 mg/kg. The ESLs for the earthworm and plant are 450 mg/kg and 220 mg/kg, respectively, which results in HQs of approximately 2 and 3. Manganese background concentrations range from 22 mg/kg to 1100 mg/kg for soil, sediment, and tuff combined. This results in manganese HQs ranging from 0.05 to 5 for these two receptors based on naturally occurring manganese concentrations: the earthworm and plant HQs for the maximum Qbt 2,3,4 background concentration (752 mg/kg) are approximately 2 and 3. It is clear that the risks (HQs) from the 95% UCL are the same as the HQs from background concentrations (i.e., the risks are not increased

above those present from naturally occurring concentrations). The 95% UCL represents the mean exposure at the site despite some concentrations being above background for a given medium. This mean exposure is used as the basis for whether potential risk exists at the site. It makes no difference whether the concentration divided by the ESL is a 95% UCL or a single concentration, the HQ is the same, and, therefore, the associated risk is the same. If the mean exposure does not add additional risks to what could in theory result from naturally occurring concentrations, then it should not be included in the overall analysis. The uncertainty analysis is designed to illustrate whether the “risks” estimated based on conservative screening values reflect potential impacts to receptors. Because in this case the manganese HQs do not reflect potential impacts to receptor, potential risk from manganese should not be included as part of the overall risk for this site.

#### **4.3.3 Toxicity-Related Parameters**

Another key uncertainty is the availability of toxicity information for receptor groups (e.g., birds, mammals, plants, and invertebrates). The toxicity data and uncertainty factors used to develop the ESLs may potentially overestimate the actual toxicity of a chemical to a receptor, particularly when those data are extrapolated from one species to another. In addition, the comparison of EPCs to ESLs assumes the chemical species or form at the site is identical to the chemical species used in the toxicity analysis. The absence of toxicity information greatly reduces the meaning of a screening assessment, and the uncertainty analysis should determine the impact of missing or incomplete toxicity information on identifying COPECs.

TFs are used to estimate the potential for accumulation of contaminants through the levels of the food chain. TFs based on linear equations are used to generate ESLs. They are not well documented, and many are based on the physical properties of a chemical instead of empirically measured values. Although the linear TFs are considered conservative, other models available can predict higher levels of accumulation. Equations based on TFs also do not account for any depuration from the organism, which tends to overestimate the concentrations at higher trophic levels. Therefore, the models and TFs used to generate the ESLs may over- or underestimate the actual concentrations within an organism, particularly at higher trophic levels.

Many sites have multiple COPECs; cumulative effects and contaminant interactions may alter the safe threshold for exposure to any or all of these COPECs. However, the ESL calculation is modeled on the assumption of the additive effects of chemicals. This assumption could overestimate or underestimate the actual impact of exposure to multiple contaminants from synergistic or antagonistic effects. No information is available for most chemicals on synergistic or antagonistic effects; therefore, almost all risk assessments assume the effects are additive when multiple chemical contaminants are present.

The ESLs also include the implicit assumption that the chemical form of the COPEC is likely to be present in the environment in the same form and with the same bioavailability as the chemical form used in toxicity studies. In general, toxicity studies use readily bioavailable forms of chemicals; the TRVs from these studies may overestimate the toxicity of the chemical form of a COPEC in the environment. Because TRVs are derived from toxicity studies with whole animals, the TRVs are based on the potential effects of both the administered chemical and the metabolic products of that chemical. The form of the chemical in the toxicity study may differ from that found in the environment, however, which means the chemical form at the site could potentially have different metabolic products.

Because of these uncertainties, ESLs for some inorganic chemicals may be below background concentrations of those chemicals. In cases where the background concentration is below the ESL, this issue should be addressed in the uncertainty section. An HQ for the background concentration may be presented to show the contribution of background to the overall estimate of potential risk at the site. If the



EPC for the site is within the range of background concentrations, the uncertainty analysis should also discuss whether the EPC indicates an elevated risk or represents an exposure similar to background across the site.

#### **4.3.3.1 COPECs without ESLs**

Some COPECs do not have ESLs for any receptor in the ECORISK Database because literature searches for relevant toxicity data for these chemicals either have not been completed or no usable toxicity data exist. To address this uncertainty, several online toxicity databases have been or can be searched to determine if any relevant toxicity information is available. The online databases typically searched include the EPA Ecotox Database, EPA Office of Pesticide Programs Aquatic Life Benchmarks, U.S. Army Corps of Engineers/EPA Environmental Residue-Effects, California Cal/Ecotox Database, Pesticide Action Network Pesticide Database, U.S. Army Wildlife Toxicity Assessment Program, U.S. Department of Agriculture Integrated Pesticide Management Database, American Bird Conservancy Pesticide Toxicity Database, and Oak Ridge National Laboratory Risk Assessment Information System. Although some COPECs do not have any relevant toxicity data in the online databases listed above, a search of the literature continues in an effort to determine if any relevant toxicity information exists.

In the absence of a chemical-specific ESL, COPEC concentrations may be compared with ESLs for a surrogate chemical. Comparison to surrogate ESLs provides an estimate of potential effects of a chemically related compound and a line of evidence to indicate the likelihood that ecological receptors are potentially impacted.

Some COPECs without ESLs do not have chemical-specific toxicity data or surrogate chemicals to be used in the screening assessments and cannot be assessed quantitatively for potential ecological risk. These COPECs are often infrequently detected across the site. In these cases, comparisons with residential human health soil screening levels (SSLs) are presented as part of a qualitative assessment. The comparison of COPEC concentrations to residential human health SSLs is a viable alternative for several reasons. Animal studies are used to infer effects on humans and constitute the basic premise of modern toxicology (EPA 1989, 008021). In addition, toxicity values derived for the calculation of human health SSLs are often based on potential effects that are more sensitive than the ones used to derive ESLs (e.g., cellular effects for humans versus survival or reproductive effects for terrestrial animals). EPA also applies uncertainty factors or modifying factors to ensure the toxicity values are protective (i.e., they are adjusted by uncertainty factors to values much lower than the study results). COPEC concentrations compared with these values are an order of magnitude or more below the SSLs, which corresponds to uncertainty factors of 10 or more. Therefore, it is assumed the differences in toxicity would not be more than an order of magnitude for any given chemical. The relative difference between values provides a weight of evidence that the potential toxicity of the COPEC is likely to be low or very low to the receptor(s). The COPECs without ESLs may be common to many of the sites and are discussed separately for each site.

#### **4.3.4 L-ESL Analysis**

Sites may have adjusted HIs greater than 1 for one or more receptors. To address these HIs and reduce the associated uncertainty, an analysis is conducted using L-ESLs calculated based on a LOAEL/LOEC rather than a NOAEL/NOEC. The L-ESLs are calculated based on toxicity information in the ECORISK Database (LANL 2016, 601838, or latest version). The analysis addresses some of the uncertainties and conservativeness of the NOAEL/NOEC-based ESLs used in the initial screening assessments. The HI analyses are conducted using the LOAEL/LOEC-based ESLs. The HQs and HIs calculated for this subset of receptors and COPECs are also adjusted using the PAUFs, if applicable, when the wildlife receptor HIs

exceed 1 using the L-ESLs. L-ESLs are presented in the ECORISK Database (LANL 2016, 601838, or latest version).

#### **4.3.5 Comparison with Previous Investigations**

A comparison of COPEC concentrations reported in the canyon investigations, where field and/or laboratory studies and tests have been conducted to provide empirical data, may be presented to reduce the uncertainty related to HIs greater than 1. The premise for this comparison is that if the field and laboratory studies/tests have not found any ecological effects on receptors at similar or higher concentrations than detected at a site, then the concentration(s) at a site would also not impact ecological receptors even though the screening HI is greater than 1.

Biota investigations have been conducted in canyon reaches in Los Alamos and Pueblo Canyons (LANL 2004, 087390); Mortandad Canyon (LANL 2006, 094161; LANL 2007, 098279); Pajarito Canyon (LANL 2009, 106939); and Sandia Canyon (LANL 2009, 107453). Field and laboratory studies included collecting and analyzing soil, sediment, and water samples; monitoring cavity-nesting birds and analyzing their eggs; trapping small mammals and analyzing whole organisms; conducting earthworm bioaccumulation tests (i.e., measuring growth and survival and analyzing whole organisms); laboratory testing of sensitive organisms; and performing seedling germination tests.

#### **4.4 Risk Interpretation**

At the completion of the screening evaluation, the risk assessor communicates the results to the risk manager, with an emphasis on the uncertainty analysis. The purpose of the communication is to provide the risk manager with sufficient information to support a risk-management decision with respect to potential ecological concerns. It is the responsibility of the risk manager to determine if sufficient information is provided to identify a risk-management strategy (in terms of ecological concerns) or if more information is needed to better inform the risk-management decision.

Some of the recommendations and risk-management strategies that could result from the screening assessment include the following:

1. Adequate information is not available to make a risk-management decision. The result would be to identify data needs, based on the results of the screening, and to develop a plan to collect additional data.
2. Adequate information is available to conclude the ecological risks are negligible and no additional investigation of ecological risk is recommended. For example, no unacceptable risks are inferred if the screening evaluation identifies no COPECs.
3. Ecological risks are not negligible, but the information is not sufficient to indicate adverse ecological effects are occurring. The risk management strategy is to reduce uncertainties in the screening assessment by conducting a baseline ecological risk assessment.
4. Sufficient lines of evidence are available to document potential or actual adverse ecological effects such that remediation is warranted.

#### **5.0 REFERENCES**

*The following reference list includes documents cited in this report. Parenthetical information following each reference provides the author(s), publication date, and ERID or ESHID. This information is also included in text citations. ERIDs were assigned by the Associate Directorate for Environmental*

Management's (ADEM's) Records Processing Facility (IDs through 599999), and ESHIDs are assigned by the Environment, Safety, and Health Directorate (IDs 600000 and above). IDs are used to locate documents in the Laboratory's Electronic Document Management System and in the Master Reference Set. The NMED Hazardous Waste Bureau and ADEM maintain copies of the Master Reference Set. The set ensures that NMED has the references to review documents. The set is updated when new references are cited in documents.

LANL (Los Alamos National Laboratory). November 2015. "Threatened and Endangered Species Habitat Management Plan for Los Alamos National Laboratory." Los Alamos National Laboratory document LA-UR-15-28610. Los Alamos, New Mexico. (LANL 2015, 602156)

Arthur III, W.J., and R.J. Gates, March 1988. "Trace Element Intake via Soil Ingestion in Pronghorns and in Black-Tailed Jackrabbits," *Journal of Range Management*, Vol. 41, No. 2, pp. 162-166. (Arthur and Gates 1988, 602506)

Baker, D.A., and J.K. Soldat, June 1992. "Methods for Estimating Doses to Organisms from Radioactive Materials Released into the Aquatic Environment," PNL-8150, UC-602, Battelle, Pacific Northwest Laboratories, Richland, Washington. (Baker and Soldat 1992, 062801)

Beyer, W.N., E.E. Connor, and S. Gerould, 1994. "Estimates of Soil Ingestion by Wildlife," *Journal of Wildlife Management*, Vol. 58, No. 2. (Beyer et al. 1994, 062785)

Bowman, J., 2003. "Is Dispersal Distance of Birds Proportional to Territory Size?," *Canadian Journal of Zoology*, Vol. 81, No. 2, pp. 195–202. (Bowman 2003, 087148)

Bowman, J., J.A.G. Jaeger, and L. Fahrig, 2002. "Dispersal Distance of Mammals is Proportional to Home Range Size," *Ecology*, Vol. 83, No. 7, pp. 2049-2055. (Bowman et al. 2002, 073475)

Buchman, M.F., 2008. "NOAA Screening Quick Reference Tables," National Oceanic and Atmospheric Administration (NOAA) OR&R Report 08-1, Office of Response and Restoration Division, Seattle, Washington. (Buchman 2008, 206414)

Chapman, P.M., July 1989. "Current Approaches to Developing Sediment Quality Criteria," *Environmental Toxicology and Chemistry*, Vol. 8, No. 7, pp. 589–599. (Chapman 1989, 062902)

Cotgreave, P., July 1993. "The Relationship between Body Size and Population Abundance in Animals," *Trends in Ecology and Evolution*, Vol. 8, No. 7, pp. 244–248. (Cotgreave 1993, 062905)

DOE (U.S. Department of Energy), July 2002. "A Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota," DOE Standard No. DOE-STD-1153-2002, U.S. Department of Energy, Washington, D.C. (DOE 2002, 085637)

DOE (U.S. Department of Energy), January 2004. "RESRAD-BIOTA: A Tool for Implementing a Graded Approach to Biota Dose Evaluation, User's Guide, Version 1," ISCORS Technical Report 2004–02, U.S. Department of Energy document DOE/EH-0676, Washington, D.C. (DOE 2004, 085639)

Dunning, J.B., Jr. (Ed.), 1993. *CRC Handbook of Avian Body Masses*, CRC Press, Inc., Boca Raton, Florida. (Dunning 1993, 073795)

EPA (U.S. Environmental Protection Agency), December 1989. "Risk Assessment Guidance for Superfund, Volume I: Human Health Evaluation Manual (Part A), Interim Final," EPA/540/1-89/002, Office of Emergency and Remedial Response, Washington, D.C. (EPA 1989, 008021)

- EPA (U.S. Environmental Protection Agency), December 1993. "Wildlife Exposure Factors Handbook," Vol. I of II, EPA/600/R-93/187a, Office of Research and Development, Washington, D.C. (EPA 1993, 059384)
- EPA (U.S. Environmental Protection Agency), January 1996. "Ecotox Thresholds," ECO Update, Intermittent Bulletin Vol. 3, No. 2, EPA 540/F-95/038, Office of Solid Waste and Emergency Response, Washington, D.C. (EPA 1996, 062792)
- EPA (U.S. Environmental Protection Agency), June 5, 1997. "Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments, Interim Final," Office of Emergency and Remedial Response, Washington, D.C. (EPA 1997, 059370)
- EPA (U.S. Environmental Protection Agency), April 1998. "Guidelines for Ecological Risk Assessment," EPA/630/R-95/002F, Risk Assessment Forum, Washington, D.C. (EPA 1998, 062809)
- EPA (U.S. Environmental Protection Agency), October 7, 1999. "Issuance of Final Guidance: Ecological Risk Assessment and Risk Management Principles for Superfund Sites," OSWER Directive No. 9285.7-28 P, Office of Solid Waste and Emergency Response, Washington, D.C. (EPA 1999, 070086)
- EPA (U.S. Environmental Protection Agency), November 2003. "Guidance for Developing Ecological Soil Screening Levels," OSWER Directive No. 9285.7-55, Office of Solid Waste and Emergency Response, Washington, D.C. (EPA 2003, 085643)
- EPA (U.S. Environmental Protection Agency), February 2005. "Guidance for Developing Ecological Soil Screening Levels," OSWER Directive No. 9285.7-55, Office of Solid Waste and Emergency Response, Washington, D.C. (EPA 2005, 089448)
- EPA (U.S. Environmental Protection Agency), April 2007. "Attachment 4-1, "Guidance for Developing Ecological Soil Screening Levels (Eco-SSLs)," *Exposure Factors and Bioaccumulation Models for Derivation of Wildlife Eco-SSLs*," OSWER Directive No. 9285.7-55, Office of Solid Waste and Emergency Response, Washington, D.C. (EPA 2007, 602500)
- Fordham, C.L., and D.P. Reagan, July 1991. "Pathways Analysis Method for Estimating Water and Sediment Criteria at Hazardous Waste Sites," *Environmental Toxicology and Chemistry*, Vol. 10, No. 7, pp. 949–960. (Fordham and Reagan 1991, 063081)
- Gallaher, B.M., and R.J. Koch, September 2004. "Cerro Grande Fire Impacts to Water Quality and Stream Flow near Los Alamos National Laboratory: Results of Four Years of Monitoring," Los Alamos National Laboratory report LA-14177, Los Alamos, New Mexico. (Gallaher and Koch 2004, 088747)
- Gallegos, A.F., G.J. Gonzales, K.D. Bennett, and L.E. Pratt, February 1997. "Preliminary Risk Assessment of the Mexican Spotted Owl under a Spatially-Weighted Foraging Regime at the Los Alamos National Laboratory," Los Alamos National Laboratory report LA-13259-MS, Los Alamos, New Mexico. (Gallegos et al. 1997, 057915)
- Gonzales, G.J., R.L. Budd, P.R. Fresquez, and R.J. Wechsler, July 2000. "Source Document: The Relationship Between Pocket Gophers (*Thomomys bottae*) and the Distribution of Buried Radioactive Waste at the Los Alamos National Laboratory," Los Alamos National Laboratory report LA-13719-MS, Los Alamos, New Mexico. (Gonzales et al. 2000, 085653)

- Gonzales, G.J., A.F. Gallegos, and T.S. Foxx, 1997. "Second Annual Review Update, Preliminary Risk Assessment of Federally Listed Species at the Los Alamos National Laboratory," Los Alamos National Laboratory document LA-UR-97-4732, Los Alamos, New Mexico. (Gonzales et al. 1997, 062879)
- Gonzales, G.J., A.F. Gallegos, T.S. Foxx, P.R. Fresquez, M.A. Mullen, L.E. Pratt, and P.E. Gomez, 1998. "A Spatially Dynamic Preliminary Risk Assessment of the Bald Eagle at the Los Alamos National Laboratory," Los Alamos Scientific Laboratory report LA-13399-MS, Los Alamos, New Mexico. (Gonzales et al. 1998, 062350)
- Gonzales, G.J., A.F. Gallegos, M.A. Mullen, K.D. Bennett, and T.S. Foxx, 1998. "Preliminary Risk Assessment of the Southwestern Willow Flycatcher (*Empidonax traillii extimus*) at the Los Alamos National Laboratory," Los Alamos National Laboratory report LA-13508-MS, Los Alamos, New Mexico. (Gonzales et al. 1998, 062349)
- Gonzales, G.J., R. Ryti, A. Gallegos, and P. Newell, March 2004. "Modeled Ecological Risk to the Deer Mouse, Mexican Spotted Owl and Western Bluebird at the Los Alamos National Laboratory Using ECORISK.7," Los Alamos National Laboratory report LA-14118, Los Alamos, New Mexico. (Gonzales et al. 2004, 085207)
- Hathcock, C.D., D.C. Keller, and L.A. Hansen, May 2015. "Sensitive Species Best Management Practices Source Document (Updated March 2015)," Los Alamos National Laboratory document LA-UR-15-20981, Los Alamos, New Mexico. (Hathcock et al. 2015, 602499)
- Higley, K.A., and R. Kuperman, August 1996. "Appendix C, Ecotoxicological Benchmarks for Radionuclide Contaminants at RFETS," in *Ecological Risk Assessment Methodology, Technical Memorandum for Ecological Chemicals of Concern (ECOCs) Screening Methodology*, RF/ER-96-0039, Kaiser-Hill/Rocky Flats Environmental Technology Site, Golden, Colorado. (Higley and Kuperman 1996, 062804)
- Hope, B.K., June 1995. "A Review of Models for Estimating Terrestrial Ecological Receptor Exposure to Chemical Contaminants," *Chemosphere*, Vol. 30, No. 12, pp. 2267–2287. (Hope 1995, 062783)
- IAEA (International Atomic Energy Agency), 1992. *Effects of Ionizing Radiation on Plants and Animals at Levels Implied by Current Radiation Protection Standards*, International Atomic Energy Agency, Vienna, Austria. (IAEA 1992, 062802)
- Jager, T., October 1998. "Mechanistic Approach for Estimating Bioconcentration of Organic Chemicals in Earthworms (*Oligochaeta*)," *Environmental Toxicology and Chemistry*, Vol. 17, No. 10, pp. 2080–2090. (Jager 1998, 062786)
- Jones, D.S., G.W. Suter, II, and R.N. Hull, November 1997. "Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Sediment-Associated Biota: 1997 Revision," Report No. ES/ER/TM-95/R4, Oak Ridge National Laboratory, Oak Ridge, Tennessee. (Jones et al. 1997, 059813)
- Kraig, D.H., R. Ryti, D. Katzman, T. Buhl, B. Gallaher, and P. Fresquez, March 2002. "Radiological and Nonradiological Effects After the Cerro Grande Fire," Los Alamos National Laboratory report LA-13914, Los Alamos, New Mexico. (Kraig et al. 2002, 085536)
- LANL (Los Alamos National Laboratory), 1990. "Annotated Bibliography of Geologic, Hydrogeologic, and Environmental Studies Relevant to Solid Waste Management Units at Los Alamos National Laboratory," Los Alamos National Laboratory document LA-UR-90-3216, Los Alamos, New Mexico. (LANL 1990, 047588)

- LANL (Los Alamos National Laboratory), April 1997. "Core Document for Canyons Investigations," Los Alamos National Laboratory document LA-UR-96-2083, Los Alamos, New Mexico. (LANL 1997, 055622)
- LANL (Los Alamos National Laboratory), September 22, 1998. "Inorganic and Radionuclide Background Data for Soils, Canyon Sediments, and Bandelier Tuff at Los Alamos National Laboratory," Los Alamos National Laboratory document LA-UR-98-4847, Los Alamos, New Mexico. (LANL 1998, 059730)
- LANL (Los Alamos National Laboratory), June 1999. "General Assessment Endpoints for Ecological Risk Assessment at Los Alamos National Laboratory," report prepared for Los Alamos National Laboratory, Los Alamos, New Mexico. (LANL 1999, 064137)
- LANL (Los Alamos National Laboratory), March 2000. "Installation Work Plan for Environmental Restoration Project, Revision 8," Los Alamos National Laboratory document LA-UR-00-1336, Los Alamos, New Mexico. (LANL 2000, 066802)
- LANL (Los Alamos National Laboratory), September 4, 2002. Personal communication between K. Olson, K. Bostick, R. Mirenda, M. Nagy, P. Schumann, and M. Tardiff, Environmental Restoration Project, Los Alamos, New Mexico. (LANL 2002, 073791)
- LANL (Los Alamos National Laboratory), April 2004. "Los Alamos and Pueblo Canyons Investigation Report," Los Alamos National Laboratory document LA-UR-04-2714, Los Alamos, New Mexico. (LANL 2004, 087390)
- LANL (Los Alamos National Laboratory), October 2006. "Mortandad Canyon Investigation Report," Los Alamos National Laboratory document LA-UR-06-6752, Los Alamos, New Mexico. (LANL 2006, 094161)
- LANL (Los Alamos National Laboratory), June 2007. "Revised Risk Assessments for Mortandad Canyon," Los Alamos National Laboratory document LA-UR-07-4010, Los Alamos, New Mexico. (LANL 2007, 098279)
- LANL (Los Alamos National Laboratory), August 2009. "Pajarito Canyon Investigation Report, Revision 1," Los Alamos National Laboratory document LA-UR-09-4670, Los Alamos, New Mexico. (LANL 2009, 106939)
- LANL (Los Alamos National Laboratory), October 2009. "Investigation Report for Sandia Canyon," Los Alamos National Laboratory document LA-UR-09-6450, Los Alamos, New Mexico. (LANL 2009, 107453)
- LANL (Los Alamos National Laboratory), September 2010. "Toxicity Reference Value Development Methods for the Los Alamos National Laboratory," Los Alamos National Laboratory document LA-UR-10-4922, Los Alamos, New Mexico. (LANL 2010, 110623)
- LANL (Los Alamos National Laboratory), October 2015. "Threatened and Endangered Species Habitat Management Plan for Los Alamos National Laboratory," Los Alamos National Laboratory document LA-UR-15-28610, Los Alamos, New Mexico. (LANL 2015, 602156)
- LANL (Los Alamos National Laboratory), October 2016. "ECORISK Database (Release 4.0)," on CD, LA-UR-16-27520, Los Alamos National Laboratory, Los Alamos, New Mexico. (LANL 2016, 601838)

- Markwiese, J., R. Ryti, R. Mirenda, P. Newell, and J. Podolsky, November 2003. "Wildlife Screening Thresholds for Volatile Organic Chemicals," poster presented at the 24th Annual Meeting of the Society of Environmental Toxicology and Chemistry, November 9–13, 2003, Austin, Texas. (Markwiese et al. 2003, 087149)
- Menzel, M.A., J.M. Menzel, J.C. Kilgo, W.M. Ford, T.C. Carter, and J.W. Edwards, September 2003. "Bats of the Savannah River Site and Vicinity," Publication no. SRO-NERP, Savannah River Site National Environmental Research Park Program, Ashville, North Carolina. (Menzel et al. 2003, 087151)
- NCRP (National Council on Radiation Protection and Measurements), 1991. *Effects of Ionizing Radiation on Aquatic Organisms*, National Council on Radiation Protection and Measurements Report No. 109, Bethesda, Maryland. (NCRP 1991, 062803)
- Newman, M.C., M.M. Keklak, and M.S. Doggett, January 1994. "Quantifying Animal Size Effects on Toxicity: A General Approach," *Aquatic Toxicology*, Vol. 28, No. 1–2, pp. 1–12. (Newman et al. 1994, 062788)
- NMED (New Mexico Environment Department), April 2000. "Guidance for Assessing Ecological Risks Posed by Radionuclides: Screening-Level Radioecological Risk Assessment," New Mexico Environment Department, Hazardous and Radioactive Materials Bureau, Santa Fe, New Mexico. (NMED 2000, 087104)
- NMED (New Mexico Environment Department), March 2017. "Risk Assessment Guidance for Site Investigations and Remediation, Volume 2, Soil Screening Guidance for Ecological Risk Assessments," Hazardous Waste Bureau and Ground Water Quality Bureau, Santa Fe, New Mexico. (NMED 2017, 602274)
- NMED (New Mexico Environment Department), March 2017. "Risk Assessment Guidance for Site Investigations and Remediation, Volume 1, Soil Screening Guidance for Human Health Risk Assessments," Hazardous Waste Bureau and Ground Water Quality Bureau, Santa Fe, New Mexico. (NMED 2017, 602273)
- Nyhan, J.W., L.W. Hacker, T.E. Calhoun, and D.L. Young, June 1978. "Soil Survey of Los Alamos County, New Mexico," Los Alamos Scientific Laboratory report LA-6779-MS, Los Alamos, New Mexico. (Nyhan et al. 1978, 005702)
- Pastorok, R.A., M.K. Butcher, and R.D. Nielsen, September 1996. "Modeling Wildlife Exposure to Toxic Chemicals: Trends and Recent Advances," *Human and Ecological Risk Assessment*, Vol. 2, No. 3, pp. 444–480. (Pastorok et al. 1996, 062784)
- Podolsky, J.S., P.G. Newell, and O.B. Myers, 2001. "A Comparison of Toxicity Reference Values and Their Derivation Methods: Implications for Ecological Risk Assessment," Los Alamos National Laboratory presentation LA-UR-01-6503, Los Alamos National Laboratory, Los Alamos, New Mexico. (LANL 2001, 072586)
- Reed, P.B.J., September 1988. "National List of Plant Species That Occur in Wetlands: 1988 National Summary," Biological Report 88(24), U.S. Department of the Interior, Fish and Wildlife Service, Washington D.C. (Reed 1988, 600338)
- Ryti, R.T., J. Markwiese, R. Mirenda, and L. Soholt, April 2004. "Preliminary Remediation Goals for Terrestrial Wildlife," *Human and Ecological Risk Assessment*, Vol. 10, No. 2, pp. 437–450. (Ryti et al. 2004, 600901)

- Sample, B.E., G.W. Suter II, R.A. Efroymsen, and D.S. Jones, May 1998. "A Guide to the ORNL Ecotoxicological Screening Benchmarks: Background, Development, and Application, Revision 1.0," Oak Ridge National Laboratory report ORNL/TM-13615, Oak Ridge, Tennessee. (Sample et al. 1998, 062807)
- Sowls, L.K., May 1957. "Reproduction in the Audubon Cottontail in Arizona," *Journal of Mammalogy*, Vol. 38, No. 2, pp. 234-243. (Sowls 1957, 602507)
- Stahl, W.R., March 1967. "Scaling of Respiratory Variables in Mammals," *Journal of Applied Physiology*, Vol. 22, No. 3, pp. 453-460. (Stahl 1967, 063119)
- Sutherland, G.D., A.S. Harestad, and K.P. Lertzman, June 2000. "Scaling of Natal Dispersal Distances in Terrestrial Birds and Mammals," *Conservation Ecology*, Vol. 4, No. 1, online. (Sutherland et al. 2000, 073460)
- USACE (U.S. Army Corps of Engineers), October 2005. *Wetland Delineation Report*. (USACE 2005, 092220)
- Whitaker, J., 1980. Excerpted pages from *The Audubon Society Field Guide to North American Mammals*, Alfred A. Knopf, New York, New York. (Whitaker 1980, 062889)



# **Appendix A**

---

## *Derivation of Ecological Screening Levels*



## A-1.0 DERIVATION OF CHEMICAL ECOLOGICAL SCREENING LEVELS

This appendix provides the basis for media-specific ecological screening levels (ESLs) and the equations that are the foundation for calculating wildlife screening levels.

### A-1.1 Soil ESLs

The parameters used to calculate exposure of ecological receptors to contaminants in soil and food in terms of the ESL are presented in Table 3.3-1 of the document. Equation 3.4-4 is rearranged to the basic equation for the soil ESL, as shown in Equation A-1.1-1:

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times [fs_i + TF_j]} \quad \text{Equation A-1.1-1}$$

where  $ESL_{ij}$  is the soil ESL for wildlife receptor  $i$  and chemical of potential concern (COPC)  $j$  (mg/kg)  
 $TRV_{ij}$  is the toxicity reference value (no observed adverse effect level [NOAEL]) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)  
 $I_i$  is the normalized daily dietary ingestion rate for wildlife receptor  $i$  (kg-food dry wt/kg-body wt/d)  
 $fs_i$  is the fraction of soil ingested by wildlife receptor  $i$ , expressed as a fraction of the dietary intake  
 $TF_j$  is a transfer factor from soil to food for COPC  $j$  (mg/kg dry wt food per mg/kg dry wt soil)

Equations for calculating wildlife ESLs for herbivore, omnivore, insectivore, and carnivore functional groups based on Equation A-1.1-1 are shown in Equations A-1.1-2 through A-1.1-5, respectively.

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times [fs_i + fp_i \times TF_{plant,j}]} \quad \text{Equation A-1.1-2}$$

where  $ESL_{ij}$  is the soil ESL for herbivore  $i$  and COPC  $j$  (mg/kg)  
 $TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)  
 $I_i$  is the normalized daily dietary ingestion rate for herbivore  $i$  (kg-food dry wt/kg-body wt/d)  
 $fs_i$  is the fraction of soil ingested by herbivore  $i$ , expressed as a fraction of the dietary intake  
 $fp_i$  is the fraction of plants in diet for herbivore  $i$   
 $TF_{plant,j}$  is a transfer factor from soil to plants for COPC  $j$  (mg/kg dry wt plant per mg/kg dry wt soil)

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times [fs_i + fp_i \times TF_{plant,j} + fi_i \times TF_{invert,j}]} \quad \text{Equation A-1.1-3}$$

where  $ESL_{ij}$  is the soil ESL for omnivore  $i$  and COPC  $j$  (mg/kg)  
 $TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)  
 $I_i$  is the normalized daily dietary ingestion rate for omnivore  $i$  (kg-food dry wt/kg-body wt/d)  
 $fs_i$  is the fraction of soil ingested by omnivore  $i$ , expressed as a fraction of the dietary intake

$fp_i$  is the fraction of plants in diet for omnivore  $i$

$TF_{plant,j}$  is a transfer factor from soil to plants for COPC  $j$  (mg/kg dry wt plant per mg/kg dry wt soil)

$fi_i$  is the fraction of invertebrates in diet for omnivore  $i$

$TF_{invert,j}$  is a transfer factor from soil to invertebrates (mg/kg dry insect wt per mg/kg dry wt soil) or soil to flesh for COPC  $j$  (mg/kg dry wt flesh per mg/kg dry wt soil)

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times [fs_i + fi_i \times TF_{invert,j}]} \quad \text{Equation A-1.1-4}$$

where  $ESL_{ij}$  is the soil ESL for insectivore  $i$  and COPC  $j$  (mg/kg)

$TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)

$I_i$  is the normalized daily dietary ingestion rate for insectivore  $i$  (kg-food dry wt/kg-body wt/d)

$fs_i$  is the fraction of soil ingested by insectivore  $i$ , expressed as a fraction of the dietary intake

$fi_i$  is the fraction of invertebrates in diet for insectivore  $i$

$TF_{invert,j}$  is a transfer factor from soil to invertebrates for COPC  $j$  (mg/kg dry wt invertebrate per mg/kg dry wt soil)

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times [fs_i + ff_i \times TF_{flesh,j}]} \quad \text{Equation A-1.1-5}$$

where  $ESL_{ij}$  is the soil ESL for carnivore  $i$  and COPC  $j$  (mg/kg)

$TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)

$I_i$  is the normalized daily dietary ingestion rate for carnivore  $i$  (kg-food dry wt/kg-body wt/d)

$fs_i$  is the fraction of soil ingested by carnivore  $i$ , expressed as a fraction of the dietary intake

$ff_i$  is the fraction of flesh in diet for carnivore  $i$

$TF_{flesh,j}$  is a transfer factor from soil to flesh for COPC  $j$  (mg/kg dry wt flesh per mg/kg dry wt soil)

The wildlife ESL models (Equation A-1.1-1 and the functional group-specific Equations A-1.1-2 through A-1.1-5) show the ESL as proportional to the effect level. Thus, larger toxicity reference values (TRVs) result in larger ESLs, which indicate the receptor may be more tolerant of the COPC. The opposite relationship holds for the variables in the denominator of the wildlife ESL model. Thus, a receptor with higher feeding rates or one that eats more contaminated prey has a lower ESL. A receptor with higher exposure will have lower ESLs for the same TRV as a receptor with lower exposure. The wildlife lowest-effect ESLs (L-ESLs) are calculated with Equations A-1.1-2 through A-1.1-5 using the lowest observed adverse effect level (LOAEL) for the TRV term. Table A-1.1-1 summarizes the input variables for the wildlife exposure models and indicates the general sources used for these variables.

**Table A-1.1-1**  
**Summary of Variables Used in the Wildlife Soil ESL Models for Chemicals**

Variable	Source
<i>TRV</i>	Receptor and COPC-specific NOAEL values are obtained from reviewing primary literature on toxicity to ecological receptors. Values for specific receptors and COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version). The wildlife L-ESLs are calculated using the LOAEL for the TRV term.
<i>fs</i>	Receptor-specific values are provided in Table 3.3-1 of the document.
<i>I</i>	Body weight (BW) normalized food intake for wildlife receptors (see values provided in Table 3.3-1). Body weight is an implicit component of this variable. For this reason, Table 3.3-1 provides BW for each receptor. Note that intake can also be expressed as a gross daily amount (in units of kg of food ingested per day). This alternate formulation of the model requires BW to be an explicit variable.
<i>fp</i>	The fraction of plants in diet is provided in Table 3.3-1.
<i>fi</i>	The fraction of invertebrates in diet is provided in Table 3.3-1.
<i>ff</i>	The fraction of flesh in diet is provided in Table 3.3-1.
<i>TF<sub>plant</sub></i>	The transfer from soil to plants is a COPC-specific value derived from site-specific studies, other empirical literature studies, and models. Values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version). The ECORISK Database must be reviewed to determine if the soil-to-plant transfer factor (TF) accounts for all complete plant exposure pathways. In particular, many plant uptake factors do not include foliar uptake. If foliar uptake represents a complete pathway for site, then the effect of not including this pathway in the plant uptake factor should be evaluated in the site-specific uncertainty analysis.
<i>TF<sub>invert</sub></i>	The transfer from soil to invertebrates is a COPC-specific value derived from site-specific studies, other empirical literature studies, and models. Values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).
<i>TF<sub>flesh</sub></i>	<p>The transfer from soil to flesh is a COPC-specific value derived from three other factors. The first factor is a fresh weight feed to muscle TF (<math>TF_{beef}</math>) derived from studies of beef cattle. The second factor is the maximum of either the moisture content– (MC–) adjusted dry weight <math>TF_{plant}</math> or the MC-adjusted dry weight <math>TF_{invert}</math>. This TF term represents the prey with the most contaminated diet. The two TFs are multiplied by a food-ingestion rate. This rate is based on a composite prey species value developed from the four potential mammalian prey species (robin, deer mouse, cottontail, and shrew). The highest food and soil intake rates among these four potential prey species were used to represent the composite prey species in the equation below: Thus,</p> $TF_{flesh} = TF_{beef} \times (I_{food} \times \text{maximum of } [TF_{plant} \times (1 - MC_{plant}) \text{ or } TF_{invert} \times (1 - MC_{invert})] + I_{soil}) / (1 - MC_{flesh}).$ <p>General parameters and values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).</p>

### A-1.2 Burrow Air ESLs (Vapor-Phase Contaminants Only)

Quantitative evaluations of ecological risk do not typically include the inhalation pathway because ingestion-related exposure is relatively more important for most chemicals. However, air exposure is potentially a significant exposure pathway for burrowing mammals at some solid waste management units and areas of concern at Los Alamos National Laboratory (LANL or the Laboratory). Gaseous or otherwise airborne contaminants can build up in burrows because the potential for dilution with the atmosphere is much more limited compared with surface conditions. Section 3.4.2 of the document describes ecological parameters affecting air intake for the gopher.

The gopher's ESL for inhalation is expressed in Equation A-1.2-1:

$$ESL_j = \frac{TRV_j \times BW}{I_{air}} \quad \text{Equation A-1.2-1}$$

where  $ESL_j$  is the soil ESL for burrow animal and COPC  $j$  (mg/m<sup>3</sup>)

$TRV_j$  is the NOAEL for burrow animal inhalation and COPC  $j$  (mg-COPC/kg-body wt/d)

$BW$  is the body weight for burrow animal (kg)

$I_{air}$  is the daily inhalation rate for the pocket gopher (m<sup>3</sup>/d)

The wildlife L-ESLs are calculated using the LOAEL rather than the NOAEL for the TRV in the same equation.

### A-1.3 Sediment ESLs for the Aquatic Community

The National Oceanic and Atmospheric Administration (NOAA) Screening Quick Reference Tables (SQuiRT) (Buchman 2008, 206414); <http://response.restoration.noaa.gov/cpr/sediment/squirt/squirt.html>) represents a nationally recognized compendium of ecological effects values in soil, sediment, and water. In selecting sediment ESLs that are protective of the aquatic community, it is first determined if a benchmark is available in the SQuiRT. Within SQuiRT, benchmarks are evaluated in the order presented in Figure A-1.3-1, based on the rigor and comprehensiveness of the data source. Preference was given to benchmarks based on publication date (the more recent are assumed to reflect the broader extent of scientific knowledge), chronic direct exposure, and nonlethal endpoint studies designed to be protective of sensitive species. Table A-1.3-1 lists the definitions of the sediment effect concentrations.

Sediment benchmarks from MacDonald et al. (2000, 205266) were selected as the first potential source of sediment ESLs protective of the aquatic community. For some contaminants, MacDonald et al. (2000, 205266) published two consensus-based benchmarks (threshold effect concentration [TEC] and probable effect concentration [PEC]) for each contaminant. The predictive ability of these benchmarks was numerically evaluated for accuracy using field data.

If a TEC and/or PEC was not available from MacDonald et al. (2000, 205266), the next potential source for freshwater sediment benchmarks was Persuad et al. (1993, 205250), which form the basis for sediment screening values used by the Canadian Council of Ministers of the Environment (CCME) (<http://st-ts.ccme.ca/>). Some of the CCME values have been periodically updated since they were first published in the early 1990s. The sediment ESL protective of the aquatic community is based on the CCME concentrations (LELs) below which concentrations are not toxic to the majority of benthic organisms. The L-ESL is based on the concentrations (severe effect level [SELs]) that are expected to be detrimental to the majority of benthic species.




Figure A-1.3-1 Sediment ESL selection process for the aquatic sediment community

**Table A-1.3-1**  
**Definitions of Terms Associated with Sediment ESLs for the Aquatic Community**

Term	Definition	Description
CCME	Canadian Council and Ministry of the Environment	Canadian environmental standards. Based on Persuad et al. (1993, 205250) and periodically updated (e.g., <a href="http://www.ccme.ca/publications/cegg_rcqe.html">www.ccme.ca/publications/cegg_rcqe.html</a> ).
CSL	Cleanup screening level	Concentration below which only minor adverse effects would occur and above which more significant adverse effects are expected.
EqP	Equilibrium partitioning	Di Toro's method (Di Toro 1985, 062876) to calculate sediment ESLs based on chemicals toxicity in water and calculated partitioning between sediment and water.
ERM	Effect range median	Concentration of a chemical in sediment above which effects are frequently or always observed or predicted among most species.
LEL	Lowest effect level	Concentration below which majority of benthic organisms tolerate.
MPC	Maximum permissible concentration	Concentration above which the risk of adverse effects is unacceptable to sedimentary ecosystems.
NC	Negligible concentration	Concentration below which potential for adverse effects are negligible.
PEC	Probable effects concentration	Consensus based concentration in sediment to which a plant or animal is directly exposed that is likely to cause an adverse effect.
SEL	Severe effect level	Concentration expected to be detrimental to the majority of benthic species.
SQS	Sediment quality standard	Concentration resulting in no adverse effects, including no acute or chronic adverse effects on biological resources.
TEC	Threshold effect concentration	Consensus-based concentration of a contaminant above which some effect will be produced and below which an effect will not be produced.
TEL	Threshold effect level	Consensus-based concentration of a contaminant below which adverse biological effects are expected to occur rarely.
TL	Threshold level	Dutch sediment standard for acceptable level of chemical in bedded sediment environment.
UET	Upper effects threshold	Concentration above which adverse impacts on the benthic community are always expected.

If a CCME value was not available in SQuiRT, the Assessment and Remediation of Contaminated Sediments (ARCS) program in the Great Lakes (Ingersoll et al. 1996, 062873) was the next potential source of sediment benchmarks. The ARCS program has sponsored numerous investigations using the amphipod *Hyalella azteca* and the midge *Chironomus riparius* in sediment bioassays. These results, along with those from other freshwater areas, were used to generate a TEL and ERM (Table A-1.3-1). The ERM values from ARCS represent studies on freshwater species and should not be confused with the marine ERM values. Marine ERM values are not used as the basis for ESLs. The next potential source of sediment benchmarks is the Dutch<sup>1</sup> sediment TL that may be used as a no observed effect concentration (NOEC) for the ESL. The TL represents concentrations that delineate the threshold below which effects are not expected. The last benchmark selected in SQuiRT is the UET, a sediment toxicity value put forth by NOAA that corresponds to a concentration above which adverse impacts on the benthic community are always expected. A UET is not suitable for a no-effect screening level but can be used for

<sup>1</sup> As reported in the SQuiRT, Dutch standards are "E.M.J. Verbruggen, R. Posthumus and A.P. van Wezel, 2001. Ecotoxicological serious risk concentrations for soil, sediment, and (ground)water: updated proposal for first series of compounds. Nat. Inst. Public Health and the Env., and subsequent updates as published elsewhere."



an LEL; consequently, the ESL is derived by taking one-tenth of the UET and the L-ESL is equal to the UET.

If sediment toxicity values were not available in SQUIRT, Michelsen (2003, 215128) was consulted. Michelsen (2003, 215128) compiled freshwater sediment toxicity results intended for use in the state of Washington, and these benchmarks are likely representative of potential for adverse effects in any freshwater stream, including those found at the Laboratory. No other neighboring state has compiled freshwater sediment toxicity values. Sediment quality values were generated using four bioassay endpoints: *H. azteca* 10-d mortality, *Chironomus* 10-d mortality, *Chironomus* 10-d growth, and Microtox 15-min luminescence bioassays. Michelsen (2003, 215128) compiles two relevant sediment benchmarks: the SQS and the CSL. The SQS corresponds to the concentration that will result in no adverse effects (i.e., equivalent to a NOAEL), including no acute or chronic adverse effects on biological resources, and the CSL corresponds to concentration below which only minor adverse effects would occur and above which more significant adverse effects are expected (i.e., equivalent to a LOAEL).

If benchmarks are not available from any of the preferred sources, values used in the Netherlands (Crommentuijn et al. 2000, 205264; Crommentuijn et al. 2000, 205265) may be considered. These values are designated as the negligible concentration (NC), which is equivalent to a NOAEL, and the maximum permissible concentration (MPC), which is equivalent to a LOAEL. If Crommentuijn et al. values are not available, the equilibrium partitioning (EqP) values are used. If the EqP is used as the ESL, then the ESL is multiplied by a factor of 10 to derive the L-ESL.

Because the sediment ESLs are broadly representative of the adverse effects of contaminants on the aquatic community, they are applied to both aquatic plants and aquatic invertebrates. The sediment ESLs described here are broadly protective of the aquatic environment.

In addition to selecting sediment ESLs that are protective of the aquatic community, the model shown in Equation A-1.3-1 is based on the transfer of contamination from sediment to invertebrates and the subsequent ingestion of the insects (by an insectivore) as contaminated food. The insectivores in this model are the bat and the swallow, and the exposure information for these receptors is provided in Table 3.3-1 of the document. Contaminant transfer to higher level carnivores is not accounted for by these ESLs and should be addressed in the uncertainty analysis.

$$ESL_{ij} = \frac{TRV_{ij}}{I_i \times f_{ij} \times TF_{invert,j}} \quad \text{Equation A-1.3-1}$$

where  $ESL_{ij}$  is the sediment ESL for insectivore  $i$  and COPC  $j$  (mg/kg)

$TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  (mg-COPC/kg-body wt/d)

$I_i$  is the normalized daily dietary ingestion rate for insectivore  $i$  (kg-food dry wt/kg-body wt/d)

$f_{ij}$  is the fraction of invertebrates in diet for insectivore  $i$

$TF_{invert,j}$  is a transfer factor from sediment to invertebrate for COPC  $j$  (mg/kg dry invertebrate wt per mg/kg dry sediment wt)

The aerial insectivore sediment L-ESLs are calculated with Equation A-1.3-2 using the LOAEL for the TRV term.

#### A-1.4 Water ESLs for the Aquatic Community

This section describes the selection process for water ESLs or benchmarks protective of the aquatic community. Values reported as chronic are used for the ESLs, and those reported as acute are used for L-ESLs. In some cases, study conditions did not match or produce data directly comparable with chronic or acute benchmarks. In these instances, and when the difference between the chronic and acute was more than tenfold, uncertainty factors were applied to the lowest acceptable study data in the order of preferred sources to obtain water benchmarks (Figure A-1.4-1). Uncertainty factors were used to convert acute values to chronic values and, conversely, when only a chronic value was available, uncertainty factors were applied to derive the acute value. Table A-1.4-1 provides definitions for terms used to develop water ESLs that are protective of the aquatic community.

For conversion of chronic values to acute, an uncertainty factor (UF) of 10 was applied. This value is consistent with the geometric mean (7.6) of the acute-chronic ratios used by the U.S. Environmental Protection Agency (EPA) to develop AWQC for primary pollutant metals (EPA 2009, 109328). The UF of 10 is within the range of 1–10 recommended by EPA in the Great Lakes Water Quality Initiative (60 Federal Register 15366, “Final Water Quality Guidance for the Great Lakes System, Final Rule”) and is supported by EPA Region 10 guidance (EPA 1997, 215127).

Values are selected from four tiers of data sources, with Tier 1 the most preferred data source. The selection process followed is shown in Figure A-1.4-1, and the data sources are as follows:


1. “Standards for Interstate and Intrastate Streams,” 20.6.4.900 of the New Mexico Administrative Code (20.6.4.900 NMAC)
2. Ambient water-quality criteria (AWQC) set forth by EPA (2009, 109328)
3. NOAA SQUIRT (Buchman 2008, 206414);  
(<http://response.restoration.noaa.gov/cpr/sediment/squirt/squirt.html>)
4. Other sources (see LANL 2016, 601838, or latest version)

Water ESLs are selected utilizing water-quality criteria (WQC) or benchmarks in the order presented above. For example, if a 20.6.4.900 NMAC criterion is available for a given constituent, then it is selected as the most relevant screening value. If no 20.6.4.900 NMAC criterion is available, the AWQC are evaluated as the next tier. Justification for selecting the above order is provided in greater detail in 20.6.4.900 NMAC and in various EPA documents (60 Federal Register 15366, “Final Water Quality Guidance for the Great Lakes System, Final Rule”; EPA 2009, 109328).

The national AWQC are developed by EPA’s Office of Water under the Clean Water Act, Section 304 (EPA 2009, 109328). New Mexico has developed similar criteria for “high quality coldwater fisheries,” as listed in 20.6.4.900 NMAC. The development of AWQC is outlined in EPA guidance (60 Federal Register 15366, “Final Water Quality Guidance for the Great Lakes System, Final Rule”). Metals are often water hardness–dependent and should be adjusted for site-specific conditions (see EPA guidance [EPA 2009, 109328], and 20.6.4.900 NMAC for explanations and delineation of methods because methods require analyte-specific information).

If New Mexico WQC or national AWQC are not available, values from SQUIRT (Buchman 2008, 206414; <http://response.restoration.noaa.gov/cpr/sediment/squirt/squirt.html>) should be reviewed for applicability. In some cases, more than one chronic value is presented for a chemical in SQUIRT. In such instances, the priority is to use ECOTOX thresholds or Tier II secondary chronic values (<http://www.esd.ornl.gov/programs/ecorisk/tools.html>), followed by values from Canada or New Zealand, with the EPA Region 5 ecological screening values (<https://www3.epa.gov/region5/waste/cars/esl.htm>) as

the last option. If toxicity information is not available in SQUIRT, other sources are consulted for water benchmarks.



**Figure A-1.4-1 Water ESL selection process for the aquatic community**

**Table A-1.4-1**  
**Definitions of Terms Associated with Water ESLs for the Aquatic Community**

Term	Definition	Description
AWQC	Ambient water-quality criteria	U.S. national recommended water-quality criteria broadly protective of aquatic species.
CCC	Criterion continuous concentration	The concentration in water expected to be protective of 95% of aquatic species over chronic exposure.
CMC	Criterion maximum concentration	The concentration in water that represents a low-level effect on the fifth percentile genus, applied as a limit on the short-term average concentration in the environment. Both the acute and chronic criteria are values not to be exceeded more than once in 3 yr. In other words, the criteria specify a magnitude, duration, and frequency to be met to protect aquatic life.
WQC	Water-quality criteria	State acute and chronic stream standards broadly protective of aquatic species.

In addition to selecting water ESLs that are protective of the aquatic community, the wildlife ESL is calculated as the oral daily dose for a COPC from water as the numerator in the hazard quotient (HQ) calculation (Equation 3.1-1 in the document) and setting the HQ to 1, then rearranging to solve for the wildlife water ESL, yielding the following equation:

$$ESL_{ij} = \frac{1000 \times TRV_{ij}}{I_i} \quad \text{Equation A-1.4-1}$$

where  $ESL_{ij}$  is the water ESL for wildlife species  $i$  and COPC  $j$  ( $\mu\text{g/L}$ )

1000 is the number of  $\mu\text{g}$  per  $\text{mg}$

$TRV_{ij}$  is the toxicity reference value (NOAEL) for wildlife receptor  $i$  and COPC  $j$  ( $\text{mg-COPC/kg-body wt/d}$ )

$I_i$  is the daily water ingestion rate for wildlife species  $i$  ( $\text{L of water/kg body wt/d}$ )

The minimum water ESL for a given chemical is the lowest of the values available for the aquatic community or wildlife. The parameter values are summarized in Table 3.3-1. The wildlife water L-ESLs are calculated with Equation A-1.4-1 using the LOAEL for the TRV term.

## A-2.0 BASIS AND DERIVATION OF RADIONUCLIDE ESLS

The derivation of ESLs for radionuclides is discussed in section 3.5 of the document. The methods followed for radionuclide ESL development are consistent with U.S. Department of Energy (DOE) guidance (DOE 2002, 085637). The equations and assumptions underlying radiological ESL development for soil, sediment, and water are presented in the following sections.

## A-2.1 Soil ESLs

### A-2.1.1 Radionuclide Concentrations in Biota

#### Plants and Invertebrates

The internal dose to plants is calculated by estimating the internal concentration or body burden and the internal dose conversion factor (DCF) (as described below). The internal plant concentration is calculated as

$$C_{plant,j} = C_{soil,j} \times TF_{plant,j} \quad \text{Equation A-2.1-1}$$

where  $C_{plant,j}$  is the internal concentration of radionuclide  $j$  in plants (pCi/g fresh wt)

$C_{soil,j}$  is the soil concentration of radionuclide  $j$  (pCi/g dry soil)

$TF_{plant,j}$  is the soil to plant transfer factor for radionuclide  $j$  (pCi/g fresh wt plant per pCi/g dry wt soil)

The same equation is used to calculate dose to soil-dwelling invertebrates, with a soil-to-invertebrate TF ( $TF_{invert}$ ) substituted in place of the soil to plant TF. Thus, the internal concentration in invertebrates is

$$C_{invert,j} = C_{soil,j} \times TF_{invert,j} \quad \text{Equation A-2.1-2}$$

where  $C_{invert,j}$  is the internal concentration of radionuclide  $j$  in invertebrates (pCi/g fresh wt)

$C_{soil,j}$  is the soil concentration of radionuclide  $j$  (pCi/g dry soil)

$TF_{invert,j}$  is the soil to invertebrate transfer factor for radionuclide  $j$  (pCi/g fresh wt invertebrate per pCi/g dry wt soil)

Values and references for TFs are presented in the ECORISK Database (LANL 2015, 600921, or latest version). When values are not available in the literature, a default value of 1 is used.

#### Wildlife

The internal dose to wildlife is calculated by multiplying the effective energy of a radionuclide by the body burden of that radionuclide in an organism. Body burden is a measure of the accumulation of a radionuclide in an organism through ingestion. The body burden calculation is presented in Equation A-2.1-3.

$$C_{wildlife,j} = C_{soil,j} \times [I_{soil} + TF_{food,j} \times I_{food}] \times TF_{blood,j} \times R_{t,j} \quad \text{Equation A-2.1-3}$$

where  $C_{wildlife,j}$  is the body burden of radionuclide  $j$  in a wildlife species (pCi/g)

$C_{soil,j}$  is the concentration of radionuclide  $j$  in soil (pCi/g)

$I_{soil}$  is the normalized daily soil ingestion rate (g of soil/g of body wt/d)

$TF_{food,j}$  is the soil to food transfer factor for radionuclide  $j$  (pCi/g fresh wt food per pCi/g dry wt soil)

$I_{food}$  is the normalized daily dietary ingestion rate (g of food [fresh wt]/g of body wt/d)

$TF_{blood,j}$  is the food to blood transfer factor (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time of radionuclide  $j$  in the organism (d)

Dietary and soil ingestion rates for each receptor are presented in Table 3.3-1 of the document. Values and supporting references for all TFs used are provided in the ECORISK Database (LANL 2016, 601838, or latest version). The retention time,  $R_t$ , is an equilibrium model that assumes the activity concentration of a radionuclide reaches steady state in an organism over time, depending upon the rate of radiological decay and metabolic elimination of the element from the organisms body. This value (modified from Baker and Soldat 1992, 062801) is calculated as

$$R_t = (1 - e^{-\lambda T_c}) / \lambda \quad \text{Equation A-2.1-4}$$

where  $R_t$  = retention time of radionuclide in the organism (d)

$$\lambda = \lambda_r + \lambda_b$$

$$\lambda_r = \ln(2) / T_r, \text{ where } T_r \text{ is the radiological half-life of the radionuclide (d)}$$

$$\lambda_b = \ln(2) / T_b, \text{ where } T_b \text{ is the biological half-life of the radionuclide (d)}$$

$$T_c = \text{exposure duration, or the average life-span of the receptor (d)}$$

Values and references for all of the parameters used in calculating  $R_t$  for each radionuclide are presented in the ECORISK Database (LANL 2016, 601838, or latest version).

#### A-2.1.2 Internal DCF

Uranium, plutonium, americium, thorium, and radium have radioactive daughters. For screening purposes, the sum of average energies per disintegration for the decay chains of all radioactive daughters for any given isotope is used. This method provides an overestimate of exposure because the lifetime of many of the biota of interest is short compared with the time for the build-up of progeny. The energy deposition for radionuclides is given in the units of million electron volts (MeV) per disintegration. To calculate internal dose, it is necessary to convert MeV/disintegration to rad/d per pCi/g, as internal radioactivity is measured in pCi/g. A combined conversion factor of  $5.11 \times 10^{-5}$  (disintegrations  $\times$  g  $\times$  rad)/(MeV  $\times$  pCi  $\times$  d) is applied to convert MeV/disintegration to rad/d per pCi/g. This conversion factor is derived in Equation A-2.1-5.

#### Equation A-2.1-5

$$5.11 \times 10^{-5} \frac{\text{disintegrations} \times \text{g} \times \text{rad}}{\text{MeV} \times \text{pCi} \times \text{day}} = 1.6 \times 10^{-6} \frac{\text{ergs}}{\text{MeV}} \times \frac{\text{rad}}{100 \text{ ergs/g}} \times \frac{\text{disintegration}}{27.03 \text{ pCi} \cdot \text{s}} \times 8.64 \times 10^4 \frac{\text{s}}{\text{day}}$$

where disintegrations is spontaneous disintegration of a radioactive substance along with the emission of ionizing radiation

erg is a unit of energy equal to a force of 1 dyne acting over 1 cm (equal to  $0.642 \times 10^{12}$  eV)

MeV is million electron volts

The relative biological effectiveness of alpha-particle emissions is about 20 times that of beta or gamma emissions, so the fraction of energy deposition from alpha particles must be taken into account in calculating the internal dose (IAEA 1992, 062802). Thus, the internal DCF to any organism from radionuclide  $j$  can be calculated as follows:

$$DCF_{int,j} = CF_i \times (f_a \times 20 + [1 - f_a]) E_j \quad \text{Equation A-2.1-6}$$

where  $DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$CF_i$  is the conversion factor between energy per disintegration and rad/d  
[value is  $5.11 \times 10^{-5}$  (disintegrations  $\times$  g  $\times$  rad)/(MeV  $\times$  pCi  $\times$  d)]

$f_a$  is the fraction of disintegrations that are alpha particles

$E_j$  is the sum of deposited energies for radionuclide  $j$  and its daughter products  
(units are MeV/disintegration)

### A-2.1.3 External Dose to Biota

The external dose to biota is the dose an organism receives from being exposed to contaminated soil and varies with several factors, including the size of the organism, the distance of the organism from the contaminated media, the geometry of the contamination within the contaminated media, and the type of radiological decay (Baker and Soldat 1992, 062801; EPA 1993, 062798). Several simplifying assumptions are made in estimating this dose. First, as indicated by the conceptual site model diagram (Figure 3.4-1), only external exposure from gamma-emitting radionuclides is considered. The basis for eliminating alpha and beta external exposure is that these particles are primarily a hazard when committed internally and have very low dose consequence when they are external to the organism (Higley and Kuperman 1996, 062804). To emphasize the protective nature of the screening levels, “worst case” assumptions are made on the size of the organism, the geometry of the contaminated source, and the location of the receptor relative to the contaminated source. Dose coefficients developed for exposure to soil assume only 180-degree exposure to the contaminated source and thus are inappropriate for modeling exposure to organisms dwelling in soil. For soil invertebrates and burrowing mammals, external dose coefficients based upon immersion in water contaminated to an infinite depth are used (EPA 1993, 062798) to provide a conservative estimate of external dose because dose resulting from immersion in contaminated soil would be less than dose from water from the higher density of soil. For terrestrial organisms living on or above the soil surface, dose coefficients for exposure to soil contaminated to an infinite depth is used (EPA 1993, 062798). As larger organisms receive a greater proportion of the external dose, the standard man is used as a default organism to conservatively represent exposure to all terrestrial receptors living on or above the soil surface. Thus, external DCF is modeled by the following equations:

Invertebrates and burrowing mammals,

$$DCF_{ext,j} = DC_{water,skin,j} \times CF_{e,w} \quad \text{Equation A-2.1-7a}$$

Terrestrial receptors on or above the soil surface,

$$DCF_{ext,j} = DC_{soil,skin,j} \times CF_{e,s} \quad \text{Equation A-2.1-7b}$$

where  $DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil)

$DC_{water,skin,j}$  is the dose coefficient for skin exposed to water contaminated to an infinite depth with radionuclide  $j$  (EPA 1993, 062798)

$CF_{e,w}$  is the conversion factor from Sv/s per Bq/m<sup>3</sup> to rad/d per pCi/g for an organism immersed in water (value is  $3.2 \times 10^{11}$ ; Equation A-2.1-8)

$DC_{soil,skin,j}$  is the dose coefficient for skin exposed to soil contaminated to an infinite depth with radionuclide  $j$  (EPA 1993, 062798)

$CF_{e,s}$  is the conversion factor from Sv/s per Bq/m<sup>3</sup> to rad/d per pCi/g for an organism on the soil surface (value is  $5.11 \times 10^{11}$ ; Equation A-2.1-9)

$CF_{e,w}$  assumes a water density of  $1.0 \times 10^3 \text{ kg/m}^3$  and is derived in the following equation:

$$CF_{e,w} = 10^3 \frac{\text{kg}}{\text{m}^3} \times 10^3 \frac{\text{g}}{\text{kg}} \times 100 \frac{\text{rad}}{\text{Sv}} \times \frac{\text{Bq}}{27.03 \text{ pCi}} \times 86400 \frac{\text{s}}{\text{d}} \quad \text{Equation A-2.1-8}$$

$CF_{e,s}$  assumes a soil density of  $1.6 \times 10^3 \text{ kg/m}^3$  and is derived in the following equation:

$$CF_{e,s} = 1.6 \times 10^3 \frac{\text{kg}}{\text{m}^3} \times 10^3 \frac{\text{g}}{\text{kg}} \times 100 \frac{\text{rad}}{\text{Sv}} \times \frac{\text{Bq}}{27.03 \text{ pCi}} \times 86400 \frac{\text{s}}{\text{d}} \quad \text{Equation A-2.1-9}$$

#### A-2.1.4 Calculations of Soil ESLs

The soil ESL is defined as the soil concentration of a given radionuclide that yields a combined internal and external dose rate of 0.1 rad/d to any organism.

For terrestrial plants, the ESL equation is written as

$$ESL = \frac{\text{Dose Limit}}{TF_{\text{plant},j} \times DCF_{\text{int},j} + DCF_{\text{ext},j}} \quad \text{Equation A-2.1-10}$$

where *Dose Limit* is 0.1 rad/d

$TF_{\text{plant},j}$  is the soil to plant transfer factor for radionuclide  $j$  (pCi/g fresh plant per pCi/g dry soil wt)

$DCF_{\text{int},j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g-fresh tissue)

$DCF_{\text{ext},j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil, assuming 360-degree exposure)

For terrestrial invertebrate receptors, the ESL equation is written as

$$ESL = \frac{\text{Dose Limit}}{TF_{\text{invert},j} \times DCF_{\text{int},j} + DCF_{\text{ext},j}} \quad \text{Equation A-2.1-11}$$

where *Dose Limit* is 0.1 rad/d

$TF_{\text{invert},j}$  is the soil to invertebrate transfer factor for radionuclide  $j$  (pCi/g fresh invertebrate per pCi/g dry soil wt)

$DCF_{\text{int},j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$DCF_{\text{ext},j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil, assuming 360-degree exposure)

For terrestrial herbivores, the ESL equation is written as

$$ESL = \frac{\text{Dose Limit}}{[I_{\text{soil},i} + TF_{\text{plant},j} \times I_{\text{plant},i}] \times TF_{\text{blood},j} \times R_{t,j} \times DCF_{\text{int},j} + DCF_{\text{ext},j}} \quad \text{Equation A-2.1-12}$$

where *Dose Limit* is 0.1 rad/d

$I_{\text{soil},i}$  is the normalized daily soil ingestion rate for organism  $i$  (g of dry soil/g of body wt/d)

$TF_{\text{plant},j}$  is the soil to plant transfer factor for radionuclide  $j$  (pCi/g fresh plant per pCi/g dry soil)

$I_{\text{plant},i}$  is the normalized daily plant ingestion rate for organism  $i$  (g of plant-fresh wt/g of body wt/d)



$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time for radionuclide  $j$  (d)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil assuming 180- or 360-degree exposure)

For terrestrial receptors with a 100% invertebrate diet, the ESL equation is written as

$$ESL = \frac{Dose\ Limit}{[I_{soil,i} + TF_{invert,j} \times I_{invert,i}] \times TF_{blood,j} \times R_{t,j} \times DCF_{int,j} + DCF_{ext,j}} \quad \text{Equation A-2.1-13}$$

where *Dose Limit* is 0.1 rad/d

$I_{soil,i}$  is the normalized daily soil ingestion rate for organism  $i$  (g of dry soil/g of body wt/d)

$TF_{invert,j}$  is the soil to invertebrate transfer factor for radionuclide  $j$  (pCi/g fresh invertebrate per pCi/g dry soil)

$I_{invert,i}$  is the normalized daily invertebrate ingestion rate for organism  $i$  (g of invertebrate-fresh wt/g of body wt/d)

$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time for radionuclide  $j$  (d)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil assuming 180- or 360-degree exposure)

For terrestrial omnivores feeding upon both plants and invertebrates, the ESL equation is written as

**Equation A-2.1-14**

$$ESL = \frac{Dose\ Limit}{[I_{soil,i} + TF_{plant,j} \times I_{plant,i} + TF_{invert,j} \times I_{invert,i}] \times TF_{blood,j} \times R_{t,j} \times DCF_{int,j} + DCF_{ext,j}}$$

where *Dose Limit* is 0.1 rad/d

$I_{soil,i}$  is the normalized daily soil ingestion rate for organism  $i$  (g of soil/g of body wt/d)

$TF_{plant,j}$  is the soil to plant transfer factor for radionuclide  $j$  (pCi/g fresh plant per pCi/g dry soil)

$I_{plant,i}$  is the normalized daily plant ingestion rate for organism  $i$  (g of plant-fresh wt/g of body wt/d)

$TF_{invert,j}$  is the soil to invertebrate transfer factor for radionuclide  $j$  (pCi/g fresh invertebrate per pCi/g dry soil)

$I_{invert,i}$  is the normalized daily invertebrate ingestion rate for organism  $i$  (g of invertebrate-fresh wt/g of body wt/d)

$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time for radionuclide  $j$  (d)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil assuming 180- or 360-degree exposure)

For terrestrial carnivores, the ESL equation is written as

$$ESL = \frac{Dose\ Limit}{[I_{soil,i} + TF_{flesh,j} \times I_{flesh,i}] \times TF_{blood,j} \times R_{t,j} \times DCF_{int,j} + DCF_{ext,j}} \quad \text{Equation A-2.1-15}$$

where  $Dose\ Limit$  is 0.1 rad/d

$I_{soil,i}$  is the normalized daily soil ingestion rate for organism  $i$  (g of soil/g of body wt/d)

$TF_{flesh,j}$  is the soil to flesh transfer factor for radionuclide  $j$  (pCi/g flesh-fresh wt per pCi/g dry soil)

$I_{flesh,i}$  is the normalized daily flesh ingestion rate for organism  $i$  (g of flesh-fresh wt/g of body wt/d)

$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time for radionuclide  $j$  (d)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry soil assuming 180- or 360-degree exposure)

The soil radiological L-ESLs are calculated with Equations A-2.1-10 through A-2.1-15 using 1 rad/d as the dose limit. Table A-2.1-1 summarizes the variables used to calculate soil ESLs for radionuclides.

**Table A-2.1-1**  
**Summary of Variables Used in Soil ESL Calculations for Radionuclides**

Variable	Source
$I_{soil}$	BW-normalized soil ingestion rate for wildlife receptors (fresh food intake $\times$ fraction of soil in diet) (LANL 2016, 601838, or latest version).
$I_{plant}$	BW-normalized plant ingestion rate for wildlife receptors (fresh food intake $\times$ fraction of plants in diet) (LANL 2016, 601838, or latest version).
$I_{invert}$	BW-normalized invertebrate ingestion rate for wildlife receptors (fresh food intake $\times$ fraction of invertebrates in diet) (LANL 2016, 601838, or latest version).
$I_{flesh}$	BW-normalized flesh ingestion rate for wildlife receptors (fresh food intake $\times$ fraction of flesh in diet) (LANL 2016, 601838, or latest version).
$R_t$	The retention time of a radionuclide in an organism. This is a COPC-specific value based upon both the radiological decay constant and the biological removal rate constant for a given radionuclide. See Equation A-2.1-4 for calculation of this variable.
$TF_{blood}$	The TF from food to blood is a COPC-specific value derived from site-specific studies, other empirical literature studies, and/or models. The TF is based on the beef TF ( $TF_{beef}$ ) in pCi/g fresh beef per pCi COPC/d and the food ingestion rate. Values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).
$TF_{plant}$	The TF from soil to plants is a COPC-specific value derived from site-specific studies, other empirical literature studies, and/or models. Values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).
$TF_{invert}$	The TF from soil to invertebrates is a COPC-specific value derived from site-specific studies, other empirical literature studies, and/or models. Values for specific COPCs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).

Table A-2.1-1 (continued)

Variable	Source
$TF_{flesh}$	The TF from soil to flesh is a COPC-specific value that may come directly from measurement or may be derived from three other factors. The first factor is a fresh weight feed to muscle TF ( $TF_{beef}$ ) derived from studies of beef cattle. The second factor is the maximum of either the MC-adjusted dry weight $TF_{plant}$ or the MC-adjusted dry weight $TF_{invert}$ . This TF term represents the prey with the most contaminated diet. The two TFs are multiplied by a food ingestion rate. This rate is based on a composite prey species value developed from the four potential mammalian prey species (robin, deer mouse, cottontail, and shrew). The highest food and soil intake rates among these four potential prey species were used to represent the composite prey species in the equation below. Thus, $TF_{flesh} = TF_{beef} \times (I_{food} \times \text{maximum of } [TF_{plant} \times (1-MC_{plant}) \text{ or } TF_{invert} \times (1-MC_{invert})] + I_{soil}) / (1-MC_{flesh})$ Values for specific COPCs are provided in the Ecorisk Database (LANL 2016, 601838, or latest version).
$f_a$	The fraction of energy deposition in an organism from alpha-particle absorption.
$DCF_{int}$	The internal DCF for a specific radionuclide. This factor considers the conversion of units of deposited energy from MeV/disintegration to rad/d per pCi/g BW and accounts for the increased biological effectiveness of alpha-particle deposition over beta or gamma deposition (Equation A-2.1-6).
$DCF_{ext}$	The external DCF for a specific radionuclide. This factor applies only to gamma emitters and is media- and COPC-specific. It contains the unit conversion factor rad/d per pCi/g dry soil and is based on assuming 180- or 360-degree exposure.

## A-2.2 Sediment ESLs

### A-2.2.1 Radionuclide Concentrations in Biota

For organisms living in or on sediment (algae, daphnid, snail, and bottom-feeding fish), internal concentration of any radionuclide is modeled as part of the water ESL development described in section 3.4.4 of the document (Baker and Soldat 1992, 062801). Thus, paired data for water and sediment are needed to assess the radionuclide dose.

For terrestrial receptors ingesting sediment invertebrates, however, the internal dose from invertebrate prey is explicitly considered in the sediment calculation, which is consistent with DOE standard DOE-STD-1153-2002 (DOE 2002, 085637). Assuming the bat and swallow are eating only flying insects that have emerged from aquatic systems (an extremely conservative assumption), the body burden for these receptors is calculated as

$$C_{organism,j} = C_{sediment,j} \times BCF_{invert,j} \times I_{food,i} \times TF_{blood,j} \times R_{t,j} \quad \text{Equation A-2.2-1}$$

where  $C_{organism,j}$  is the internal concentration of radionuclide  $j$  (pCi/g of organism)

$C_{sediment,j}$  is the concentration of radionuclide  $j$  in sediment (pCi/g)

$BCF_{invert,j}$  is the sediment to invertebrate bioconcentration factor for radionuclide  $j$  (g of invertebrate-fresh wt/g dry sediment)

$I_{food,i}$  is the normalized daily dietary ingestion rate of organism  $i$  (g of food [dry wt]/g of body wt/d)

$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time of radionuclide  $j$  in the organism (d)

Values and references for the TFs and bioconcentration factors (BCFs) are provided in the ECORISK Database (LANL 2016, 601838, or latest version).

### A-2.2.2 DCFs

For aquatic receptors, internal DCFs are identical to those used for terrestrial receptors. For organisms that reside in, on, or near sediment (algae, snail, and fish), external dose is estimated the same as for terrestrial receptors living in or on soil. As with terrestrial receptors, external dose is deemed significant only for gamma emitters.

Internal dose to terrestrial receptors from sediment is assumed to come entirely from uptake from the food chain. Because these receptors have limited contact with sediment, it is assumed the external dose to terrestrial receptors is insignificant and all dose received is internal.

### A-2.2.3 Calculations of Sediment ESLs

The sediment ESL is defined as the sediment concentration of a given radionuclide that yields a combined internal and external dose rate of 0.1 rad/d to a particular receptor. For receptors that spend at least part of their lives in close association with sediment, the ESL equation is

$$ESL = \frac{Dose\ Limit}{BCF_i \times DCF_{int,i} + DCF_{ext,j}} \quad \text{Equation A-2.2-2}$$

where *Dose Limit* is 0.1 rad/d

$BCF_i$  is the bioconcentration factor for sediment for organism  $i$  (pCi/g-fresh wt per pCi-COPC/g dry sediment)

$DCF_{int,i}$  is the internal dose conversion factor for sediment and is set to zero for sediment as internal dose is modeled via water exposures

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide  $j$  (rad/d per pCi/g dry sediment assuming 180-degree exposure)

For the terrestrial receptors feeding primarily on emergent aquatic invertebrates, with little contact with the sediment itself, the ESL equation is

$$ESL = \frac{Dose\ Limit}{I_{food,i} \times BCF_{invert,j} \times TF_{blood,j} \times R_{t,j} \times DCF_{int,j}} \quad \text{Equation A-2.2-3}$$

where *Dose Limit* is 0.1 rad/d

$I_{food,i}$  is the normalized daily dietary ingestion rate for organism  $i$  (g of invertebrate-fresh wt/g of body wt/d)

$BCF_{invert,j}$  is the invertebrate bioconcentration factor for radionuclide  $j$  (pCi/g invertebrate-fresh wt per pCi/g dry sediment)

$TF_{blood,j}$  is the food to blood transfer factor for radionuclide  $j$  (unitless)

$R_{t,j}$  is the retention time for radionuclide  $j$  (d)

$DCF_{int,j}$  is the internal DCF for radionuclide  $j$  (rad/d per pCi/g fresh tissue)

The sediment radiological L-ESLs are calculated with Equation A-2.2-3 using 1 rad/d as the dose limit.

## A-2.3 Water ESLs

### A-2.3.1 Radionuclide Concentrations in Biota

For organisms immersed in water (algae, daphnid, snail, and fish), the internal concentration of any radionuclide is modeled by applying a simple BCF (Baker and Soldat 1992, 062801):

$$C_{organism,j} = C_{water,j} \times BCF_{organism,j} \quad \text{Equation A-2.3-1}$$

where  $C_{organism,j}$  is the internal concentration of radionuclide  $j$  (pCi/g of organism)

$C_{water,j}$  is the concentration of radionuclide  $j$  in water (pCi/mL)

$BCF_{organism,j}$  is the bioconcentration factor for radionuclide  $j$  in the organism (pCi/g fresh wt per pCi/mL water)

For wildlife, it is assumed the major exposure pathway to radionuclides in water is through ingestion of contaminated water. The body burden from drinking water containing radionuclides is calculated as

$$C_{organism,j} = C_{water,j} \times I_{water} \times TF_{blood,j} \times R_{t,j} \quad \text{Equation A-2.3-2}$$

where  $C_{organism,j}$  is the internal concentration of radionuclide  $j$  (pCi/g of organism)

$C_{water,j}$  is the concentration of radionuclide  $j$  in water (pCi/mL)

$I_{water}$  is the normalized daily water ingestion rate (mL of water/g of body wt /d)

$TF_{blood,j}$  is the water to blood transfer factor for radionuclide  $j$  (pCi/g fresh blood per pCi/g fresh food)

$R_{t,j}$  is the retention time of radionuclide  $j$  in the organism (d)

Values and references for the TFs and BCFs are provided in the ECORISK Database (LANL 2016, 601838, or latest version).

### A-2.3.2 DCFs

For aquatic receptors, the internal DCFs are identical to those used for terrestrial receptors. For organisms immersed in water (algae, daphnid, snail, and fish), the external dose coefficients of EPA guidance (EPA 1993, 062798) are used to estimate external dose. Coefficients used are for skin immersed in water contaminated to an infinite depth. A DCF of  $3.2 \times 10^{11}$  is used to convert the dose coefficients from Sv/s per Bq/m<sup>3</sup> to rad/d per pCi/g.

Internal dose to terrestrial receptors from water is assumed to come entirely from water ingestion. Because of the limited amount of perennial surface water at the Laboratory and the conservative model used to calculate internal dose to terrestrial receptors, external dose is assumed to be insignificant and all dose received assumed to be internal.

### A-2.3.3 Water ESL Calculations

The water ESL is defined as the water concentration (pCi/L) of a given radionuclide that yields a combined internal and external dose rate of 0.1 rad/d to a particular receptor. For aquatic receptors that spend at least part of their lives immersed in water, the ESL equation is

$$ESL = \frac{Dose\ Limit}{(BCF_{i,j} \cdot DCF_{int,j} + DCF_{ext,j})/1000} \quad \text{Equation A-2.3-3}$$

where *Dose Limit* is 0.1 rad/d

$BCF_{i,j}$  is the bioconcentration factor for organism *i* and radionuclide *j* (pCi/g fresh wt per pCi/mL water)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide *j* (rad/d per pCi/g fresh tissue wt)

$DCF_{ext,j}$  is the external dose conversion factor for radionuclide *j* (rad/d per pCi/mL water, assuming 360-degree exposure)

1000 is the number of mL per L

For the terrestrial receptors drinking contaminated water, the ESL equation is

$$ESL = \frac{Dose\ Limit}{(I_{water,i} \times TF_{blood,j} \times R_{t,j} \times DCF_{int,j})/1000} \quad \text{Equation A-2.3-4}$$

where *Dose Limit* is 0.1 rad/d

$I_{water}$  is the normalized daily water ingestion rate (mL of water/g of body wt per d)

$TF_{blood,j}$  is the water to blood transfer factor for radionuclide *j* (unitless)

$R_{t,j}$  is the retention time for radionuclide *j* (d)

$DCF_{int,j}$  is the internal dose conversion factor for radionuclide *j* (rad/d per pCi/g fresh tissue wt)

1000 is the number of mL per L

The water radiological L-ESLs are calculated with Equation A-2.3-4 using 1 rad/d as the dose limit.

### A-3.0 REFERENCES

*The following reference list includes documents cited in this appendix. Parenthetical information following each reference provides the author(s), publication date, and ERID or ESHID. This information is also included in text citations. ERIDs were assigned by the Associate Directorate for Environmental Management's (ADEM's) Records Processing Facility (IDs through 599999), and ESHIDs are assigned by the Environment, Safety, and Health Directorate (IDs 600000 and above). IDs are used to locate documents in the Laboratory's Electronic Document Management System and in the Master Reference Set. The New Mexico Environment Department Hazardous Waste Bureau and ADEM maintain copies of the Master Reference Set. The set ensures that NMED has the references to review documents. The set is updated when new references are cited in documents.*

Baker, D.A., and J.K. Soldat, June 1992. "Methods for Estimating Doses to Organisms from Radioactive Materials Released into the Aquatic Environment," PNL-8150, UC-602, Battelle, Pacific Northwest Laboratories, Richland, Washington. (Baker and Soldat 1992, 062801)

- Buchman, M.F., 2008. "NOAA Screening Quick Reference Tables," National Oceanic and Atmospheric Administration (NOAA) OR&R Report 08-1, Office of Response and Restoration Division, Seattle, Washington. (Buchman 2008, 206414)
- Crommentuijn, T., D. Sijm, J. de Bruijn, M. van den Hoop, K. van Leeuwen, and E. van de Plassche, October 2000. "Maximum Permissible and Negligible Concentrations for Metals and Metalloids in the Netherlands, Taking into Account Background Concentrations," *Journal of Environmental Management*, Vol. 60, No. 2, pp. 121–143. (Crommentuijn et al. 2000, 205264)
- Crommentuijn, T., D. Sijm, J. de Bruijn, K. van Leeuwen, and E. van de Plassche, April 2000. "Maximum Permissible and Negligible Concentrations for Some Organic Substances and Pesticides," *Journal of Environmental Management*, Vol. 58, No. 4, pp. 297–312. (Crommentuijn et al. 2000, 205265)
- Di Toro, D.M., 1985. "A Particle Interaction Model of Reversible Organic Chemical Sorption," *Chemosphere*, Vol. 14, No. 10, pp. 1503–1538. (Di Toro 1985, 062876)
- DOE (U.S. Department of Energy), July 2002. "A Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota," DOE Standard No. DOE-STD-1153-2002, U.S. Department of Energy, Washington, D.C. (DOE 2002, 085637)
- EPA (U.S. Environmental Protection Agency), 1993. "External Exposure to Radionuclides in Air, Water, and Soil," EPA-402-R-93-081, Federal Guidance Report No. 12, Office of Air and Radiation, Washington, D.C. (EPA 1993, 062798)
- EPA (U.S. Environmental Protection Agency), June 1997. "EPA Region 10 Supplemental Ecological Risk Assessment Guidance for Superfund," EPA 910-R-97-005, U.S. EPA Region 10 Office of Environmental Assessment Risk Evaluation Unit, Seattle, Washington. (EPA 1997, 215127)
- EPA (U.S. Environmental Protection Agency), 2009. "National Recommended Water Quality Criteria," available online at <http://water.epa.gov/scitech/swguidance/standards/current/index.cfm>, Office of Water, Office of Science and Technology, Washington, D.C. (EPA 2009, 109328)
- Higley, K.A., and R. Kuperman, August 1996. "Appendix C, Ecotoxicological Benchmarks for Radionuclide Contaminants at RFETS," in *Ecological Risk Assessment Methodology, Technical Memorandum for Ecological Chemicals of Concern (ECOCs) Screening Methodology*, RF/ER-96-0039, Kaiser-Hill/Rocky Flats Environmental Technology Site, Golden, Colorado. (Higley and Kuperman 1996, 062804)
- IAEA (International Atomic Energy Agency), 1992. *Effects of Ionizing Radiation on Plants and Animals at Levels Implied by Current Radiation Protection Standards*, International Atomic Energy Agency, Vienna, Austria. (IAEA 1992, 062802)
- Ingersoll, C.G., P.S. Haverland, E.L. Brunson, T.J. Canfield, F.J. Dwyer, C.E. Henke, N.E. Kemble, D.R. Mount, and R.G. Fox, 1996. "Calculation and Evaluation of Sediment Effect Concentrations for the Amphipod *Hyaella azteca* and the Midge *Chironomus riparius*," *Journal of Great Lakes Research*, Vol. 22, No. 3, pp. 602–623. (Ingersoll et al. 1996, 062873)
- LANL (Los Alamos National Laboratory), October 2016. "ECORISK Database (Release 4.0)," on CD, LA-UR-16-27520, Los Alamos National Laboratory, Los Alamos, New Mexico. (LANL 2016, 601838)

- MacDonald, D.D., C.G. Ingersoll, and T.A. Berger, July 2000. "Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems," *Archives of Environmental Contamination and Toxicology*, Vol. 39, No. 1, pp. 20–31. (MacDonald et al. 2000, 205266)
- Michelsen, T., September 2003. "Development of Freshwater Sediment Quality Values for Use in Washington State, Phase II Report: Development and Recommendation of SQVs for Freshwater Sediments in Washington State," report prepared for the Washington Department of Ecology by Avocet Consulting, Publication Number 03-09-088, Kenmore, Washington. (Michelsen 2003, 215128)
- Persaud, D., R. Jaagumagi, and A. Hayton, August 1993. "Guidelines for the Protection and Management of Aquatic Sediment Quality in Ontario," Ministry of Environment and Energy, Ontario, Canada. (Persaud et al. 1993, 205250)



## **Appendix B**

---

### *Ecological Scoping Checklist*



**B-1.0 PART A—SCOPING MEETING DOCUMENTATION**

<b>Site Identification (Include Aggregate Area)</b>	
<b>Form of Site Releases (Solid, Liquid, Vapor)</b> Describe known or suspected <u>mechanisms</u> of release (spills, dumping, material disposal, outfall, explosive testing, etc.) and describe potential <u>areas</u> of release. Reference map if appropriate.	
<b>Directly Impacted Media</b> Indicate all that apply.	Surface soil – Surface water/sediment – Subsurface – Groundwater – Other, explain –
<b>Vegetation Class Based on Geographic Information System (GIS) Vegetation Coverage</b> Indicate all that apply.	Water – Bare Ground/Unvegetated – Spruce-fir-aspen-mixed conifer – Ponderosa pine – Piñon-juniper/juniper savannah – Grassland-shrubland – Developed – Burned –
<b>Threatened and Endangered Species Habitat</b> If applicable, list threatened and endangered species known or suspected of using the site for breeding or foraging.	
<b>Neighboring/Contiguous/Upgradient Sites</b> Include a summary of chemicals of potential concern and the type of releases if impacting site. (Use this information to evaluate the need to aggregate sites for scoping and screening.)	
<b>Surface Water Erosion Potential</b> Indicate if erosion is present and type; terminal point of surface water transport; slope; and surface water run-on sources. Indicate if best management practices (BMPs) are in place or are needed.	

**B-2.0 PART B—SITE VISIT DOCUMENTATION**

Site ID	
Date of Site Visit	
Site Visit Conducted by	

**Receptor Information:**

Estimate cover	Relative vegetative cover (high, medium, low, none) = Relative wetland cover (high, medium, low, none) = Relative structures/asphalt, etc., cover (high, medium, low, none) =
Field notes on the GIS vegetation class	
Are ecological receptors present at the site? (yes/no/uncertain) Describe the general types of receptors present at the site (terrestrial and aquatic), and note the quality of habitat present at the site.	

**Contaminant Transport Information:**

Surface Water Transport Field notes on the erosion potential and BMPs, including a discussion of the terminal point of surface water transport (if applicable).	
Are there any off-site transport pathways (surface water, air, or groundwater)? (yes/no/uncertain) Provide explanation	

**Ecological Effects Information:**

<b>Physical Disturbance</b> (Provide list of major types of disturbances, including erosion and construction activities; review historical aerial photos where appropriate.)	
<b>Are there obvious ecological effects?</b> (yes/no/uncertain) Provide explanation and apparent cause (e.g., contamination, physical disturbance, other).	

**Adequacy of Site Characterization:**

<b>Do existing or proposed data provide information on the nature and extent of contamination?</b> (yes/no/uncertain) Provide explanation	
<b>Do existing or proposed data for the site address potential transport pathways of site contamination?</b> (yes/no/uncertain) Provide explanation	

**No Exposure/Transport Pathways:**

If there are no complete exposure pathways to ecological receptors on-site and no transport pathways to off-site receptors, do not complete Part C. Provide explanation/justification for proposing an ecological "No Further Action" recommendation.
---

***Additional Field Notes:***

**Provide additional field notes on the site setting and potential ecological receptors, if appropriate.**

### **B-3.0 PART C—ECOLOGICAL PATHWAYS CONCEPTUAL EXPOSURE MODEL**

Provide answers to Questions A to V to develop the Ecological Pathways Conceptual Exposure Models (use to complete figures at end of Part C).

Answer all questions with drop-down menu choices. When finished, select the entire document using control A, and press F9. This will update all the fields in the models to reflect the questions. You can also click in individual fields in the models and press F9 to update.

#### **Question A:**

Could soil contaminants reach receptors through vapors?

- Determine the volatility of the hazardous substance (volatile chemicals generally have Henry's law constant  $>1\text{E-}05$  atm-m<sup>3</sup>/mol and molecular weight  $<200$  g/mol).
- In the case of burrowing animals, the contamination would have to occur in the depth interval where burrows are present (near surface to 5 ft below ground surface).

Answer (likely/unlikely/uncertain):

Provide explanation:

#### **Question B:**

Could the soil contaminants reach receptors through fugitive dust carried in air?

- Soil contamination would have to be on the actual soil surface to become available for dust.
- In the case of dust exposures to burrowing animals, the contamination would have to occur in the depth interval where the burrows occur.

Answer (likely/unlikely/uncertain):

Provide explanation:

#### **Question C:**

Can contaminated soil be transported to aquatic communities?

If erosion is an off-site transport pathway, determine the terminal point to see if aquatic receptors could be impacted by contamination from the site.

Answer (likely/unlikely/uncertain):

Provide explanation:

**Question D:**

**Is contaminated groundwater potentially available to biological receptors through seeps or springs or shallow groundwater?**

- The potential exists for contaminants to migrate through groundwater and discharge into habitats and/or surface waters.
- Contaminants may be taken up by terrestrial and rooted aquatic plants whose roots are in contact with groundwater present within the root zone.
- Terrestrial wildlife receptors generally will not come in contact with groundwater unless it is discharged to the surface.

**Answer (likely/unlikely/uncertain):**

**Provide explanation:**

**Question E:**

**Is infiltration/percolation from contaminated subsurface material a viable transport and exposure pathway?**

- The potential for contaminants to migrate to groundwater.
- The potential for contaminants to migrate to groundwater and discharge into habitats and/or surface waters.
- Contaminants may be taken up by terrestrial and rooted aquatic plants whose roots are in contact with groundwater present within the root zone.
- Terrestrial wildlife receptors generally will not come in contact with groundwater unless it is discharged to the surface.

**Answer (likely/unlikely/uncertain):**

**Provide explanation:**

**Question F:**

**Might erosion or mass-wasting events be a potential release mechanism for contaminants from subsurface materials or perched aquifers to the surface?**

- This question is applicable only to release sites located on or near the mesa edge.
- Consider the potential erosion of surficial material and the geologic processes of canyon/mesa edges.

**Answer (likely/unlikely/uncertain):**

**Provide explanation:**



**Question G:**

Could airborne contaminants interact with receptors through the respiration of vapors?

- Contaminants must be present as volatiles in the air.
- Consider the importance of the inhalation of vapors for burrowing animals.
- Foliar uptake of vapors is typically not a significant exposure pathway.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Terrestrial Animals:

Provide explanation:

**Question H:**

Could airborne contaminants interact with plants through the deposition of particulates or with animals through the inhalation of fugitive dust?

- For this exposure pathway to be complete, contaminants must be present as particulates in the air or as dust.
- Exposure through the inhalation of fugitive dust is particularly applicable to ground-dwelling species that would be exposed to dust disturbed by their foraging or burrowing activities or by wind movement.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Terrestrial Animals:

Provide explanation:

**Question I:**

Could contaminants interact with plants through root uptake or rain splash from surficial soil?

- Contaminants in bulk soil may partition into soil solution, making them available to roots.
- Exposure of terrestrial plants to contaminants may occur through particulates deposited on leaf and stem surfaces by rain striking contaminated soil (i.e., rain splash).

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Provide explanation:

**Question J:**

Could contaminants interact with receptors through food web transport from surficial soil?

- The chemicals may bioaccumulate in animals.
- Animals may ingest contaminated food.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question K:**

Could contaminants interact with receptors through the incidental ingestion of surficial soil?

- Incidental ingestion of contaminated soil could occur while animals grub for food resident in the soil, feed on plant matter covered with contaminated soil, or groom themselves.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question L:**

Could contaminants interact with receptors through dermal contact with surficial soil?

- Exposure through dermal contact would generally be limited to organic contaminants that are lipophilic and can cross epidermal barriers.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question M:**

Could contaminants interact with plants or animals through external irradiation?

- External irradiation is most relevant for gamma-emitting radionuclides.
- Burial of contamination attenuates radiological exposure.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Terrestrial Animals:

Provide explanation:

**Question N:**

Could contaminants interact with plants through direct uptake from water and sediment or sediment rain splash?

- Contaminants may be taken up by terrestrial plants whose roots are in contact with surface waters.
- Terrestrial plants may be exposed to particulates deposited on leaf and stem surfaces by rain striking contaminated sediment (i.e., rain splash) in an area that is only periodically inundated with water.
- Contaminants in sediment may partition into soil solution, making them available to roots.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Provide explanation:

**Question O:**

Could contaminants interact with receptors through food web transport from water and sediment?

- The chemicals may bioconcentrate in food.
- Animals may ingest contaminated food.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question P:**

Could contaminants interact with receptors through the ingestion of water and suspended sediment?

- If sediment is present in an area that is only periodically inundated with water, terrestrial receptors may incidentally ingest sediment.
- Terrestrial receptors may ingest water-borne contaminants if contaminated surface waters are used as a source of drinking water.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question Q:**

Could contaminants interact with receptors through dermal contact with water and sediment?

- If sediment is present in an area that is only periodically inundated with water, terrestrial species may be dermally exposed during dry periods.
- Terrestrial organisms may be dermally exposed to water-borne contaminants as a result of wading or swimming in contaminated waters.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Animals:

Provide explanation:

**Question R:**

Could suspended or sediment-based contaminants interact with plants or animals through external irradiation?

- External irradiation is most relevant for gamma-emitting radionuclides.
- Burial of contamination attenuates radiological exposure.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Terrestrial Plants:

Terrestrial Animals:

Provide explanation:

**Question S:**

Could contaminants bioconcentrate in free-floating aquatic plants, attached aquatic plants, or emergent vegetation?

- Aquatic plants are in direct contact with water.
- Contaminants in sediment may partition into pore water, making them available to submerged roots.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Aquatic Plants/Emergent Vegetation:

Provide explanation:

**Question T:**

Could contaminants bioconcentrate in sedimentary or water-column organisms?

- Aquatic receptors may actively or incidentally ingest sediment while foraging.
- Aquatic receptors may be directly exposed to contaminated sediment or may be exposed to contaminants through osmotic exchange, respiration, or ventilation of sediment pore waters.
- Aquatic receptors may be exposed through osmotic exchange, respiration, or ventilation of surface waters.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Aquatic Animals:

Provide explanation:

**Question U:**

Could contaminants bioaccumulate in sedimentary or water-column organisms?

- Lipophilic organic contaminants and some metals may concentrate in an organism's tissues.
- Ingestion of contaminated food may result in bioaccumulation through the food web.

Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):

Aquatic Animals:

Provide explanation:

**Question V:**

**Could contaminants interact with aquatic plants or animals through external irradiation?**

- External irradiation is most relevant for gamma-emitting radionuclides.
- The water column acts to absorb radiation; therefore, external irradiation is typically more important for sediment-dwelling organisms.

**Provide quantification of exposure pathway (0=no pathway, 1=unlikely pathway, 2=minor pathway, 3=major pathway):**


**Aquatic Plants:**

**Aquatic Animals:**

**Provide explanation:**


# **Ecological Scoping Checklist** **Terrestrial Receptors** **Ecological Pathways Conceptual Exposure Model**

**NOTE:**  
 Letters in circles refer  
 to questions on the  
 scoping checklist.



# **Ecological Scoping Checklist** **Aquatic Receptors** **Ecological Pathways Conceptual Exposure Model**

**NOTE:**  
 Letters in circles refer  
 to questions on the  
 scoping checklist.





## **SIGNATURES AND CERTIFICATION**

**Checklist completed by:**

**Name (printed):** \_\_\_\_\_

**Name (signature):** \_\_\_\_\_

**Organization:** \_\_\_\_\_

**Date completed:** \_\_\_\_\_

**Checklist reviewed by:**

**Name (printed):** \_\_\_\_\_

**Name (signature):** \_\_\_\_\_

**Organization:** \_\_\_\_\_

**Date reviewed:** \_\_\_\_\_





### Legend

- |   |                                      |
|---|--------------------------------------|
|  | Cerro Grande extent                  |
|  | LANL boundary                        |
|  | Paved road                           |
|  | 20 ft Contour                        |
|  | 100 ft Contour                       |
| Description   |                                      |
|  | Aspen-Riparian-Wetland               |
|  | Cerro Grande Fire high burn severity |
|  | Grass species                        |
|  | Mixed conifer-Spruce-Fir             |
|  | Open Water                           |
|  | Pinon-Juniper                        |
|  | Ponderosa Pine                       |
|  | Shrub species                        |
|  | Urban-Sparse-Bare rock               |

## DATA SOURCES

Title; Owner; ID; Intended Scale; Publication Date

1991 Hypsography (100ft); Los Alamos National Laboratory, Remediation Services Project;  
NA; Unknown; 1991.

1991 Hypsography (20ft); Los Alamos National Laboratory, Remediation Services Project; NA; Unknown; 1991.

Boundary of Department of Energy Property In and Around the Los Alamos National Laboratory;  
Site and Project Planning Group; NA; Unknown; February 1, 2003.

Cerro Grande Extent, Los Alamos National Laboratory, Earth and Environmental Sciences  
GISLab; NA; Unknown; April 5, 2002.

Roads, surfaced; Los Alamos County; ER2004-0071; Unknown; 2004; Provided by the Los Alamos National Laboratory Remediation Services Project.

Vegetation Taxonomy; Los Alamos National Laboratory, Environmental Stewardship Ecology Group; NA; Unknown; June 19, 2003.

LA-UR-15-27577

Cartography by: Doug Walther  
Date: December 21, 2004  
GISLab M#: 201262  
Request #: 13572, 13580, 1369

State Plane Coordinate System, New Mexico Central Zone  
1983 North American Datum. Units Feet.

Grid displays New Mexico State Plane coordinates in feet.


1:24,000

DISCLAIMER OF LIABILITY:

Neither the United States Government nor the University of California nor any of their agents or employees makes any warranty, express or implied, including the warranties of merchantability and fitness for a particular purpose, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information disclosed herein.

Work performed on behalf of the RRES Remediation Services Project, Los Alamos  
Los Alamos, NM 87545

--



Los Alamos  
NATIONAL LABORATORY