LA-UR-12-23076

Approved for public release; distribution is unlimited.

Title: Update on the Electron Source Model

Author(s): Cowee, Misa
Winske, Dan

Intended for: DTRA Review

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Update on the Electron Source Model

M. M. Cowee (ISR-1) and D. Winske (XCP-6)
Los Alamos National Laboratory

DTRA Review
July 19, 2012

Abstract

- We summarize work done in FY12 on the Los Alamos Electron Source Model (ESM), which predicts the distribution of beta-decay electrons after a high altitude nuclear explosion (HANE) as a function of L, energy, and pitch angle. In the last year we have compared model results with data taken after the Russian 2 HANE test of 1962 and presented results at the HEART conference. We discuss our future plans to continue comparison with HANE data and to develop the code to allow a more complex set of initial conditions.

Outline

1. Review of the Electron Source Model (ESM) calculations
2. Summary of ESM runs in FY12
3. Plans for future development of code
4. Plans for future comparisons with HANE test data

1. Description of the ESM: Basic calculation

The basic calculation is as follows:

1. The location and yield of the burst and the debris ion thermal velocity (or mass) determine the spatial extent of the debris ions (fission fragments) immediately after the burst (i.e. the burst region size).
 - Debris ion thermal velocity is 1500 km/s (default)
 - Total debris ion mass is determined from thermal velocity (1500 km/s default), yield and kinetic fraction of the yield (typically 0.25 default)
 - Burst region size is determined by the minimum of the equal mass radius or the magnetic confinement radius
 - Debris ions are uniformly spread throughout the burst region (default)
2. The debris ions immediately begin to β -decay and produce MeV electrons (β^-) on the same field line as the debris ions.
 - β^- are produced isotropically in local pitch angle
3. For times less than the burst expansion time, τ_B , all the debris ions are assumed to be in the burst region and β -decay there; τ_B is calculated as the radius of the burst region divided by the thermal velocity of the debris ions

1. Description of the ESM: Basic calculation (cont.)

3. (continued) At times greater than τ_B , some of the debris ions continue to remain in the burst region but most of them stream out of the burst region along field lines.
4. Of these streaming debris ions, a few will mirror and be trapped but most are directly lost to the atmosphere.
 - We can set the fraction of debris ions lost to atmosphere, the fraction that remains in burst region, and the fraction that mirrors
 - Debris ions stream down field lines

1. Description of the ESM: Basic calculation (cont.)

5. The start/stop times for the debris ion β^- -decay are calculated from their streaming velocity and the length of the field line. The number of β^- is then calculated using the well-known expression for the β^- -decay rate from radioactive fission fragments.
6. Assuming the top of the atmosphere (default 80 km) is a hard cutoff for loss of electrons, we can calculate the loss cone angle and determine the number of β^- that will be trapped and their equatorial pitch-angles.
7. With the flux tube volume and the number of trapped β^- , we calculate the flux of trapped β^- at early times. To calculate a later-time (i.e. 'longitudinally averaged') flux, we use the volume of the L shell over all longitudes.
 - We can also approximate loss due to drift past the South Atlantic Anomaly by imposing a higher altitude hard cutoff for the top of the atmosphere
8. Assuming the trapped β^- have the expected fission spectrum (Carter, et al. 1959), we have determined the flux of trapped β^- , $f(L, E, \alpha)$ which is input to DREAM (Dynamic Radiation Environment Assimilation Model).

2. Summary of ESM runs in FY12

- To validate ESM against HANE data, we used the Russian 2 burst because it is considered the best observed of the HANE tests
 - + Used data from the Artificial Radiation Belt Database (ARBDB) for comparison
 - + Showed that artificially distributing the debris ions in L (e.g. presumption of the “jetting” mechanism, could yield fluxes in general agreement with data)
 - Comparison with measurements in the different energy ranges suggested that the measurements in the lowest energy bin included a contribution from ambient electrons that were heated by the burst
 - + This work was presented at HEART 2012 and paper submitted to conf proceedings
- Then, we looked into doing similar comparison for Starfish data, but the lack of a background measurement for Starfish introduces too much uncertainty ...
 - + Unless there is a basic physics understanding of “jetting”, comparing model results to Starfish is not effectively in constraining the model
 - Plots showing background-subtracted Starfish fluxes are created using NASA’s AE-8 electron flux maps which effectively represent average conditions. Real conditions at a given time can vary significantly from these averages, so any structure in these background-subtracted Starfish fluxes could be a product of this.

3. Plans for future development of the code

ESM improvements:

- + Realistic magnetic field model
- + Realistic atmosphere density model
- + Variable initial debris ion velocity and spatial distribution
- + Realistic calculations for debris ion streaming and mirroring
- + Time and energy-dependent β -decay rates
- + Users select which models they would like to use for the environment (magnetic field, atmosphere density) and the calculations (debris ion loading in velocity and space in burst region)
- + The LANLGeomag suite has the magnetic field models built in and routines to do relevant calculations (e.g. field line tracing, McIlwain L)

UNCLASSIFIED

3. Plans for future ESM and HANE test data comparison

- We would like to continue comparison of ESM with HANE data:
 - + ESM is a very flexible code and we can easily carry out sensitivity studies for the input parameters, and identify particular features of the resulting distributions which agree with HANE test observations
 - + In particular, we could look at how the initial distribution of debris ions along the field lines affects the structure of the fluxes in L and B/B_{eq}
 - We do not have an accepted physical mechanism which can distribute debris ions over such a large region of space. If we can better constrain the characteristics of the initial debris ion distribution, this may suggest a mechanism.
 - + Additionally, we can include the effects of pitch-angle scattering by ambient, natural-environment plasma waves on the ESM fluxes using the DREAM diffusion code
 - + These results could also be used to aid the re-interpretation of the HANE test data
 - They can indicate possible pitch-angle distributions for the HANE electrons for mapping the observed fluxes along the magnetic field lines, and calculating the trapped beta inventory
- However, work is currently being done to improve the data (e.g. Roth, McDaniels?) so it may be better to wait until we these efforts are done
- Additionally, the UCLA laser experiments and LLNL 3D simulations may yield other physics which should be included in the ESM

Extra slides follow

UNCLASSIFIED

Description of the ESM: Magnetic field model

- We use a dipole magnetic field, where quantities can be determined analytically
 - + λ is north latitude
 - + ϕ is east longitude
 - + r is radial distance from the center of the earth

- In this geometry, magnetic field lines are defined by

$$r = LR_E \cos^2 \lambda$$

- + where L is the McIlwain parameter that describes the distance in R_E where a field line crosses the magnetic equator

- Figure shows

- + magnetic field lines at various L (solid lines)
- + lines of latitude (dashed lines)
- + Earth surface at $r = R_E$ (solid line)
- + top of the atmosphere at $r = R_a$ (dotted line)

UNCLASSIFIED

2. Description of the ESM: Ionosphere profile

- We consider a three-region ionospheric model:

- + low altitude region
 - below $h_c = 180$ km
 - scale height $h_1 = 10$ km
- + high altitude region
 - above $h_d = 320$ km
 - scale height $h_3 = 100$ km
- + intermediate altitude region
 - between h_c and h_d
 - scale height $h_2 = 20$ km

- At 400 km altitude, we assume that the plasma consists of O^+ with number density $3 \times 10^5 \text{ cm}^{-3}$ or mass density of $8 \times 10^{-18} \text{ g/cm}^3$ (Brecht et al., 1995; Dyal, 2006)
- We also include a simple latitude dependence, taking the ionosphere mass density to vary as $\cos^{-2} \lambda$ for $0^\circ \leq \lambda \leq 85^\circ$ (e.g., Denton et al., 2004).

2. Description of the ESM: Burst region size

- Based on hybrid simulation results from Winske and Gary (2007), the size of the burst region is determined by the minimum of the so-called “equal mass radius” or “magnetic confinement radius”:
 - + Equal mass radius, R_m , corresponds to a volume of background ionospheric plasma (‘a’) equal to the total mass of the debris ions (‘d’):

$$M_d = \frac{4\pi}{3} n_a m_a R_m^3$$

- + Magnetic confinement radius, R_b , corresponds to a volume of magnetic field energy equal (‘B’) to the kinetic energy of the debris ions (‘d’):

$$\frac{1}{2} M_d U_d^2 = \frac{4\pi}{3} R_b^3 \frac{B_B^2}{8\pi}$$

- + These quantities are calculated above and below the burst point by integrating the atmosphere density or magnetic field energy along a line from the center of the earth through the burst point

2. Description of the ESM: Burst region size (cont.)

- For burst locations below ~800 km, the equal mass radius is smaller than the magnetic confinement radius above the burst point

- Figure shows burst regions for varying burst altitude. The “cube-shaped” burst region in red is approximated in the model as the volume shown in blue defined by:

- + min/max L of burst region
- + min/max λ of burst region
- + longitudinal width of burst region set equal to latitudinal width
- + Burst defined by burst yield (y_B), height (h_B), and north latitude (λ_B)

UNCLASSIFIED

2. Description of the ESM: L, λ calculation grid

- The regions within the burst flux tube (defined by $L_{\min}$ to $L_{\max}$) at λ north and south of the burst region are called the conjugate regions
 - + North ('N') and South ('S') conjugate regions extend from the edges of the Burst ('B') region ($\lambda_{\min}$ or $\lambda_{\max}$) to the top of the atmosphere (80 km altitude)
- Calculation grid:
 - + the range $L_{\min}$ to $L_{\max}$ is divided into L subregions
 - + the N and S conjugate regions are further divided into λ subregions with maximum width of 0.5° .

2. Description of the ESM: Debris ion distribution

- At $t < \tau_B$, the debris ions (shaded region) are all present in the burst region volume, uniformly distributed
 - + Expansion velocity, τ_B , is equal to the burst region radius divided by the debris ion thermal velocity
- At $t > \tau_B$, debris ions stream out of burst region along the field lines and are lost to the atmosphere
 - + Some debris ions can remain in the burst region and do not stream away
 - + Some streaming debris ions are trapped rather than lost to atmosphere
- The times when debris ions stream through the λ subregions is used for the start/stop times in the β -decay equation
 - + For simplicity, half of the debris ions that do not remain in the burst region are distributed to each of the N and S conjugate regions and are assumed to be uniformly distributed throughout those volumes