LA-UR-13-22558

Approved for public release; distribution is unlimited.

Title: CMIME Update on Metal-Metal Oxide Atomistic Models

Author(s): Valone, Steven M.
Baskes, Michael I.
Gibson, Joshua
Allen, Jonathan R.
Dunlap, David H.
Atlas, Susan R.

Intended for: Project update; want to send vugraphs to colleague at UNM

Issued: 2013-04-10

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

CMIME Update on Metal-Metal Oxide Atomistic Models

Steve Valone

LANL

Mike Baskes, UCSD & LANL
Josh Gibson, UNT (aka Srini connection)
Jon Allen, Prof. Dave Dunlap, Prof. Susan Atlas
UNM Physics

Multiple Materials, Multiple Models

Nano-scale oxides
strengthen steels

HRTEM of Fe-
oxide
interface in
12CrODS

*Oka et al. Met
Trans (2007)*

Osman & Jeff

Metals

- Metallic bonding
- No charge transfer

EAM, MEAM, F-S,
COMB3, ...

Covalent

- Angular bonding
- No charge transfer

Tersoff, ReaxFF,
COMB, BETM, ...

Ceramics

- Ionic bonding
- Charge transfer controlled
heuristically, no physical basis

ReaxFF, COMB, QTPIE,
QEq, BETM, ...

Metals: Embedded Atom Models

$$E = \sum_A \left(F_{EAM}(\bar{\rho}_A) + \frac{1}{2} \sum_{B \neq A} \phi_{AB}(R_{AB}) \right)$$

- EAM energy

$$F_{MEAM}(\bar{\rho}) = A \frac{\bar{\rho}}{\rho_0} \ln \frac{\bar{\rho}}{\rho_0}$$

$$F_{F-S}(\bar{\rho}) = A \sqrt{\frac{\bar{\rho}}{\rho_0}}$$

- Embedding Function

$$\phi(R) = \frac{2}{Z} \{ E^u(R) - F(\bar{\rho}^u(R)) \}$$

- Pair Potential

Metal Oxides

**Dislocations loops in ceria
can be cation vacancies**

**HRTEM of stacking fault in
ceria created by e^- irradiation**

Yasunaga et al. Nucl Instr Meth B (2008)

**Radiation damage produces
high-energy defects**

Oxy
Hf
Mg

Valone et al. 2009

Ionic Materials: Buckingham, ReaxFF, COMBS ...

$$\phi_{AB}^{ionic} = A^{Buck} \exp\left(-\frac{R_{AB}}{\rho_{AB}^{Buck}}\right) - \frac{C_6}{R_{AB}^6} + \frac{q_A q_B}{\epsilon_0 R_{AB}}$$

$$F_{IM}(q) = E_A^0 + \chi_A q + 1/2 \eta_A q^2$$

$$E = \sum_A \left(F_{IM}(q_A) + \frac{1}{2} \sum_{B \neq A} \phi_{AB}(R_{AB}) \right)$$

- **Buckingham:**
Coulombic interactions between atoms
Dispersion interactions
 - Constant charges
 - No bond-breaking
- **Qeq, ...: Variable charges**
IM – Iczkowski-Margrave
 χ : electronegativity $\sim$ Fermi level
 η : hardness $\sim$ charge transfer gap
- **Same general form as EAM**

Charge Flow at Interfaces: Charge-Dependent Models

Phillpot et al.: Si/SiO₂
(PRB, 2007)

Charge-flow Regulation:

Development of space charge at hetero-interface

Need to attenuate flow: COMBS by higher-order charge-dependencies

Nano-scale oxides
strengthen steels

HRTEM of Fe-oxide
interface in 12CrODS

Oka et al. Met Trans (2007)

Issues :

Physical basis for charge flow regulation?

What is a metal, insulator atomistically?

General model reduces to metals and ceramics?

FH Density Matrix for Diatomic: Chemical Potential and Charge Equilibration

Chemical Potential (dE/dq): Varies greatly among models. PPLB is correct limit for dissociated (gas-phase) atoms. IM is linear.

Attenuate charge flows: Crossing of black line with colored curves

Model Hamiltonian Approaches

Drautz, 2007

Fragment model H

↑ cp, gaps, hopping
 $\hat{H}_{el}$ & $\bar{E}$ – fragment

↑
 $\hat{H}_{el}$

Fragment model potl

smv et al.

$$E = \sum_A \left(F + \frac{1}{2} \sum_{B \neq A} \phi_{AB} \right)$$

TB vs FH

What is the energy of the **next electron** added to the system?

What is the energy of the **atoms** if another electron added to the system?

Many e⁻ Hamiltonians in Fragments

$$\hat{H}_{\text{el}} \psi_i = \sum_A \left(\hat{H}_A(\mathbf{z}_A) + \frac{1}{2} \sum_{B \neq A} \hat{V}_{AB}(\mathbf{z}_A, \mathbf{z}_B) \right) \psi_i$$

- Objects defined by Hamiltonians
- “Atom-in-Molecule” Hamiltonian well-known
 - Moffitt (1951)
 - Always assigned number of electrons N_A equal to nuclear charge Z_A
 - Makes sense physically, but “ $N_A = Z_A$ ” assumption **problematic**

$$|AB\rangle = C_0 |A^0 B^0\rangle + C_+ |A^+ B^-\rangle + C_- |A^- B^+\rangle$$

Pauling, 1929, Slater 1931, Mulliken, 1934

Fragment Hamiltonian (FH) Model:

Fragment E + Fragment-Fragment

Fragment:

$$E_A = \text{tr } H_A \Gamma_A(C_+, C_0, C_-)$$

H matrix

$$H_A = H_{00}^A + \left(\text{Diagram} \right)_{A_i}$$

States of atoms paramount

Density Matrix

$$\Gamma_A(C_+, C_0, C_-) = |A\rangle\langle A| = \begin{pmatrix} C_+C_+ & C_0C_+ & C_+C_- \\ \dagger & C_0C_0 & C_0C_- \\ \dagger & \dagger & C_-C_- \end{pmatrix}_A$$

Fragment – Fragment:

$$V_{AB} = V_{00}^{AB} + \begin{pmatrix} V_{++} - V_{00} & V_{0+} & V_{+-} \\ \dagger & 0 & V_{0-} \\ \dagger & \dagger & V_{--} - V_{00} \end{pmatrix}_{AB}$$

Recovers energy scales:
Ionization, Attachment,
Gaps, Hopping

smv, JCTC, JPCL 2011

$$\mu_A = -(\mathcal{I}_A + \mathcal{E}_A)/2$$

$$U_A = (\mathcal{I}_A - \mathcal{E}_A)/2$$

Fragment
CP
Gaps

$$\Gamma_{AB}(C_+, C_0, C_-) = (|B\rangle\langle A| + |A\rangle\langle B|)/2$$

STEM-EELS: Fe-TiO₂

Effect of the presence of iron on the titanium valence, points outside this region are noisy and difficult to quantify

Model H, Spectral Density & Density of States

$$\bar{E} \approx \sum_A \sum_{\zeta} \left(\varepsilon_{A\zeta A\zeta} C_{A\zeta}^\dagger C_{A\zeta} + \frac{1}{2} \sum_{B \neq K} \sum_{\zeta, \zeta'} V_{A\zeta B\zeta'} C_{A\zeta}^\dagger C_{B\zeta'} \right)$$

$$EC_{A\zeta} = \varepsilon_{A\zeta A\zeta} C_{A\zeta} + \frac{1}{2} \sum_{B \neq K} \sum_{\zeta, \zeta'} V_{A\zeta B\zeta'} C_{B\zeta'}$$

• Structure of tight-binding H, but FH means something different

$$\Gamma(z) = -\frac{1}{\pi} \text{Im} G(z^+)$$

$$\rightarrow \text{tr} \delta(zS - H_\Gamma - H_* - V_{\Gamma*}) \Gamma$$

- Related to vast knowledge, data of electron spectroscopy, electronic structure calculations
- Not used in typical atomic-scale models

- Green fcn introduces contributions from the spectrum of H
- Density of states (energy) $\Gamma(z)$ observable and computable
“Spectral Density Matrix”

Electronic Spectra and Densities of States

XPS, BIS and XAS of UO_2
Tobin & Yu, PRL (2011)

See also Zaanen, Sawatzky & Allen, 1985

DOS changes with vol, new information

Matrix of Charge-State Variables

Link to traditional atomistic models

$$\Gamma_A(q, \tau) = \begin{pmatrix} 1 - \tau & \sqrt{(1 - \tau)(\tau + q)/2} & \sqrt{(1 - \tau)(\tau - q)/2} \\ \dagger & \dfrac{\tau + q}{2} & \dfrac{\sqrt{\tau^2 - q^2}}{2} \\ \dagger & \dagger & \dfrac{\tau - q}{2} \end{pmatrix}_A$$

$$q_A \equiv C_{A^+}^2 - C_{A^-}^2$$

- **Charge:** Difference between occupation numbers
– Not unique: choose definition

$$\tau_A \equiv C_{A^+}^2 + C_{A^-}^2$$

- **Ionicity** related to metallic character
- Most empirical potentials only use q or τ (ie $\bar{\rho}$)

Unified Embedding Energy

$$F_{\text{FH}}(q, \tau) = F_{\text{FH}}^{(\text{gap})}(q, \tau) - F_{\text{FH}}^{(\text{hop})}(q, \tau)$$

Gap + 2e⁻ 1e⁻

$$F_{\text{FH}}^{(\text{gap})}(q, \tau) = E_0 - \mu q + U\tau + W_{2e} \sqrt{\tau^2 - q^2}$$

$$F_{\text{FH}}^{(\text{hop})}(q, \tau) = W_{1e}^+ \sqrt{(1 - \tau)(\tau + q)} + W_{1e}^- \sqrt{(1 - \tau)(\tau - q)}$$

smv, JCTC, JPCL 2011

- Arrange into two embedding energies, two variables:

1. Total charge fluctuations (τ)
2. Imbalance in charge fluctuations (q)

- Mulliken Electronegativity:

$$\mu^* = -(I^* + E^*)/2$$

- Fragment Hardness, Local “Hubbard U”:

$$U^* = (I^* - E^*)/2$$

* means environmental influence

$$F_{\text{F-S}}(\tau) \sim (W_{1e}^+ + W_{1e}^-) \sqrt{\rho}$$

$$E(q) = E_0 + \chi q + \eta q^2$$

Hardness/gap 2nd order in q

Embedding for Elements: Ni

$$F_{\text{FH}}(0, \tau) = \underbrace{F_{\text{FH}}^{(\text{gap})}(0, \tau)}_{\text{Gap} + 2e^-} - \underbrace{F_{\text{FH}}^{(\text{hop})}(0, \tau)}_{1e^-}$$

$$F_{\text{FH}}^{(\text{gap})}(0, \tau) = E_0 + (U + W_{2e})\tau$$

$$F_{\text{FH}}^{(\text{hop})}(0, \tau) = (W_{1e}^+ + W_{1e}^-) \sqrt{(1 - \tau)\tau}$$

• $q = 0$

1. Terms simplify

2. Other models are special cases

• Database from E-vol curves of Ni

Gaps and Gap Closure in Ni

- Key quantity for gap closure combines gap U and widths W^{1e-} and W^{1e+}

$$F_{\text{FH}}(\bar{\rho}) = U F_{\text{FH}}^{(\text{gap})}(\bar{\rho}) - 2(W^{1e-} + W^{1e+}) F_{\text{FH}}^{(\text{hop})}(\bar{\rho})$$

Surfaces and Stacking Faults in Ni: Truth in Advertising

- Expect things to be off a bit
- Stacking fault OK – it's 3d
- Surfaces not so great – 2d
- Need the other gadgetry:
2nd, 3rd neighbors; Angular dependencies

Lattice Distortions, Phase Changes, Defects

B1

B2

B3

- FH model sensitive to different crystal structures (neighbors, deformations), similar to MEAM, NMEX (Taylor), A-EAM, COMBS

$$W_A^{1e+} \sim \frac{L_{A^+}}{\sqrt{L_{A^+}}} w_A^{1e} = \sqrt{L_{A^+}} w_A^{1e}$$

Need dimensional dependence, shape changes

$$U_A^* (h_{BAB'}^+, h_{BAB'}^-, \tau_B, \tau_{B'})$$

Oxides: Change model of charge
Go back toward variables for model H

Combine fits from elements

Three-body

shape-dependent

$$\tau = \frac{\bar{\rho}_+^2 + \bar{\rho}_-^2}{1 + \bar{\rho}_+^2 + \bar{\rho}_-^2}$$

$$q = \frac{\bar{\rho}_+^2 - \bar{\rho}_-^2}{1 + \bar{\rho}_+^2 + \bar{\rho}_-^2}$$

Unified Model of Bonding in FH View

Metals & Ceramics

**Ionicity
orthogonal to
net charge**

FH Mapping to TB Model: 1D, ∞ Chain

TB

1-band: $E(k) = E_0 + 2V \cos ka$

FH analog

3-bands: 3 states per site

$$0 = \left| E(k) - \left(\tilde{E}_0 + 2\tilde{V} \cos ka \right) \right|$$

- Periodic potential

$$V(x) = \sum_m v(x - ma)$$

- Individual electrons move in potential
- Fill e^- levels successively (fermionic)

- Multiple bands:
Integer charges replace principle quantum number as band index

- States of **fragments**

Gap Closure in FH Model: 1D, Periodic, ∞ Chain

For 1D chain

- Problem suggested by Adrian
- Multiband TB: metal or insulator depending on filling
- Hubbard model: No gap closure unless $U \rightarrow 0$
cf Lieb & Wu, 1968
- **Atomistic FH model:** Gaps open, close with “local U ” and hopping

$$I = 2 \quad V_{00} = -0.5$$

$$\mathcal{E} = 1 \quad V_{+-} = -0.0$$

FH DOS

Summary: FH Model Potentials

- **Model Hamiltonian for atomic scale**
 - States of ATOMs, rather than states of e^- s
 - Tight-binding-like in character
 - Unify models for oxides and metals
- **Introduces local Hubbard energy or chemical hardness**
 - Some connection to 3-band tight-binding model
 - Gaps and gap closure in an atomistic model
- **Ni model developed, applied to surfaces, stacking fault E, paper about to be submitted**
- **Apply FH model to NiO based on Ni, O**
- **Over winter holiday discovered that we missed a term**

References

- [1] RP Iczkowski and JL Margrave, J Am Chem Soc **83**, 3547 (1961).
- [2] HO Pritchard and FH Sumner, Proc Roy Soc, London A **235**, 136 (1956).
- [3] JP Perdew, RG Parr, M Levy, JL Balduz, Jr, Phys Rev Lett **49**, 1691 (1982).
- [4] SM Valone and SR Atlas, J Chem Phys **120**, 7262 (2004).
- [5] SM Valone, J Li, and S Jindal, IJQC **108**, 1452 (2008).
- [6] RP Feynman, Statistical Mechanics: A Set of Lectures, (Addison-Wesley, Reading MA, 1998), Ch. 2
- [7] SM Valone, J Chem Theory Comput **7**, 2253 (2011)
- [8] SM Valone, J Phys Chem Lett **2**, 2618 (2011)

Reduction to Fragment Variables

$$|AB\rangle = C_0|A^0B^0\rangle + C_+|A^+B^-\rangle + C_-|A^-B^+\rangle$$

$$|\psi\rangle \equiv \sum_i c_i |i\rangle$$

$$|A\rangle \equiv \sum_{\zeta} c_{A^{\zeta}} |A^{\zeta}\rangle$$

$$|A^{\zeta}\rangle \equiv \sum_i \delta_{\zeta, \zeta_i^A} c_i |i\rangle / c_{A^{\zeta}}$$

$$c_{A^{\zeta}}^2 \equiv n_{A^{\zeta}} = \sum_i \delta_{\zeta, \zeta_i^A} c_i^2$$

- Generalize paradigm for diatomic
- Collect VB states (N-e⁻) and expansion coefficients according to fragment properties (integer charges)
- Can only define occupation numbers, not coefficients

$$|\psi\rangle = |A\rangle = C_0 \underset{\hat{H}_{A^0}}{|A^0\rangle} + C_+ \underset{\hat{H}_{A^+}}{|A^+\rangle} + C_- \underset{\hat{H}_{A^-}}{|A^-\rangle}$$

Model Hamiltonian Approach

Full, Many-electron $\hat{H}$

$$\hat{H}_{\text{el}} = \sum_{\mathbf{e}^- \mathbf{s}} \left(T_i + \sum_{\text{atoms } A} V_{iA} + \frac{1}{2} \sum_{j \neq i} V_{ij} \right) + \hat{V}_{\text{NN}}$$

$$\begin{aligned} \hat{H}_{\text{el}} &\rightarrow \hat{h}_{\text{TB}} + U \hat{D}_H \\ &= - \sum_{R_A \neq R_B, \sigma} t(R_A - R_B) \hat{c}_{R_A, \sigma}^\dagger \hat{c}_{R_B, \sigma} \end{aligned}$$

$$+ U \sum_A \hat{n}_{A\downarrow} \hat{n}_{A\uparrow}$$

Gebhard

Tight-bonding, Hubbard, DFT H's
One-electron granularity

Fragment $\hat{H}$

Atomic granularity

$$\hat{H}_{\text{el}} \equiv \sum_A \left(\hat{H}_A + \frac{1}{2} \sum_{B \neq A} \hat{V}_{AB} \right)$$

$$E = \sum_A \left(F + \frac{1}{2} \sum_{B \neq A} \phi_{AB} \right)$$

High Resolution STEM Images: Annular Dark and Bright Field at the Interface

Jeff & Osman

Dark Field, High Z Contrast

Bright Field, Low Z Contrast

Missing Element: Charge Fluctuations

- Distribute electrons among fragments: Charge fluctuations
- Vary integer number of electrons assigned to fragments
→ Open system
- Superpose states with different numbers of e^- on sites