LA-UR-18-21136

Approved for public release; distribution is unlimited.

Title: Quantum Effects on Cosmology: Probing Physics Beyond the Standard Model with Big Bang

Author(s): Paris, Mark W.

Intended for: Report

Issued: 2018-02-13

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Quantum Effects on Cosmology: Probing Physics Beyond the Standard Model with Big Bang Nucleosynthesis

M. Paris (T-2, LANL), E. Grohs (T-2)

V. Cirigliano (T-2), S. Shalgar (T-2), L. Johns
(UCSD) & G. Fuller (UCSD)

Overview & Motivation

Develop new computational/code capability for *precision* **BBN***

- Cosmological observations of light primordial abundances are getting **very good**: $\sim 1\%$ level
- Improved modeling for applications (**security**, **energy**, **safety**, **basic res.**) requires *precision* ($< \text{few } \%$) nuclear cross sections
- **Objective**: **Verification & Validation** of **light nuclear reaction cross sections independent of other (e.g. fission) cross sections**
- The **Early Universe** becoming a highly constrained environment
 - ▣ Big Glass (Extremely Large Telescopes: E-ELT, GMT, TMT): $\sim 10\times$ light
 - ▣ Next generation **Cosmic Microwave Background radiation (CMB-S4)**
 - ▣ Observables: **CMB** anisotropies/polarizations; **D**, $^3\text{He}/^4\text{He}$, ^4He , ^7Li

***BBN = 'big bang' or 'primordial nucleosynthesis'**

Collaboration

-LDRD support: Exploratory Research (20170430ER) & Center for Space & Earth Sciences (CSES)

□ University

▣ George Fuller (UCSD), Luke Johns (UCSD)

□ LANL

▣ Vincenzo Cirigliano, E. Grohs, S. Shalgar, M. Paris

Publications:

- [1] Vincenzo Cirigliano, Mark W. Paris, and Shashank Shalgar. Effect of collisions on neutrino flavor inhomogeneity in a dense neutrino gas. *Physics Letters B*, 774:258 – 267, 2017.
- [2] Shashank Shalgar. Multi-angle calculation of the matter-neutrino resonance near an accretion disk. *Journal of Cosmology and Astroparticle Physics*, 2018(02):010, 2018.
- [3] E. Grohs, George M. Fuller, C. T. Kishimoto, and Mark W. Paris. Lepton asymmetry, neutrino spectral distortions, and big bang nucleosynthesis. *Phys. Rev.*, D95(6):063503, 2017.
- [4] E. Grohs and George M. Fuller. Insights into neutrino decoupling gleaned from considerations of the role of electron mass. *Nuclear Physics B*, 923:222 – 244, 2017.

The BBN Epoch

$$T_{\text{initial}} \sim 10 \text{ MeV} \Rightarrow T_{\text{final}} \sim 1 \text{ keV}$$

$$\text{Large entropy: } \eta = \frac{n_b}{n_\gamma} \sim 10^{-10}$$

BBN briefly

Precision effects
viz. BURST

□ Early FLRW universe

- Homogeneous & isotropic (??)
- Hubble expansion drives out-of-equilibrium dynamics
- Cooling thermonuclear fusion reactor
 - nuclei bathed in neutrinos/photons
 - produces ^4He (Y_p), D, ^3He & ^7Li
 - NB: out-of-equilibrium

□ Status of observation

- [now] $\mapsto$ [with extremely large telescopes]
- $Y_p \sim 5\% \mapsto \sim 1\%$
- $\text{D}/\text{H} \sim 2\% \mapsto \lesssim \text{O}(1\%)$
- $^7\text{Li} \sim \text{O}(50\%) \mapsto \text{O}(\text{few } \%)$

7Li problem

BBN briefly

Precision effects
viz. BURST

- Early **FLRW** universe
 - ▣ Homogeneous & isotropic
 - ▣ Hubble expansion drives out-of-equilibrium dynamics
 - ▣ Cooling thermonuclear fusion reactor
 - nuclei bathed in neutrinos/photons
 - produces ${}^4\text{He}$ (Y_p), D & ${}^7\text{Li}$
 - NB: out-of-equilibrium
- Status of observation
 - ▣ [now] $\mapsto$ [with extremely large telescopes]
 - ▣ $Y_p \sim 5\% \mapsto \sim 1\%$
 - ▣ $\text{D}/\text{H} \sim 2\% \mapsto \lesssim \text{O}(1\%)$
 - ▣ ${}^7\text{Li} \sim \text{O}(50\%) \mapsto \text{O}(\text{few } \%)$

Self-consistent neutrino transport

□ “Self-consistent”

- Previous approaches evolve neutrinos and ‘post-process’ BBN
 - $\Delta Y_p \sim 0.05\%$: very small; probably unmeasurable
- Current approach solves neutrino energy transport and BBN *concurrently*
 - $\Delta Y_p \sim 1\%$: possibly observable with ELT's

□ Computational approach

- Solve the neutrino quantum kinetic equations
 - Allow lepton number asymmetry
- Describe e^+/e^- /photon/baryon plasma in terms of equilibrium distributions for all times/temperatures

Neutrino quantum kinetics (flavor dynamics)

$$\left[\partial_t - pH[\rho; t] \partial_p \right] \rho(p, t) - i \left[\left(H_V + H_{\nu\nu} + H_{\text{matter}} \right), \rho(p, t) \right] = \mathcal{C}[\rho, \bar{\rho}]$$

$$C_{\nu_1}^{(r)}[f_j] = \frac{1}{2E_1} \int \frac{d^3p_2}{(2\pi)^3 2E_2} \frac{d^3p_3}{(2\pi)^3 2E_3} \frac{d^3p_4}{(2\pi)^3 2E_4} \\ \times (2\pi)^4 \delta^{(4)}(p_1 + p_2 - p_3 - p_4) S_r \langle |\mathcal{M}_r|^2 \rangle \\ \times F_r(p_1, p_2, p_3, p_4),$$

Collision integral

- ▣ Reduce 5d→2d
- ▣ Gauss-quadrature/Boole's rule

Integrate as ODE

- ▣ xform: $p/T_{\text{cm}}; T_{\text{cm}} \sim 1/a(t)$
- ▣ Fifth-order Runge-Kutta-Cash-Karp

r	Process	$G_F^{-2} S_r \langle \mathcal{M}_r ^2 \rangle$
1	$\nu_i + \nu_i \leftrightarrow \nu_i + \nu_i$	$2^6 (P_1 \cdot P_2)(P_3 \cdot P_4)$
2	$\nu_i + \nu_j \leftrightarrow \nu_i + \nu_j$	$2^5 (P_1 \cdot P_2)(P_3 \cdot P_4)$
3	$\nu_i + \bar{\nu}_i \leftrightarrow \nu_i + \bar{\nu}_i$	$2^7 (P_1 \cdot P_4)(P_2 \cdot P_3)$
4	$\nu_i + \bar{\nu}_j \leftrightarrow \nu_i + \bar{\nu}_j$	$2^5 (P_1 \cdot P_4)(P_2 \cdot P_3)$
5	$\nu_i + \bar{\nu}_i \leftrightarrow \nu_j + \bar{\nu}_j$	$2^5 (P_1 \cdot P_4)(P_2 \cdot P_3)$
6	$\nu_e + e^- \leftrightarrow e^- + \nu_e$	$2^5 [(2 \sin^2 \theta_W + 1)^2 (P_1 \cdot Q_2)(Q_3 \cdot P_4) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_3)(Q_2 \cdot P_4) \\ - 2 \sin^2 \theta_W (2 \sin^2 \theta_W + 1) m_e^2 (P_1 \cdot P_4)]$
7	$\nu_{\mu(\tau)} + e^- \leftrightarrow e^- + \nu_{\mu(\tau)}$	$2^5 [(2 \sin^2 \theta_W - 1)^2 (P_1 \cdot Q_2)(Q_3 \cdot P_4) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_3)(Q_2 \cdot P_4) \\ - 2 \sin^2 \theta_W (2 \sin^2 \theta_W - 1) m_e^2 (P_1 \cdot P_4)]$
8	$\nu_e + e^+ \leftrightarrow e^+ + \nu_e$	$2^5 [(2 \sin^2 \theta_W + 1)^2 (P_1 \cdot Q_3)(Q_2 \cdot P_4) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_2)(Q_3 \cdot P_4) \\ - 2 \sin^2 \theta_W (2 \sin^2 \theta_W + 1) m_e^2 (P_1 \cdot P_4)]$
9	$\nu_{\mu(\tau)} + e^+ \leftrightarrow e^+ + \nu_{\mu(\tau)}$	$2^5 [(2 \sin^2 \theta_W - 1)^2 (P_1 \cdot Q_3)(Q_2 \cdot P_4) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_2)(Q_3 \cdot P_4) \\ - 2 \sin^2 \theta_W (2 \sin^2 \theta_W - 1) m_e^2 (P_1 \cdot P_4)]$
10	$\nu_e + \bar{\nu}_e \leftrightarrow e^- + e^+$	$2^5 [(2 \sin^2 \theta_W + 1)^2 (P_1 \cdot Q_4)(P_2 \cdot Q_3) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_3)(P_2 \cdot Q_4) \\ + 2 \sin^2 \theta_W (2 \sin^2 \theta_W + 1) m_e^2 (P_1 \cdot P_2)]$
11	$\nu_{\mu(\tau)} + \bar{\nu}_{\mu(\tau)} \leftrightarrow e^- + e^+$	$2^5 [(2 \sin^2 \theta_W - 1)^2 (P_1 \cdot Q_4)(P_2 \cdot Q_3) \\ + 4 \sin^4 \theta_W (P_1 \cdot Q_3)(P_2 \cdot Q_4) \\ + 2 \sin^2 \theta_W (2 \sin^2 \theta_W - 1) m_e^2 (P_1 \cdot P_2)]$

Neutrino non-equilibrium results

Employing the time- and momentum-dependent co-rotating frame transformation largely decouples the ultra-high frequency neutrino vacuum oscillations.

$$\tilde{\rho}(p, t) = e^{-itH_V(p, t)} \rho(p, t) e^{+itH_V(p, t)}$$

The figure illustrates the effect of flavor oscillations on the non-equilibrium neutrino distribution functions.

BBN self-consistently coupled to ν transport

BBN/ ν transport

TABLE V. Changes in primordial abundances in BBN for Coulomb and radiative corrections. The first column gives the processes used for a given run. Rows correspond to various corrections as “CC” for Coulomb corrections; “0T” for zero-temperature radiative corrections; “Trans” for neutrino transport calculation with computational parameters as given in Table IV. The notation for the relative changes is the same as in Table IV. Row 4 is our (Q) baseline.

Processes	Y_P	δY_P	$10^5 \times D/H$	$\delta(D/H)$	$10^5 \times {}^3\text{He}/H$	$\delta({}^3\text{He}/H)$	$10^{10} \times {}^7\text{Li}/H$	$\delta({}^7\text{Li}/H)$
None	0.2438	0	2.627	0	1.049	0	4.277	0
CC	0.2474	1.463×10^{-2}	2.647	7.898×10^{-3}	1.052	2.737×10^{-3}	4.317	9.344×10^{-3}
0T	0.2442	1.454×10^{-3}	2.629	7.816×10^{-4}	1.049	0.0	4.281	9.365×10^{-4}
CC, 0T	0.2478	1.613×10^{-2}	2.650	8.719×10^{-3}	1.052	3.021×10^{-3}	4.321	1.030×10^{-2}
Trans	0.2440	4.636×10^{-4}	2.636	3.686×10^{-3}	1.050	1.209×10^{-3}	4.260	-3.916×10^{-3}
CC, 0T, Trans	0.2479	1.644×10^{-2}	2.659	1.236×10^{-2}	1.053	4.209×10^{-3}	4.304	6.231×10^{-3}

$$\delta Y_P = +1.64\%$$

$$\delta(D/H) = +1.24\%$$

*Factor of 20 times
larger than previous
estimates*

- “All”: all antineutrinos/neutrinos on charged leptons and n & p
- Non-linear dependence on included processes
- Observable with next-generation instruments

Conclusion/Outlook

- Include full standard model/quantum theory:
 - ▣ Essential to include *neutrino quantum kinetics*
 - ▣ Flavor oscillations: we've shown for the first time that in the self-consistent approach the neutrino flavors have not 'equilibrated' as BBN is starting
- Solved a long-standing problem of computing the effect of neutrino oscillations in a self-consistent approach using a new technique:
 - ▣ time- and momentum-dependent co-rotating frame
- Other work
 - ▣ testing the validity of the assumption of homogeneity and isotropy in the early universe
 - neutrino flavor oscillations can (do?) affect this
 - ▣ test the affect of neutrino collisions in extreme astro. environments like supernovae