LA-UR-20-29022

Approved for public release; distribution is unlimited.

Title: COVID-19 Modeling for Pandemic Response

Author(s): McCabe, Kirsten J.

Intended for: Directors colloquium

Issued: 2020-11-04

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

GLOBAL DISEASE MODELING
& FORECASTING CENTER

COVID-19 Modeling for Pandemic Response

Kirsten Taylor-McCabe
Global Security Program Office
November 3, 2020

Managed by Triad National Security, LLC for the U.S. Department of Energy's NNSA

Some key facts about the COVID-19 Pandemic

- The Coronavirus disease 2019 (COVID-19) is caused by SARS-CoV-2 and is spreading from person to person mainly through respiratory droplets, aerosols and fomites.
- Spread is more likely when people are in close contact with one another.
- Symptoms may appear 2-14 days after exposure to the virus.
- U.S. number of known cases is doubling about **every three months**.
- ***This is a new virus and we are still learning.***

Cases

9.28M

+74,236

Nov 2, 2020

Deaths

231K

+427

What we need to know

. ...science for decision making

Scientist at LANL are involved in the global effort to understand this outbreak.

LANLs Core Team Disciplines

Joint DOE Effort for Pandemic Modeling and Analysis Overview

ORNL, ANL, SNL, LANL

- Recognized potentially very high values of *effective* reproduction number (R_{eff}) very early in pandemic.
- Careful modeling on NY, Seattle, Wuhan, and Italy enabled determinants of force of infection to be dis-entangled and thus translated from scenario to scenario.
- Provided early posting of statistical models, translated to hospital and ICU loadings.
- Provided detailed comparisons of impact of bars and schools re-opening on COVID spread.
- Recognition of Descartes Labs cell phone data in capturing by-county variation in R_{eff} .
- Quantified contact tracing as 40-60% effective when 300 tracers per 2M population were deployed, as independent mitigation to masks / social distancing / closures.
- Developed systems-engineering model relating testing turnaround time, sensitivity & specificity, contact tracing & isolation to consequence.

Modeling provides a framework to evaluate forecasts, potential actions, and specific scenarios

- Outbreak surveillance, Hot Spots, Mobility factors/impacts, Transmission links, School re-opening, Non-congregate housing needs, Short-term hospital resource needs, vaccine distribution strategies, impacts of mitigations/contact tracing and more.

Models Improve Understanding and Support Pandemic Response Decisions: Forecasting

COVID-19 Cases and Deaths Forecasts

- Forecasts up to six weeks ahead for all US states and many countries. Forecasts are probabilistic, providing a range of possible future outbreak trajectories.

- CDC-cited model

6-Week Forecast of Confirmed Cases for New Mexico Based on Data as of 2020-10-25			
Week	Best Case (5th Percentile)	Middle Case (50th Percentile)	Worst Case (95th Percentile) [^]
2020-10-25		41,863*	
2020-11-01	44,864	47,190	50,410
2020-11-08	46,922	51,939	59,788
2020-11-15	48,553	56,072	69,805
2020-11-22	49,656	59,651	79,804
2020-11-29	50,470	62,610	89,660
2020-12-06	51,107	65,313	99,499

*Last reported confirmed cases count
[^]Closest-matching scenario

- New Mexico Department of Health Operational Support

- <https://covid-19.bsvgateway.org>

Models Improve Understanding and Support Pandemic Response Decisions: Forecasting

- Infectious disease forecasting that blends the structure of epidemiological models with the flexibility of machine learning.

$$\hat{y} = f(X, \beta)$$

- Model has been successfully applied to real-time influenza and COVID-19 forecasting. Can be extended to many diseases and geographies.

So what?

We are on track to exceed ICU beds by November 8. This is using the updated LANL forecasting model COFFEE.

Support Pandemic Response Decision

Non-Congregate Shelter Forecast Estimates to facilitate planning action.

The daily number of cases is expected to range between 80 and 90 for the middle case scenario in the next few weeks

COVID-19 across New Mexico

A 7-day moving window comparison
Oct 26, 2020

So what?

- MOST New Mexicans live in a county with currently accelerating growth and high per-capita case counts
- No counties are decelerating right now

Manore, Del Valle & Team

EpiGrid: Impacts of Assumptions and Interventions

EpiGrid

- Readily parameterized medium-grained epidemiological model that links geography, disease and epidemic progression, and mitigations.
- Model is enabling timely and accurate decision support.
- Has been utilized in over 200 scenarios by DoD and spans 12+ diseases.

Improving disease emergence situational awareness & preparation

- A direct means of incorporating crucial information such as mobility as an independent estimator of the force of infection.

- “What if” planning - such as school reopening and understanding and reconciling differences between administrative districts (counties)

<https://cvm modeling.nmhealth.org/medical-advisory-team/modeling-updates/>

Impacts of Workforce Reopening's

EpiCast

- A fine-grained hybrid-agent epidemic model with diurnal agent travel and contagion compartments that allow the analysis of the importance of contact-networks, travel, and detailed intervention strategies for the control of outbreaks and epidemics
- 2000-person communities in 65,433 census tracts
- Explicit model of geography, demographics (i.e., age), worker/household/school/ community contacts, and mitigations
- Captures workforce by 3-digit NAICS

LANL-specific modeling support

So what?

Relaxation of interventions can result in a second wave of infection

Spatial Differentiation as a Function of Demographics and Public Health Orders

Counties clustered based on similar income, population size, age, and education

There is a correlation between cases and bar and restaurant public health orders – the number of cases increased as bar and restaurant orders were relaxed

COVID-19 Studies on Drivers of Pandemic Outcomes

Modeling epidemics for decision support with infrastructure analysis (MEDIAN)

- Understand interdependencies between critical infrastructures, contact tracing, testing and diagnostics to uncertainty in the primary drivers of pandemic outcomes and mitigation impacts.
- Contact tracing and testing is vital but the acceleration of cases is still driven by isolation and quarantine.

Mining Nontraditional Data (e.g., Google, Twitter) to Measure Human Behavior

- Analyze trends geospatially and over time using natural language processing (e.g., sentiment analysis, topic modeling) and machine learning.
- Understand evolution of important words to observe important shifts (e.g., patterns in tweets suggesting vaccine adverse beliefs) that can inform communication strategies.

COVID-19 Impacts on Behavioral Health

Reactions and Behavioral Health Symptoms in Disasters

National disasters such as pandemics impact mental and behavioral health, including depression, anxiety, domestic violence, and substance abuse.

These impacts often occur in phases (see top left).

a. Static vulnerability index

b. Vulnerability index, week 22

We are analyzing state data about mental health along with socio-economic factors to estimate vulnerability and help determine future needs.

Vulnerability index based on socio-demographics, changing in time with COVID-19 cases

During late June, 40% of U.S. adults reported struggling with mental health or substance use^{*}

^{*}Based on a survey of U.S. adults aged ≥18 years during June 24-30, 2020
[†]In the 30 days prior to survey

For stress and coping strategies: bit.ly/dailylifecoping

CDC.GOV

bit.ly/MMWR81320

MMWR

We are investing LDRD funds in research on COVID transmission and outbreaks in the state

Approach

- SARS-CoV-2 mutations reflect a tree of relatedness among cases, and hence transmission links

Methods

- Tree-based analyses reveal when the virus moved into/out of NM, and hence NM transmission clusters

We have found we can:

- Identify many transmission links
- Suggest additional cases likely to be part of outbreaks
- Show that outbreaks often result from multiple introductions (rather than having a single source)

Cases (red circles) are mixed among cases from elsewhere, showing many introductions of the virus into the state

Inset: Cluster sub-tree shows transmission link between two designated outbreaks

LANL High Performance Computing: Why is SARS-CoV-2 more invasive than SARS-CoV?

Forming a Tighter Grip
Revealed by extra long MD simulations

Human ACE-2

SARS-CoV-2

- The receptor binding domain of SARS-CoV-2 has evolved to have a stronger grip on Human receptors ACE2 than SARS-CoV.
- Loop with F486 on CoV-2 forms a sticky anchor and maintains interaction with ACE-2
- Loop with L486 on CoV dislodges easily
- Structural modeling allowed LANL to confirm a utilizing a SARS-like model rather than a MERS-like model in early Jan 2020

Tung and Team and Ngo, Jha and Team

The D614G mutation

SARS-CoV-2 Spike is the protein that mediates virus entry into cells **and it's the prime target of COVID vaccines.**

A single mutation in Spike, **D614** to **G614**, has become the dominant form of the virus in the pandemic.

The **G614** form is more infectious.

The **G614** form is associated with higher levels of viral RNA in the upper respiratory tract of infected people.

The **G614** was not associated with increased hospitalization.

Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus.

Korber B, Cell. 2020 Aug 20;182(4):812-827.e19.

Korber and Team

The D614G mutation

By mid-summer 2020, the G614 form had become the globally dominant form of the virus

Oct. 27, 2020 data : 10 day windows, from SARS-CoV-2 sequences in GISAID

LANL built computation tools to geographically/temporally track mutations in the COVID-19 pandemic
cov.lanl.gov

Fueled by **GISAID**

Virus-Containing Droplet Plume Modeling

DOE Office of Science: Viral Fate
& Transport for Covid-19

How far can virus-filled droplets travel?

- LANL's QUIC plume model is used to account for dispersion and deposition of droplets accounting for air currents, turbulence, gravitational settling & droplet evaporation.
- With a 0.5 m/s air current and medium to low humidity, 100-um droplets can travel **10+ meters** before settling out onto surfaces

Is outdoor dining safe?

1-person
breathing
in center
of each
courtyard

- For deeper courtyards, QUIC dispersal calculations suggest weaker ventilation, leading to airborne concentrations that are similar to those found in fast-food indoor restaurants.

Conclusions

- We do not know everything, but leveraging international networks and community tools will tell us what needs to be done to beat this outbreak.
- Decision makers need facts supported by science and data.
- A layered approach is the best approach.
- LANL has used multiple models to study the impacts of different planning assumptions and interventions (e.g., reopening, physical distancing, vaccination, virus properties, supply chain, hospitalizations) as well as to understand the biology and movement of the virus.
- We need to maintain capabilities for the next disease outbreak.