LA-UR-21-20119

Approved for public release; distribution is unlimited.

Title: IC project: Exploring Planet-disk Interaction in Dusty Protoplanetary
Disks and Their Observational Implications

Author(s): Li, Yaping

Intended for: Report

Issued: 2021-01-07

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

IC project: Exploring Planet-disk Interaction in Dusty Protoplanetary Disks and Their Observational Implications

PI: Yaping Li

Managed by Triad National Security, LLC for the U.S. Department of Energy's NNSA

Planet-induced Vortices with Dust Coagulation (**low α , high m_p**)

- Dust size is quite non-uniform within vortex region
- Dust size growth facilitates the increase of dust/gas ratio.
- Vortex lifetime can be significantly impacted by dust feedback when coagulation included.

YPL et al. 2020a

Planet-induced Vortices with Dust Coagulation

Clumpy at longer wavelength, while rings at (sub)mm band for coagulation models consistent with LkCa 15 (Isella+2014; testing with new JVLA observations). YPL et al. 2020a

Multi-ring Morphology with Dust Coagulation (**low α , low m_p**)

- If the planet doesn't open a gap quickly enough (thick disk or low planet mass), an inner ring disappears due to dust coagulation and subsequent radial drift.

Laune et al. 2020

Giant planet accretion constrained by tidal barrier

- In weakly α disks, protoplanets' accretion rate steeply decreases with $M_p \rightarrow$ asymptotic $M_p \sim M_J$, consistent with Kepler observations.
- $10 M_J$ fall-off for known exoplanets $\rightarrow$ natal disks have $\alpha \sim 10^{-3}$ and $H/R \sim 0.03 - 0.05$.

Outward migration of giant planets

At the edges of a deep gap, low order ILR dominates over OLR.

- a steep Σ_g profile significantly suppresses of migration rate or even reversal of the migration direction: a new mechanism for retention of long-period giants.

Chen et al. 2020b