LA-UR-21-24321

Approved for public release; distribution is unlimited.

Title: (U) ATCC-10 LSCI: Active learning for atomic plutonium potentials.

Author(s): Nebgen, Benjamin Tyler
Smith, Justin Steven
Barros, Kipton Marcos

Intended for: Report

Issued: 2021-05-04

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

(U) ATCC-10 LSCI: Active learning for atomic plutonium potentials.

Ben Nebgen, Justin Smith,
Kipton Barros

5/5/2021

Managed by Triad National Security, LLC for the U.S. Department of Energy's NNSA

Scale Bridging for Materials Properties

Active Learning for Dataset Generation

An Active Learning system was developed for the automated exploration of phase space. This system builds a training dataset for an Machine Learned interatomic potential with little to no human intervention.

Using Query by Committee (a UQ method), we intelligently select configurations to augment the ML training set intelligently.

Smith, J. S.; Nebgen, B.; Mathew, N.; Chen, J.; Lubbers, N.; Burakovsky, L.; Tretiak, S.; Nam, H. A.; Germann, T.; Fensin, S.; Barros, K., Automated discovery of a robust interatomic potential for aluminum. *Nature Comm.* **2021**, 12.

Aluminum Shock

This methodology was used to build a ML interatomic potential for Al during Sierra open.

The potential was used to perform a 1.2 M atom shock simulation. Forces were verified by DFT.

Potentials Developed in ATCC-10: Au, Ag, and Pu

Au, Ag, and Pu ML potentials developed during ATCC-10 show good agreement with Density Functional Theory (DFT) properties.

Ag-ML potential melting point computed to be ~ 980 °C, experimental is 962 °C (very good agreement).

Pu in good agreement with DFT properties, ATCC-11 will further refine with higher accuracy methods.

Outlook and Lessons Learned

- **ML potentials represent about a x10,000,000 speedup over Quantum Mechanical Methods**
 - ML potential (ANI+LAMMPS) can run on 1280 Sierra nodes efficiently.
- **Sierra is an incredibly powerful resource.**
 - Development of Pu potential is computationally ~100 times more difficult than development of Au/Ag. **Only possible on a computer like Sierra.**
 - Distributed GPU architecture (Sierra) works extremely well for both Active Learning and domain decomposition (small regions of space on each GPU node).
- **Difficult to estimate time until completed and validated Pu-alloy potential. Between end of 2021 to end of 2022.**
 - Pu requires transfer learning. Collaboration with Gabi Kotliar (Rutgers).
 - Smith et al “Approaching coupled cluster accuracy with a general-purpose neural network potential through transfer learning” *Nat. Comm.* 2019, 10, 2903.
- **Future Work**
 - Incorporating experimental data in ML potential training (ongoing LDRD-DR project building capability).